



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.

We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>





[Redacted]

[Redacted]

FERNS OF BRITISH INDIA.



ADIANTUM HISPIDULUM. (Sw.)



191 150.

A.S. CAPTAIN 1257419

ADIANTUM

HANDBOOK
TO THE
FERNS OF BRITISH INDIA,
CEYLON AND THE MALAY PENINSULA.

BY

COL. R. H. BEDDOME, F.L.S.,
Late Conservator of Forests, Madras.

*Author of "The Ferns of British India," "The Ferns of Southern India,"
"The Flora Sylvatica of Southern India," &c., &c.*

WITH THREE HUNDRED ILLUSTRATIONS.

CALCUTTA :

THACKER, SPINK AND CO.;

BOMBAY : THACKER & CO., LIM.; MADRAS : HIGGINGBOTHAM & CO.;

LONDON : W. THACKER & CO.

1883.

191 ; 166 .



LONDON
WILLIAM RIDER AND SON, PRINTERS.
BARTHOLOMEW CLOSE.

P R E F A C E.

THIS handbook is a digest of the information on Indian Ferns contained in Sir W. Hooker's "*Species Filicum*," the "*Synopsis Filicum*," Mr. J. Smith's "*Historia Filicum*," also of Mr. Clarke's "*Ferns of Northern India*," and of the author's works, "*The Ferns of Southern India*" and "*The Ferns of British India*."

The author's former works were compiled in India, without access to libraries or to any herbarium with typical specimens. He has now had the advantage of going thoroughly through the Wallichian ferns at the Linnæan, the large collections at Kew and in the British Museum, and of comparing his own collections with typical forms; he is, besides, indebted for much assistance to Messrs. Baker and Clarke, especially in the determination of all critical forms, so that he has been able, he believes, to clear up many doubts and to correct many errors.

The generic nomenclature is, with few exceptions, that of the "Synopsis Filicum," but the sub-genera have been raised to the rank of genera ; this avoids the inconvenience of double generic names, and may be admitted on this score even by those who do not consider the wide difference in habit of such genera as *Phegopteris*, *Polypodium*, *Pleopeltis*, and *Drynaria* of any value generically, and who base their genera on the sori alone, without reference to habit, venation, or vernation.

The geographical limits of the work comprise the whole of British India, Ceylon and the Malay Peninsula.

WEST HILL, PUTNEY,
May, 1833.

TABLE OF THE
ORDERS, TRIBES, AND GENERA OF FERNS
AS FOLLOWED IN THIS WORK.

ORDER FILICES.

SUB-ORDER I.—GLEICHENIACEÆ.

	GENUS.	PAGE.
Capsules opening vertically, surrounded by a broad transverse complete ring	1 GLEICHENIA	2

SUB-ORDER II.—POLYPODIACEÆ.

Capsules surrounded by a jointed, vertical and elastic ring.

A. INVOLUCRATÆ. *Sori with an indusium, except in Alsophila.*

TRIBE I.—CYATHEÆ.

Sori globose, capsules sessile or stalked, often on a raised receptacle. Indusium (wanting in Alsophila) inferior, including the sorus, lateral and resembling a scale, or cup-shaped, often when young enveloping the sorus, eventually opening at the summit, or breaking down with a more or less regular margin, caudex generally arborescent.

Receptacle elevated, indusium globose, inferior, quite covering the sorus when young	2 CYATHEA	5
Indusium a cup-like scale below the sorus, but never quite covering it, veins free	3 AMPHICOSMIA	8
Indusium none	4 ALSOPHILA	11
Receptacles not elevated, indusium inferior globose .	5 DIACALPE	13
Indusium umbrella-shaped, six-lobed	6 MATONIA	19

	GENUS.	PAGE.
TRIBE II.—DICKSONIÆ.		
Sori globose, indusium inferior, subglobose, free, sometimes covering the whole sorus, closed, at length bursting irregularly, more frequently cup-shaped entire or with two lips.		
Fertile fronds contracted with revolute margins, indusium hemispherical, very fugacious or wanting, veins free	7 STRUTHIOPTERIS	20
Indusium thin, subglobose, open at the top and lacinate	8 WOODSIA	20
Indusium medial on a vein, thick globose pedicellate, bursting into two lips	9 PERANEMA	22
Indusium apical on a vein, two-valved	10 CIBOTIUM	24
Indusium apical on a vein, cup-shaped	11 DENNSTÆDTIA	24
TRIBE III.—HYMENOPHYLLÆ.		
Filmy ferns, indusium inferior, apical on a vein, two-lipped or tubular.		
Indusium more or less two-lipped	12 HYMENOPHYLLUM	28
Indusium tubular, the mouth truncate or slightly two-lipped	13 TRICHOMANES	36
TRIBE IV.—DAVALLIÆ.		
Indusium squamiform, suborbicular or tubular, open at the apex, fastened broadly at the base, and sometimes at the sides.		
Indusium apical on a vein, suborbicular coriaceous, attached by a broad base, the sides free	14 HUMATA	46
Indusium as in Humata, but thin and smaller and narrower	15 LEUCOSTEGIA	48
Indusium a subcylindric cyst formed of the substance of the frond, apical on the veins	16 PROSAPTIA	56
Indusium apical, tubular, attached by the base and sides	17 DAVALLIA	58
Indusium apical, half cup-shaped, attached by the base and sides	18 MICROLEPIA	62

TABLE OF THE GENERA OF FERNS.

ix

	GENUS.	PAGE.
Indusium apical, compound, suborbicular, only open at the top	19 STENOLOMA	. 68
Indusium medial on a vein membranaceous, hood-like	20 CYSTOPTERIS	. 70

TRIBE V.—LINDSAYEÆ.

Indusium apical on the veins, in a continuous or interrupted line, the outer valve being the margin of the frond, the inner membranaceous.

Veins free	21 LINDSAYA .	. 72
Veins anastomosing	22 SCHIZOLOMA	. 77

TRIBE VI.—PTERIDEÆ.

Indusium oblong or linear, formed of the more or less changed and reflexed margin of the frond, opening inwardly.

Indusium globose to linear usually many and distinct, sometimes confluent and continuous, bearing the capsules on its under side, veins free	23 ADIANTUM	. 82
Indusium roundish and distinct, or more or less confluent but not continuous, capsules on the frond	24 CHEILANTHES	. 88
Indusium as in Pteris, the ultimate segments of fronds small and very narrow	25 ONYCHIIUM	. 95
As in Pellæa, but fronds dimorphic	26 CRYPTOGRAMME	98
Indusium quite continuous, sori at first dot-like, but soon running into a line	27 PELLÆA	98
Indusium quite continuous, sori linear continuous, occupying a slender filiform receptacle in the axis of the indusium, veins free	28 PTERIS	104
As in Pteris, but lowest veins meeting and forming an arch	29 CAMPTERIA	. 116
As in Pteris, but veins all anastomosing without free included veinlets	30 DORYOPTERIS	. 120
As in Pteris, but veins all anastomosing with some free included veinlets	31 LITOBROCHIA	. 120

TABLE OF THE GENERA OF FERNS.

	GENUS.	PAGE.
Anomalous	32 CERATOPTERIS	123
Indusium continuous, sori linear, continuous, covering all or nearly all the space between the midrib and margin, fronds dimorphous, capsule ring vertical	33 LOMARIA	125
As in Lomaria, but capsule ring oblique	34 PLAGIOGYRIA	127

TRIBE VII.—BLECHNEÆ.

Indusium linear or oblong parallel with the midrib and opening towards it, not near the margin.

Indusium membranaceous, distinct from the margin of the frond, parallel with and usually contiguous to the midrib, veins free	35 BLECHNUM	130
As in Blechnum, only veins anastomosing	36 BLECHNIDIUM	133
Indusium linear-oblong, thick, in single rows parallel with and near the midrib, sori sunk in cavities	37 WOODWARDIA	135
Indusium lunate, thin, in one or more rows, parallel with and near the midrib, sori not sunk	38 DOODIA	135

TRIBE VIII.—ASPLENIEÆ.

Indusium linear or oblong or horseshoe-shaped, opening towards the midrib, sometimes double, sori attached to the veins.

Indusium linear, single, veins free, their apices combined by a transverse marginal vein	39 THAMNOPTERIS	137
Indusium linear or oblong single, veins free	40 ASPLENIUM	141
As in Asplenium, but indusium more or less curved	41 ATHYRIUM	161
As in Asplenium, but indusium double	42 DIPLAZIUM	174
As in Diplazium, but veins anastomosing	43 ANISOGONIUM	191
As in Asplenium, but veins anastomosing obliquely only towards margin of frond	44 HEMIDICTYUM	194
Indusium sausage-shaped, veins anastomosing and forming elongated areoles	45 ALLANTODIA	195
Indusium linear elongated, submarginal, fronds fan-like	46 ACTINIOPTERIS	197

TABLE OF THE GENERA OF FERNS.

xi

	GENUS.	PAGE.
TRIBE IX.—ASPIDIEÆ.		
Indusium superior, elliptical, subglobose or reniform fixed either by the centre or a sinus.		
Indusium elliptical, emarginate at the base, attached longitudinally to the linear receptacle, veins free	47 DIDYMOCHLÆNA	199
Indusium of Didymochlæna habit and venation of Nephrodium	48 MESOCHLÆNA	199
Indusium orbicular, veins free	49 POLYSTICHUM	201
Indusium orbicular, veins generally anastomosing acutely with free veinlets from their junction	50 CYRTOMIUM	211
Indusium peltate orbicular, or reniform, veins copiously anastomosing with free included veinlets	51 ASPIDIUM	212
Indusium reniform, veins only slightly anastomosing	52 PLEOCNEMIA	223
Indusium reniform, veins free	53 LASTREA	228
Indusium reniform, one or more of the lower veinlets anastomosing with the corresponding one of the next group	54 NEPHRODIUM	267
As in Lastrea, but always simply pinnate, the pinnæ articulated and furnished with white dots above	55 NEPHROLEPIS	282
As in Lastrea, but wide creeping and scandent with jointed stems and entire fronds	56 OLEANDRA	285
B. EXINVOLUCRATÆ.— <i>Sori without an indusium.</i>		
TRIBE X.—POLYPODIEÆ.		
Sori on the back of the lobes, round or rarely somewhat oblong.		
Stipes continuous with the caudex, habit of Lastrea, veins free	57 PHEGOPTERIS	288
Stipes continuous with caudex, venation and habit of Nephrodium	58 GONIOPTERIS	296
Stipes continuous with caudex, venation and habit of Aspidium	59 DICTYOPTERIS	298
Stipes articulated with caudex, veins free	60 POLYPODIUM	302
Veins forming regular areoles, the lower one or more with a free included soriferous veinlet	61 GONIOPHLEBIUM	316

	GENUS.	PAGE.
Areoles copious, each with two or more sori, fronds always simple and woolly beneath	62 NIPHOBOLUS	. 324
Fronds flabellate in two halves, veins copiously anas- tomosing	63 DIPTERIS	. 334
Fronds either with the base oak-leaf-like, or with separate sterile oak-leaf-like small fronds	64 DRYNARIA.	. 338
Fronds various, veins copiously anastomosing with free included veinlets	65 PLEOPELTIS	. 344

TRIBE XI.—GRAMMITIDEÆ.

Sori on the back of the lobes more than twice as long as broad, usually linear.

As in Cheilanthes, but without an indusium, sori though oblong or roundish at first, soon confluent into a continuous line	66 NOTHOLÆNA	. 373
Sori linear, close to midrib on both sides, fronds grass-like	67 MONOGRAMME	. 375
Sori linear or linear-oblong, otherwise as in Phegop- teris	68 LEPTOGRAMME	. 377
As in Nephrodium, but sori linear	69 STEGNOGRAMME.	380
Habit of Cheilanthes, sori linear, veins free	70 GYMNOGRAMME.	380
Habit of Polypodium, sori linear, veins generally united by transverse veinlets near margin	71 SYNGRAMME	. 384
As in Pleopeltis, with copiously anastomosing veins, sori linear, stipes articulate with caudex	72 SELLIGUEA	. 389
As in Selliguela, but stipes not articulate and fronds leathery	73 LOXOGRAMME	. 392
Anomalous, veins arcuately anastomosing at the base, forming costal areoles	74 BRAINEA	. 395
As in Goniopteris, but sori elongated or confluent .	75 MENISCIMUM	. 397
Sori carried along the veins imperfectly articulated, fronds simple	76 ANTROPHYUM	. 401
Grass-like ferns, sori in continuous marginal lines, veins free	77 VITTARIA	. 404

TABLE OF THE GENERA OF FERNS.

xiii

	GENUS.	PAGE.
Sori in a continuous linear, or interrupted central or submarginal line, veins reticulated, fronds pinnate .	78 TÆNITIS .	410
As in Tænitis, but fronds simple and dimorphous .	79 DRYMOGLOSSUM .	410
Veins copiously reticulated and completely soriferous.	80 HEMIONITIS .	413

TRIBE XII.—ACROSTICHEÆ.

Sori not confined to the veins, but spread entirely over the frond beneath.

Fronds simple, veins free	81 ELAPHOGLOSSUM .	416
Fronds pinnate, Lomaria-like, veins all simple or sometimes arcuate at the base ; scandent ferns .	82 STENOCHLÆNA .	421
Fronds pinnate, not Lomaria-like, veins all simple ; not scandent	83 POLYBOTRYA .	424
Similar to Elaphoglossum, but with a connecting vein running round the margin	84 ACONIOPTERIS .	427
Veins copiously anastomosing with free included veinlets	85 GYMNOPTERIS .	429
Veins copiously anastomosing without free included veinlets. Sea-coast fern	86 ACROSTICHUM .	440
Drynaria-like fronds, pinnæ articulated, but stipe adherent to caudex	87 PHOTINOPTERIS .	442
Fronds with stag's-horn-like divisions, vernation articulate, sori in large patches only on the fertile fronds	88 PLATYCERIUM .	444

SUB-ORDER III.—OSMUNDACEÆ.

Capsules two-valved, opening across the apex, furnished with a short horizontal ring

89 OSMUNDA .	447
--------------	-----

SUB-ORDER IV.—SCHIZÆACEÆ.

Capsule two-valved, opening down the side, crowned by a complete operculiform ring.

Capsules sessile in 2-4 rows on one side of close spikes

90 SCHIZÆA .	452
--------------	-----

	GENUS.	PAGE
Capsules smooth, very abundant in a branched panicle	91 ANEMIA .	453
Capsules solitary in the axils of large imbricated clasping involucres ; scandent	92 LYGODIUM .	453

SUB-ORDER V.—MARATTIACEÆ.


Capsule opening by a slit down one side or a pore at the apex, without a ring.

Capsules sessile, very close together, but not concrete	93 ANGIOPTERIS .	458
Capsules concrete, in boat-shaped spore-cases . . .	94 MARATTIA .	460
Capsules concrete in raised circular masses which are hollow in the centre	95 KAULFUSSIA .	462

SUB-ORDER VI.—OPHIOGLOSSACEÆ.


Capsules deeply two-valved, opening down the side nearly to the base, without a ring.

Capsules sessile in two rows on a narrow close spike .	96 OPHIOGLOSSUM .	462
Capsules in small crested clusters forming a loose spike	97 HELMINTHO-STACHYS .	467
Capsules in two rows on the face of spikes which form a compound panicle	98 BOTRYCHIUM .	469



J. B. GUTHRIE, D. C. L. 1884

GLEICHENIA GLAUCA. (Hook.)



CYATHEA HOOKERI. (Thunb.)

HANDBOOK TO THE FERNS OF BRITISH INDIA, CEYLON, AND THE MALAY PENINSULA.



ORDER FILICES.

CAPSULES (sporangia) one-celled, usually membranaceous and dorsal (on the back of the frond), or marginal, encircled by a complete or incomplete, jointed, elastic ring, collected in clusters (sori) of a definite but varied form, with or without a covering (indusium) or panicled or spicate, rarely laxly scattered; or the capsules have the ring obsolete, or none, or the ring is confined to the apex of the capsule, where it forms a longitudinally striated crown, bursting vertically; or the capsules are sunk in a many-celled, fleshy or corky receptacle of varied forms, opening by pores or clefts on the upper surface; spores minute, various in form; vernation generally circinate, rarely erect.

SUB-ORDER I.—GLEICHENIACEÆ.

Sori dorsal; of few (2–10) capsules; capsules opening vertically surrounded by a broad, transverse, complete ring; indusium none; caudex generally creeping; stipes often dichotomous; fronds rigid, generally large, and dichotomously branched, frequently bearing axillary buds; vernation circinate.

GENUS I.—GLEICHENIA. (*Smith.*)

(In honour of Gleichen, a German botanist).

Sori of few sessile capsules, situated on a lower exterior veinlet ; caudex generally creeping ; frond rarely unbranched, generally dichotomously divided, often proliferous from the axils of the forks ; pinnæ deeply pinnatifid, with the segments small and concave, sub-orbicular, or pectinate with elongated lobes.

§ Sori solitary at the apex of a veinlet.

1. GLEICHENIA CIRCINATA. (*Sw.*) Fronds dichotomous, divaricated, lobes of the pinnules bead-like, ovate, or subrotund, the margins slightly recurved ; capsules 3-4 ; branches and rachis glabrous, or more or less scaly. *Bedd. F. B. I. t.* 177, as *semivestita* Labill. *Hook. Sp. Fil. p.* 11.

Malacca.

(Also in Australia, Tasmania, New Zealand, and New Caledonia.)


§ Sori near the middle, or at the forking, of the veinlets.

2. GLEICHENIA GLAUCA. (*Hook.*) A large straggling fern many feet long ; stipes stout forked ; primary pinnæ opposite, secondary pinnæ alternate, close, 6-8 inches long by 1-2 broad, lanceolate, acuminate, pinnatifid nearly to the rachis, the segments or pinnules oblong-lanceolate, but obtuse, more or less glaucous beneath, glabrous or tomentose, and paleaceous on the stipes and rachis ; capsules 3-5, often mixed with hairs. *Bedd. F. B. I. t.* 30, as *gigantea* Wallich. *G. longissima*, *Bl. Hook. Sp. Fil. p.* 12. *G. gigantea*, *Wall. Cat.* *Polypodium glaucum*, *Thunb. Fl. Jap.* 338, which is the oldest specific name.

Bhotan and Sikkim, alt. 4,500-7,500 feet. Very common about Darjeeling, Khasya, 3,500-5,000. Common in Nepal, Malay Peninsula.

(Also in China, Japan, the Malay Islands, Sandwich Islands, Tropical Australia, Polynesia, West Indies, and Tropical America.)

3. GLEICHENIA NORRISII. (*Mett.*) Branches of the frond one-jugate, oblong-lanceolate, 2-3 feet long, pinnæ lanceolate, the lower distinctly stalked, spreading, 6-9 inches long, cut down to a narrow



GLEICHENIA LINEARIS (Burm.)

wing into close entire ligulate blunt segments, upper gradually smaller, reflexed; texture subcoriaceous, both surfaces glabrous, the lower green or glaucous; veinlets once forked; sori medial, 12–20 to a segment. *Bedd. F. B. I. t.* 346. *Hook. Sp. Fil. p.* 449.

Malacca.

4. *GLEICHENIA LINEARIS* (*Burm. under Polypodium*). Stipes zigzag, repeatedly di- or trichotomous, the ultimate branches bearing a pair of forked pinnæ; a distinct pair of pinnæ also arises from the base of the forked branches, pinnules usually glaucous beneath, usually entire with recurved margins, texture hard, veinlets usually three-branched from the base, the centre one being often forked or pinnate. *Bedd. F. S. I. t.* 74, as *dichotoma*. *Gl. dichotoma*, *Willd. Hook. Sp. Fil. p.* 15. *Polypodium lineare*, *Burm. Fil. Ind.* 235, *t.* 67, fig. 2, oldest specific name. This fern has long been known under the name of *dichotoma*, which, however, must be dropped, as there is an older specific name.

Mountains of southern India and Ceylon, up to 6,000 feet, Sikkim, Bhotan, Nepal, Kumaon, Khasya, &c., up to 5,000 feet. Malay Peninsula.

(Also in Japan, Tropical Australia, America, Polynesia)

SUB-ORDER II.—POLYPODIACEÆ.

Sori dorsal or marginal, subglobose, of many capsules, with or without an indusium, usually pedicellate, more or less completely surrounded with a jointed vertical and elastic ring, and bursting transversely (except in *Hymenophyllæ*).

A.—*INVOLUCRATÆ*.—Sori furnished with an indusium (except in *Alsophila*), Tribes *Cyatheæ* to *Aspidiæ*.

TRIBE I.—CYATHEÆ.

Sori dorsal, globose, often at or near the forking of a vein; capsules numerous, often very compact sessile or stalked, generally on an elevated receptacle, often mixed with hairs, obovate usually with a broad, vertical, or sub-oblique elastic ring; indusium (none in

Alsophila) inferior, including the sorus, lateral and resembling a scale on the under side of the sorus, or cup-shaped, often when young enveloping the sorus, eventually opening at the summit, or breaking down with a more or less regular margin ; caudex very often arborescent.

GENUS II.—CYATHEA. (*Smith.*)

(*Kyathos*, a cup—the form of indusium.)

Sori on a vein, or in the axil of the forking of a vein, receptacle elevated, globose, or elongated ; indusium globose inferior, covering the whole sorus, afterwards breaking at the summit, and forming a more or less persistent cup, even or irregular, at the margin ; generally arborescent ; stipes often aculeated ; fronds simple, pinnate, or decompoundly pinnate.


* *Fronds undivided.*

1. CYATHEA SINUATA. (*Hook. and Grev.*) Caudex erect, 2–4 ft. long, about 1 inch in diameter ; frond simple, 2–3 ft. long, 12½ inches wide, elongate-lanceolate, sinuate, acuminate, tapering at the base ; veins pinnate ; veinlets soriferous near middle. *Bedd. F. S. I. t. 259 ; Hook. Sp. Fil. p. 16.*

Ceylon, in the Singhe-Rajah Forest.

** *Fronds pinnate.*

2. CYATHEA BRUNONIS. (*Wall.*) Stipes 12 feet long ; fronds 2–3 ft long, pinnate glossy ; pinnæ 6–14 inches long, alternate, between membranous and coriaceous, oblong-lanceolate, acuminate with a long narrow point obliquely truncated at the base and shortly



No. 2.

CYATHEA SINUATA. (*Hook.*)

petiolate, sinuate-crenate; veins three-branched or pinnate, the veinlets occasionally anastomosing amongst themselves, or with the next group; sori medial on the veinlets; indusium very persistent. *Bedd. F. B. I. t. 87; Hook. Sp. Fil. p. 16.*

Penang and Malacca.

(Also in the Malay Islands.)


3. *CYATHEA HOOKERI*. (*Thw.*) Small, but with a trunk-like caudex $1\frac{1}{2}$ inch thick; stipes short black, muricated at the base and sub-paleaceous; fronds coriaceous-membranaceous, 2-3 feet long, 4-5 inches wide, elongate-lanceolate, acuminate, pinnate pinnatifid at the apex; pinnæ from a broad base, which is more or less auricled, lanceolate acuminate, sessile or sub-sessile, coarsely dentate-pinnatifid, more or less entire towards the apex and base, and the lower ones gradually diminishing in size and obtuse at their apex; veins pinnate; sori medial on the veinlets; indusium soon breaking up and becoming cup-like. *Bedd. F. B. I. t. 260. Thw. En. Pl. Zy. p. 396. Hook. Sp. Fil. p. 16.*

Ceylon, in the Singhe-Rajah Forest.

*** *Fronds decomponently pinnate.*

4. *CYATHEA SPINULOSA*. (*Wall.*) A tall tree fern; stipes and main rachis beneath, strongly aculeate, dark purple; fronds glabrous, tripinnatifid; main rachis and rachis of pinnules ferruginous above; rachis of pinnules and main vein of segments scaly below, but the latter glabrous above; segments falcate-oblong acute, serrulate, the margin more or less recurved; veinlets once-forked, or more rarely three-branched; sori copious near the costules or main veins; indusium completely covering the sorus when young, soon breaking irregularly. *Bedd. F. S. I. t. 57. Hook. Syn. Fil. p. 23.*

The Wynad at 3,000 feet elevation, North and South Canara Coorg, Jeypore Hills (Vizag), Nepal, Jaintea Hills. The Wynad specimens are in every way identical with those from Northern India collected by Wallich, and Mr. Clarke is in error in stating that the South Indian plant is a *Hemitelea*, as on comparing with him the specimens he had examined at Kew I found that they were "*Alsophila latebrosa*," and that this *Cyathea* was not represented



CYATHEA SPINULOSA. (Wall.)

from S. India in the *Cyathea* bundles at Kew, although there was a specimen of it from Dalzell elsewhere, labelled "*Lastrea also-philoides*," and a specimen from Canara, mixed with *Hemitelea*. I find no 3-branched veinlets in my specimens; but I have no barren pinnæ in which only they occur. Except in being much more aculeate, over-ripe examples are with difficulty distinguished from *Alsophila latebrosa*; it is however perfectly glabrous on the segments beneath, whereas in the *Alsophila* there is generally some pubescence on the costules, and minute hairs on the veinlets. I have never found this fern much above 3,000 feet elevation, but the *Alsophila* is common at the highest altitudes on the Nilgiris, Pulneys and Anamallays.


GENUS III.—AMPHICOSMIA. (*Fée.*)

(*Amphi*, around; *kosmos*, world; in allusion to its wide distribution.)

Sori globose, dorsal, on a veinlet; receptacle elevated; indusium a cup-like scale below the sorus, but never completely covering the capsules as in *Cyathea*. Arborescent ferns like *Cyathea*, and a connecting link between that genus and *Alsophila*, very difficult to determine except with unripe sori; veins all free; fronds ample decompose. (*Hemitelea* only differs in having the costal veins anastomosing. None are found in India.)

1. *AMPHICOSMIA WALKERÆ*. (*Hook.*) Stipes unarmed or slightly mucronate; fronds ample bi-tripinnate, thick, firm, very coriaceous; primary pinnæ $1\frac{1}{2}$ foot long, secondary pinnæ rather remote, 3–4 inches long, deeply, nearly to the costa pinnatifid, or again pinnate, oblong, narrow-acuminate; the costa, as is the rachis, hairy above and scaly below; pinnules or lobes oblong, very obtuse, entire or crenate; veinlets once-forked, three-branched or even pinnate; sori close to the costules; indusium cup-shaped under the sorus. *Hook. Sp. Fil.* p. 30. *Bedd. F. S. I. tab.* 261.

Ceylon, centre of the island at the higher elevations. Var. β *tripinnata* has the ultimate pinnules lobed, but there are intermediate forms.



AMPHICOSMIA WALKERÆ. (Hook.)

2. AMPHICOSMIA DECIPIENS. (*J. Scott*, under *Hemitelea*.) A lofty tree fern, very prickly on the main and secondary rachises, tertiary rachis (that of the pinnules) bullate-scaly, not pubescent; pinnules glabrous, or nearly so, segments sometimes much narrowed where fertile; veinlets forked, or often three-branched and even four-branched in the sterile portions; sori in two rows close along the main veins or costa of the segment; involucre a hemispheric cup. *J. Scott*, in *Lin. Trans.* xxx. 33, t. 14; *Bedd. F. B. I.* t. 311. Shows the veins and bullate scales correctly, but not the involucre.

Sikkim and Bhotan, 1,000–4,000 feet. Khasya below Nung-klow. Differs from the next in being much more prickly and in the segments being generally longer and narrower. I do not feel certain it is distinct from *Cyathea spinulosa*, as I have not seen the involucre in very young state.

3. AMPHICOSMIA BRUNONIANA. (*Wall.* under *Alsophila*.) A large tree fern; main rachis somewhat prickly, secondary rachis slightly muricated or smooth; rachis of the pinnules crisped-pubescent; segments minutely hairy on the veinlets (under the lens), not contracted in fruit; veinlets forked, or three-branched veinlets very rare; sori and involucre as in the last. *Clarke*, p. 430. *Alsophila Brunoniana*, *Wall.* *Alsophila latebrosa*, var. *hemitelioides*, *J. Scott*, l. c. 34.

Sikkim and Bhotan, 4,000–7,500 feet; East Nepal; Khasya 3,000–5,000 feet, very common.

The specimen for which Mr. Clarke gives the locality “Deccan” is a specimen of *Cyathea spinulosa*, collected at Mendeb in Canara by Dr. Richie.

Mr. Clarke’s variety β *Scottii*, “segments of the fertile pinnules unusually large, deeply crenated, with many three-branched veinlets,” is founded on a single pinna in the Kew Herbarium, collected by Scott in Sikkim and labelled by him *latebrosa*, var. β ; it is perhaps a new species.

4. AMPHICOSMIA ALTERANS. (*Hook*, under *Alsophila*.) Rachis glabrous, smooth, turning brown; fronds subcoriaceous, glabrous, sub-

tripinnate ; primary pinnæ short, petiolate very remote, 10–14 inches long, 2–3 inches wide, oblong acuminate, deeply pinnatifid, pinnate near the base ; segments 1–1½ inch long, oblong acute subfalcate entire or serrate, all horizontal, the extremity suddenly pinnated with oblong undivided sessile pinnules serrated at the margin ; veins once forked (rarely twice-forked) ; sori in the axil of the fork, or in the lower veins, considerably above it ; involucre a hemispherical cup completely surrounding the base of the sorus. *Bedd. F. B. I. t. 236*, under *Alsophila*. *Hook. Syn. Fil. p. 41*.

Penang. My specimens distinctly show the indusium.

GENUS IV.—ALSOPHILA. (*Brown.*)

(*Alsos*, grove ; *phileo*, to love.)

Sori globose, dorsal, on a vein or at the forking of a vein ; receptacle generally elevated, often villous ; involucre none, but the bullate scales along the costa of the segments may often be mistaken for such. Arborescent ferns ; fronds decomponently pinnate ; veins never anastomosing, simple forked or pinnate.

1. *ALSOPHILA LATEBROSA*. (*Hook.*) A lofty tree fern ; stipes aculeate at the base, muricated upwards, dark mahogany-brown ; primary pinnæ oblong acuminate, 12–24 inches long, 6–8 inches wide ; pinnules lanceolate-acuminate, 3–4 inches long, cut down nearly to the rachis into linear-oblong, blunt slightly-toothed segments on each side ; rachis of pinnules beneath glabrous or pubescent ; veins generally minutely hairy (under the lens) ; veinlets forked ; numerous bullate scales along the costa of the segments beneath and also along the rachis of the pinnules ; sori occupying the lower two-thirds of the segment. *Bedd. F. S. I. t. 58*. *Hook. Syn. Fil. p. 43*.

Nilgiris and all the western mountains in the Madras Presidency and the Shevaroyes, up to 7,000 feet ; but often found as low as 3,000 ; Sikkim, Bhotan and Khasya, 3,500–5,000 feet ; also throughout the Malay Peninsula and Penang. I cannot distinguish the two varieties given by Mr. Clarke. The South Indian specimens

vary from quite or nearly quite glabrous to as pubescent as those from Sikkim, and the Penang specimens seem quite identical.

2. *ALSOPHILA GLAUCA*. (*J. Sm.*) A lofty tree fern; more or less prickly; fronds subcoriaceous, glabrous, glaucous beneath; pinnæ 2-3 feet long, pinnules 4-5 inches long, $\frac{1}{2}$ to 1 inch broad, deeply, nearly to the costa pinnatifid; segments linear-oblong subfalcate, when fertile often contracted, entire or distantly crenulate; veins forked, and often three-branched; receptacle naked or hairy. *J. Sm. in Hook. Jour. Bot.* iii. 419. *Bedd. F. B. I. t.* 86 (as *A. Brunoniana*). *Alsophila contaminans*, *Hook. Syn. Fil. p.* 41. *Chnoophora glauca*, *Bl. En. Fil. Jav.* 443.

Sikkim, Bhotan, Assam, Khasya, Cachar, Sylhet, Chittagong, Burmah and Malay Peninsula, up to 4,000 feet.

(Also in the Malay Islands.)

3. *ALSOPHILA ORNATA*. (*J. Scott.*) A lofty tree fern; stipes paleaceous at the base, slightly muricated, and, as well as the main rachises, glossy purplish-brown; fronds herbaceous-membranaceous; primary pinnæ oblong-lanceolate, rather abruptly contracted into a short, deeply pinnatifid apex; pinnules nearly all petiolate lanceolate, oblong acuminate, deeply, nearly to the costa pinnatifid; segments ovate in the sterile, narrow oblong in the fertile, slightly falcate, serrated; costa of the pinnules somewhat hairy above and scaly below; veinlets forked or a few three-branched. *J. Scott, in Trans. Lin. Soc.*, xxx. 36, t. 16 A. *Bedd, F. B. I. t.* 342

Sikkim, on the banks of the Rungbee, below the Government Cinchona plantation, at 2,500 feet elevation, called Dang pashin by the Lepchas.


4. *ALSOPHILA ANDERSONI*. (*J. Scott*) A tall tree fern; stipes muricated and rough, clothed more or less with lanceolate subulate deciduous scales; rachises tawny villous below, somewhat scabrous above; fronds ovate-lanceolate, herbaceous-membranaceous, pilose; primary pinnæ oblong-lanceolate acuminate; pinnules pinnatifid nearly to the rachis with an attenuated serrated apex; segments broad-oblong, slightly falcate, sharply serrated; veinlets simple or

once forked, 10-12 pair to each segment, all except the upper two or three with a single sorus a little below the middle; the costules, veins, and veinlets beneath all furnished with long weak white hairs, which are also present in a less degree on the upper surface. *J. Scott, in Trans. Lin. Soc. xxx. t. 17. Bedd. F. B. I. t. 310.*

Sikkim, in the tropical valleys below the Government Cinchona plantation, alt. 1,000-2,500 feet. Khasya. Called Pulai-nock by the Lepchas.

5. *ALSOPHILA OLDHAMI.*

(*Bedd.*) A tree fern, 6-10 feet, unarmed; principal rachises half an inch thick, deep purple, asperous; fronds herbaceous-membranaceous; primary pinnæ about 2 feet long by 8-9 inches broad, oblong acuminate; secondary 4-5 inches long by about 1 inch broad, lanceolate acuminate; cut down nearly to the rachis into numerous linear oblong blunt-toothed segments on each side; costa hairy above, scaly below; veins simple, or once-forked; sori in two



No. 5.

ALSOPHILA OLDHAMI. (Bedd.)

parallel rows below the middle of the veinlets. *Bedd. F. B. I. t. 343, Alsophila comosa, Scott, l. c. t. 16 B., not Wall.*

Khasya hills; Sikkim, in the forest between Darjeeling and Surail, 5,000-6,000 feet; Moulmein.

6. *ALSOPHILA COMOSA. (Hook.)* A tree fern, unarmed; stipes paleaceous, with long linear-subulate scales, pale brown, as well as the rachises, which are tawny-villous on the upper side; fronds

herbaceous, glabrous, and scaleless; primary pinnæ 14–16 inches long, 6–8 inches wide; pinnules oblong-acuminate from a rather broad and truncated sessile or shortly petioled base, serrated two-thirds of the way to the costule, pinnatifid with very narrow sinuses; lobes close-placed, oblong, serrated; veins, simple or rarely forked; sori about half-way between the margin and the costule; receptacle small, very slightly elevated. *Hook. Syn. Fil. p. 41. Bedd. F. B. I. t. 84.*

Singapore.


(Also in the Malay Islands.)

7. *ALSOPHILA COMMUTATA*. (*Mett.*) A tree fern; stipes paleaceous, at the base tuberculate-asperous; fronds bi-pinnate, partial rachis slightly squamose; pinnules all petiolate, sterile ones oblong-lanceolate, fertile portions often contracted, coriaceous, glossy as if varnished, ending in an acuminate serrulated point, pinnatifid scarcely half-way down to the rachis; segments ovate-obtuse, serrate, the margins thickened, or very slightly recurved; veins simple or often forked; sori frequently confined to the lower part of the pinnules, and a little distant from the costa of the segments. *Hook. Syn. Fil. p. 43. Bedd. F. B. I. t. 235* (as squamulata.)

Malacca, on Mount Ophir.

8. *ALSOPHILA GLABRA*. (*Hook.*) A tree fern; stipes asperous, paleaceous at the base, the rachises purple-ebeneous, polished, glabrous, or with a few scales, or with adpressed pubescence; fronds subcoriaceous-membranaceous; primary pinnæ $1\frac{1}{2}$ – $2\frac{1}{2}$ feet long, deeply pinnatifid at the apex; pinnules, upper ones sessile, lower ones petiolate from a truncated base, oblong acuminate, 3–6 inches long, 5–9 inches wide, pinnatifid $\frac{1}{4}$ to $\frac{1}{2}$ way to the costule, sometimes more in Himalayan examples; segments triangular or rounded, rarely ovate, serrated; veinlets all simple; sori arranged in the shape of a V inverted, not extending to the apex of the lobe. *Hook. Syn. Fil. p. 43. Bedd. F. S. I. t. 60.*

All the western hills of the Madras Presidency up to 4,000 feet;



ALSOPHILA CRINITA. (Hook.)

North Arcot and Cuddapa Hills, Jeypore, &c., and probably general through the northern Circars in moist shady places on the hills; Himalayas, very common in Sikkim, Nepal, &c.; Burmah and Ceylon.

(Also in the Malay Islands and S. China.)


The hairs of the rachis are exaggerated in my figure quoted above.

9. *ALSOPHILA CRINITA*. (*Hook.*) A lofty tree fern; stipes and main rachis stramineous, brown, hispid-paleaceous, and strongly muricate; spines short, tipped with a black gland; fronds subcoriaceous; rachises above pilose, beneath and on the costæ densely paleaceous-crinite; scales sometimes short and minute, generally elongate and adpressed; primary pinnæ 2 feet long, 10 inches wide; pinnules oblong acuminate, sessile pinnatifid nearly to the costules; lobes narrow-oblong, sub-obtuse, falcate, the margins recurved, serrated; costa and veins often pilose; veinlets forked; sori covering the whole under side of the lobes mixed with ciliate scales. *Hook. Syn. Fil. p. 42. Bedd. F. S. I. t. 59.*

Nilgiris, Pulneys, and Anamallays, 5,000–7,000 ft. elevation. Ceylon central provinces. By far the most beautiful of all the tree ferns.

10. *ALSOPHILA ALBO-SETACEA*. (*Bedd.*) Main rachis purplish, muricated, and (in age) only slightly scaly; pinnæ long petioled; rachis of pinnæ very scaly; pinnules with the one-two lowest segments free, the rest cut down nearly to the rachis, oblong sub-falcate, slightly crenate; veinlets all forked from near the base, one branch often (or both rarely) again forked; costæ sparingly clothed beneath with long white setaceous hairs, which are also present in a less degree on the costa and costules above; costules beneath furnished with deciduous bullate scales, which often have a hyaline setaceous point at the apex; sori copious. *Bedd. Supplmt. to Ferns, p. 2* (not figured).

Nicobar Islands (Kurz).



DIACALPE ASPIDIOIDES. (Bl.)

GENUS V.—DIACALPE. (*Bl.*)

(*Dia* and *kalpe*, a vessel, in allusion to the cup-shaped indusium.)

Sori globose, the receptacles small, scarcely elevated; indusium inferior (*i.e.*, below the sorus), globose, hard-membranaceous, entire, a length bursting very irregularly at the summit; capsules numerous, nearly sessile, ring broad, veinlets simple, free.

1. DIACALPE ASPIDIODES. (*Bl.*) A large compound fern; stipe scaly, particularly near the base; fronds tripinnate, submembranaceous, rachis of pinnæ pubescent with adpressed scales; pinnæ alternate or sub opposite, segments of pinnules oblong-cuneate, lobed, and decurrent, glabrous, or furnished with long weak scales; veinlets not reaching the margin; sori medial on the anterior lower veinlets, involucre bursting irregularly, or sometimes into two regular lips. *Bl. En. Pl. Jav. Fil.* 241. *Hook. Syn. Fil.* p. 45. *Bedd. F. S. I. t.* 257.

Nepal to Bhotan, Khasya, 4,000–6,000 feet. Ceylon, higher parts of central provinces. Tenasserim on Mooleyit 5,000–6,000 feet. Tavoy.

(Also in the Malay Islands.)

2. DIACALPE FÆNICULACEA. (*Hook. under Polystichum.*) Rhizome creeping; stipes tufted, 6–12 inches long, densely clothed below with lanceolate reddish-brown scales; fronds 1–2 feet long, 9–12 inches broad, deltoid-lanceolate, 4–5-pinnatifid, lower pinnule 6–8 inches long, 3–4 inches broad, pinnules sub-deltoid, 2–3 inches long, secondary pinnules cut down to the rachis, their lower divisions often again pinnatifid; segments linear, very acute, subcoriaceous and very glossy, with a few scattered long scale-like hairs beneath; indusium leathery, dark purple, spherical, attached by the point under its centre, not stalked, splitting irregularly from above. *Clarke, p.* 434. *Hook. Syn. Fil.* p. 256. *Bedd. F. B. I. tab.* 36, under *Lastrea* (indusium incorrect as to shape).

Sikkim, 7,000–10,000 feet elevation, abundant at Buckeem; Upper Ratong, 8,000–9,000 feet.

GENUS VI.—MATONIA. (*Br.*)

(After Dr. Maton, a London physician.)

Receptacle of the sori expanded into a firm membranaceous, umbrella-shaped obscurely six-lobed stipitate involucre, which covers and encloses six large sessile capsules; veins forked free, except those round the sori, which are closely reticulated.

1. MATONIA PECTINATA. (*Br.*)

Rhizome creeping; stipe slender, 6–8 feet high; fronds fan-like, conjugate-subpedately-flabellate, the pinnæ produced on the anterior or upper side of the divergent branches, rigid-coriaceous, linear, pinnatifid nearly to the costa, glabrous, often glaucous beneath, 1–2 feet long; sori situate at the posterior base of the segments. *Br. in Wall. Pl. As. Rar.* 1, t. 16. *Hook. Syn. Fil.* p. 45. *Bedd. F. B. I. t.* 186.

Malacca, on Mount Ophir; one of the rarest and handsomest of ferns.

(Also in Borneo.)



Nº 4

MATONIA PECTINATA. (*Br.*)

TRIBE II.—DICKSONIEÆ.

Sori globose, on the back or apex of a vein; indusium inferior, subglobose, free, sometimes covering the whole sorus, closed, or at length bursting irregularly; more frequently cup-shaped, entire, or with two lips; caudex arborescent in Cibotium; veins free, or anastomosing.

GENUS VII.—STRUTHIOPTERIS. (*Willd.*)

(*Struthios* ostrich, the fronds like feathers of the bird.)

Sori dorsal on the veins of the changed and contracted pinnæ of the fertile frond, and quite concealed by the revolute margins; indusium very thin hemispherical, very fugacious, or wanting; caudex erect or creeping; fronds stipitate, dimorphous, fertile ones pinnate; pinnæ torulose or flattish; veins all free pinnate. (Differs from *Onoclea* in having free veins.)

1. STRUTHIOPTERIS ORIENTALIS. (*Hook.*) Fronds ovate-oblong, not attenuated at the base; fertile ones oblong, contracted; pinnæ linear-oblong, flattened, two-edged, the broad refracted margins covering the whole back, dark brown, glossy, at length spreading, and torn at the margin. *Hook. Syn. Fil. p. 46. Bedd. F. B. I. t. 171.*

Sikkim, elevation 900–1,200 feet, Khasya, Assam.
(Also in Japan and Western China.)

GENUS VIII.—WOODSIA. (*Br.*)

(In honour of Joseph Wood, a British botanist.)


Sori globose; indusium inferior, soft, membranaceous, calyciform, or more or less globose, and sometimes enclosing the sorus, at length opening at the top, the margin lacinate or fringed; veins free, simple, or forked. Small herbaceous ferns, the stipes tufted, often jointed.

1. WOODSIA HYPERBOREA. (*Br.*) Glabrous, or with the stipes, rachis and costa beneath slightly hairy and scaly; fronds 5 inches long by $\frac{1}{2}$ inch broad, linear-lanceolate, pinnate; pinnæ cordate-ovate, pinnatifid, with few broadly obovate entire lobes, the lower ones distinct; indusium smaller than the sorus, but fringed with long hairs which extend beyond it. *Hook. Syn. Fil. p. 46. Clarke, p. 434.*

Kashmir, Sind Valley, elevation 8,000 feet; collected only by Mr. Levinge.

(Also in Alpine and Arctic Europe and North Asia.)

№ 9.



STRUTHIOPTERIS ORIENTALIS. (Hook.)

2. *WOODSIA LANOSA*. (*Hook.*) Like *hyperborea*, but the frond densely ferruginous, with soft long hairs and scales, stipe also villous or lanate, but becoming glabrous in age. Perhaps a very tomentose variety of No. 1. *Hook. Syn. Fil. p. 47. Bedd. F. B. I. t. 341.*

Only in Northern India—mountains of Kumaon; Pindari, 12,000 feet; above Namik, 11,600 feet; Sikkim Lachen, 14,000 to 16,000 feet; Mount Donkia, 18,000 feet.

3. *WOODSIA ELONGATA*. (*Hook.*) Glandulosely pilose; fronds up to 12 inches long by 1–1½ inch broad, narrow oblong pinnate, pinnæ rather distant alternate sessile from a broadish base, oblong obtuse, pinnatifid not more than half down, lobes short, rounded, dentate; sori globose, much elevated; indusium larger than the sorus, very membranaceous, globose, soon bursting with a lacerated apex, not ciliated. *Hook. Syn. Fil. p. 47. Bedd. F. B. I. t. 14.*

Himalayas, from Dhurmsala to Sikkim, 8,000 to 12,000 feet; common.

GENUS IX.—PERANEMA. (*Don.*)

(*Peri*, around; *nema*, a thread; the sorus on a thread-like stalk.)

Sori globose, on the back of a vein or veinlet; indusium inferior, globose, coriaceous, pedicellate, at first enclosing the whole sorus, at length bursting vertically into two spreading lips; veins free, forked, clavate; fronds large, decomposed. *Sphæropteris*, *Wallich.*

1. *PERANEMA CYATHEOIDES*. (*Don.*) Fronds up to 4–6 feet high, stipate, deltoid decomposed, ultimate pinnules sessile, linear-oblong, obtuse, half an inch long, pinnatifid, repand, or crenate; main rachis scaly or pubescent. *Don. Fr. Fl. Nep. p. 12. Bedd. F. S. I. t. 73. Sphæropteris barbata, Wall. Hook. Syn. Fil. p. 49.*

Nepal and Bhotan, 6,000–10,000 feet; Khasya, 4,500–6,000 feet; Anamallays, 6,000 feet.



А. С. СЕРГЕЕВ И С. С. СЕРГЕЕВ

WOODSIA ELONGATA. (L'noč.)

GENUS X.—CIBOTIUM. (*Kaulf.*)(*Kibotos*, a casket—form of the indusium.)

Sori at the apex of a vein, intra-marginal; indusium distinctly two-valved, the outer valve coriaceous, distinct from the substance of the frond; veins free, simple, forked, or pinnate; arborescent, with large decompound coriaceous fronds. *Only differs from Dicksonia in the outer valve of the indusium being distinct from the substance of the frond.*

1. CIBOTIUM BAROMETZ. (*Link.*) Arborescent; fronds tri-pinnate, lower pinnæ ovate-lanceolate, 1–2 feet long, 6–12 inches broad; pinnules linear-acuminate, cut down within a short distance of the rachis above, and sometimes quite down to it at the base, segments linear-oblong, acute, subfalcate, upper surface naked, shining, lower glaucous, sometimes furfuraceous; sori 2–12 to a lobe; the valves nearly equal, transversely oblong. *Link, Fil. Sp. p. 166. Hook. Syn. Fil. p. 49. C. glaucum, Bedd. F. B. I. t. 83.*

Mishmee, Assam, Tavoy.

(Also in the Malay Islands and South China.)

GENUS XI.—DENNSTÆDTIA. (*Bernh.*)

(In honour of Dennstadt, a German writer.)

Sori at the apex of a vein; indusium cup-shaped, not at all, or very indistinctly, two-valved; veins free, simple, forked, or pinnate; not arborescent; fronds herbaceous or membranaceous. *Patania Presl. a name of later date. Dicksonia Sec. Patania. Hook.*


1. DENNSTÆDTIA SCABRA. (*Wall. under Dicksonia.*) Rhizome wide-creeping, villous; main rachis scabrous, often wavy; fronds very variable in size, up to $2\frac{1}{2}$ –3 feet long, and 2 feet broad; deltoid or ovate, bipinnate, surface beneath with scattered glistening hairs,



PERANEMA CYATHEOIDES. (Don.)

lower pinnæ 4–8 inches long, lanceolate, pinnules quite distinct, 1–1½ inches long, the lower ones cut down nearly to the rachis into numerous pinnatifid oblong deltoid segments; sori 2–6 to the lower segments; indusium cup-shaped, subglobose. *Wall. Cat. Dicksonia*, *Hook. Syn. Fil. p. 54.* *Dennstædtia deltoidea*, *Bedd. F. S. I. t. 258.*

Ceylon, central provinces; Himalayas, Kumaon to Bhotan, 4,000–8,000 feet; Khasya, 4,000–6,000 feet; Birma and the Malay Peninsula generally, very common.



Nº 12


DENNSTÆDTIA
SCABRA. (*Wall.*)

2. DENNSTÆDTIA ELWESII. (*Baker, under Patania.*) Fronds lanceolate, tripinnatifid, 1–1½ foot long; rachis stramineous glabrous; pinnæ lanceolate, nearly sessile, 3–4 inches long, ¾–7⁄8 inch broad, the rachis flattened; pinnules 20–30-jugate, lanceolate, sessile cuneate at the base, 1½–2 lines broad, deeply pinnatifid; segments ligulate, erecto-patent, many entire parallel, lowest on the upper side often bifid, both sides glabrous; veins solitary in the centre of each segment; sorus covering the entire apex of the segment. *Hook. Syn. Fil. p. 54.* *Bedd. Suppl. to Ferns, t. 347.*

Sikkim, near Lachen, 8,500 feet; Choongtam, 9,000 feet.

3. DENNSTÆDTIA APPENDICULATA. (*Wall. under Dicksonia.*) Fronds oblong or lanceolate, bipinnate, 12–18 inches long, 6 inches broad; rachis hairy; pinnæ close together, their rachises parallel, lower ones linear-lanceolate, 3–4 inches long, cut down to the rachis; pinnules linear-oblong, ½ inch long, 2 lines broad, pinnatifid into linear segments, under surface glandulosely pilose; sori 2–6 to a pinnule placed at the base of the sinuses; indusium cup-shaped, globose. *Wall. Cat. 65.* *Dicksonia (Patania) appendiculata.* *Hook. Syn. Fil. p. 54.* *Bedd. F. B. I. t. 82.*

Senchal above Darjeeling, 8,500 feet; Lachen Valley, in Sikkim, Nepal; Kumaon Gori Valley, 5,500 feet; Banks of the Vishnugunga, above Panchkisar, 8,000–9,000 feet.



CIBOTIUM BAROMETZ. (Linn.)

TRIBE III.—HYMENOPHYLLÆ.


Sori terminal or marginal from the apex of a vein; receptacle elongated, often filiform and long, and exserted more or less, clothed especially below with sessile orbicular imbricated subpeltate compressed capsules, surrounded by a complete transverse ring, opening vertically; indusium inferior, various in shape, generally of the same texture as the frond. Small, often epiphytal ferns, herbaceous-membranaceous, more or less laxly cellular, variously costate and veined. Caudex frequently long, creeping, and filiform.

GENUS XII.—HYMENOPHYLLUM. (L.)

(*Hymen*, membrane; *phyllon*, leaf.)

Sori marginal, more or less sunk in the frond or exserted, terminating a costa or vein; indusium inferior, more or less deeply two-lipped or two-valved, toothed, fringed, or entire; receptacle elongated, columnar, exserted, or included; capsule mostly orbicular, depressed, attached by the centre, furnished with a broad transverse ring, opening irregularly at the apex. Small, often minute ferns, growing on trunks of trees and damp rocks in moist places, generally on the mountains; fronds delicately membranaceous, simple or compound, costate, or with simple or branched, never anastomosing, veins.

* Margin of the frond entire.




Nº 14.

HYMENOPHYLLUM
PARVIFOLIUM. (*Baker.*)

1. HYMENOPHYLLUM PARVIFOLIUM. (*Baker.*)

A very tiny species, rhizome slender, creeping, tomentose; stipes about 1 line long; frond very minute, 2-3 lines long, 1 line broad, linear-oblong, margin entire; simple or 2-3-cleft at the apex, sometimes half-way down, furnished only with a central costa and a few faint, irregular, free spurious venules, the margin not thickened; sori solitary, terminal, the cuneate base sunk in the frond divided



A. CATTILLO & E. C. S.

DENNSTÆDTIA APPENDICULATA. (Wall.)

about half way down ; valves rounded entire. *Baker, Syn. Fil.* p. 56. *Bedd. F. B. I. t.* 225.

Moulmein.


2. HYMENOPHYLLUM TENELLUM. (*Kuhn.*) Rhizome wide-creeping hairy or with fine hair-like scales ; stipes $\frac{1}{2}$ – $1\frac{1}{2}$ inch long ; slender, somewhat hairy, fronds 1–2 inches long, $\frac{3}{4}$ –1 inch broad, oblong to deltoid, tripinnatifid, margin entire, main rachis winged, more or less furnished beneath with ferruginous hair-like scales ; pinnæ 4–6 jugate, deltoid, cuneate-truncate at the base on the lower side, lower pinnules 1–2, cleft at tip with ligulate blunt lobes, $\frac{1}{4}$ line broad, partial rachises and veins more or less hairy beneath, glabrous above ; sori 2–6 to a pinna, terminal in the lobes ; involucre with hemispherical lobes, and a cuneate entire immersed base. *Baker, Syn. Fil.* p. 57. *H. polyanthos*, *Bedd. F. S. I. t.* 267. *H. emersum*, *Baker, Syn. Fil.* 1st ed. p. 457.

Ceylon. (C. P. 3,360.) Differs from any form of *polyanthos* in having the rachises and veins hairy beneath.

3. HYMENOPHYLLUM EXSERTUM. (*Wall.*) Stipe 1–2 $\frac{1}{2}$ inches long, slender ; fronds 2–6 inches long, 1–2 inches broad, lanceolate-oblong, ovate or short triangular bipinnatifid ; main rachis winged above or throughout, more or less densely clothed with deciduous ferruginous hairs ; lower pinnæ rhomboidal, lanceolate-acuminate, divided more than half way down to the rachis into simple or forked linear pinnules, slightly hairy on the principal veins ; sori 2–8 to a pinna, terminal or axillary in the segments on both sides, involucre divided down nearly to the base ; valves oblong, bluntly denticulate at the apex. *Hook. Syn. Fil.* p. 58. *Bedd. F. S. I. t.* 9.

Himalayas and Khasya Hills, 2,000 to 9,000 feet ; Birma ; all the Western Ghats of the Madras Presidency ; Ceylon, central provinces. Very common.

4. HYMENOPHYLLUM POLYANTHOS. (*Sw.*) Stipes 2–3 $\frac{1}{2}$ inches long, wingless or narrowly winged towards the apex ; fronds glabrous, or nearly so, 2–8 inches long by 1–3 inches broad, not crisped, ovate-



HYMENOPHYLLUM EXSERTUM. (Wall.)

oblong tripinnatifid ; main rachis with a narrow wing ; lower pinnae triangular-rhomboidal, divided down to a narrow centre into several pinnules on each side, the lowest of which are cuneate or rhomboidal, deeply pinnatifid, ultimate segments linear, $1-1\frac{1}{2}$ line long, less than $\frac{1}{2}$ a line broad ; sori 2-12 to a pinna, terminal or axillary on the segments on both sides ; involucre small, divided down nearly to the base ; valves ovate or rounded entire or slightly denticulate. *Swartz, Syn. Fil. t. 149. Hook. Syn. Fil. p. 60. H. microglossum, V.D.B. Hook. Syn. Fil. p. 59. H. polyanthos β . minor. Bedd. F. B. I. t. 306. H. Blumeum, Spreng. Bedd. F. S. I. t. 266.*

Himalayas and Khasya mountains, 1,000-12,000 feet elevation. Western Ghats of the Madras Presidency ; Ceylon ; Birma.

(Also widely distributed in all tropical and subtropical regions throughout the world.)

The typical form is only from Northern India, and is a very large fern.

VAR. β MICROGLOSSUM is from the western ghats of South India. *Bedd. F. B. I. t. 306.* A small form with the stipe more or less winged ; but Mr. Clarke sends me a fern under this name from Bhotan, which is intermediate between this and the type with some of the fronds running very close to Blumeum.

VAR. γ BLUMEANUM. Fronds very narrow and elongate, and pinnae sometimes simply pinnatifid only. *Bedd. F. S. I. t. 266.*

Ceylon ; Tinnevely and Travancore mountains. This has generally been considered a quite distinct species by botanists in Ceylon and South India, but Mr. Clarke says it runs into the type in Northern India, and cannot be separated. With only Ceylon and South Indian specimens in view, it is difficult to consider them all forms of one species, but after seeing the Himalayan forms, I quite agree with Mr. Clarke that they cannot be separated as species.

5. HYMENOPHYLLUM JAVANICUM. (*Spreng.*) Glabrous or nearly so ; stipe 2-4 inches long, erect, margined above with a broad, crisped wing ; fronds 4-8 inches long, 3-4 inches broad, triangular,

tripinnatifid; main rachis with a broad crisped wing; lower pinnæ $1\frac{1}{2}$ –2 inches long, triangular-rhomboidal, erect-patent, divided down to a narrow crisped centre into several plane or slightly crisped pinnules on both sides, the lower of which are several times deeply pinnatifid; sori 6–20 to a pinna, terminal and axillary on the segments on both sides; involucre about as broad as the segments, divided down nearly to the base; valves orbicular, entire, or denticulate. *Spreng. Syst.* iv. 132. *Hook. Syn. Fil.* p. 60. *H. crispatum* (Wall.), *Bedd. F. S. I. t.* 207.

Himalayas, Nepal, and Bhotan, 5,000–8,000 feet; Khasya, 3,500–5,500 feet; Western Ghats of the Madras Presidency; Ceylon, central provinces; Birma and Malay Peninsula.

(Also in Australia, New Zealand, Mauritius, Bourbon, and the Philippines.)

VAR. β BADIUM. (*Hook. and Grev.*) Frond hardly or not at all crisped. *H. badium*, *Hook. Syn. Fil.* p. 60. *Bedd. F. B. I. t.* 282.

Sikkim, Malay Peninsula, Tenasserim on the Mooleyit Mountain.

6. HYMENOPHYLLUM CILIATUM. (*Sw.*) Stipe 1–2 inches long, furnished with tufts of stellate hairs, decurrently winged above; wing ciliate; frond oblong, acuminate, tripinnatifid, 2–6 inches long, 1–2 inches broad at the centre; main rachis broadly winged throughout ciliated and hairy like the stipe; lower pinnæ oblong or rhomboidal, with a broad central undivided portion, and numerous erect-patent simple or forked ciliated linear segments 2–3 lines long; sori 2–12 on a pinna, placed at the end of the lateral segments on both sides, involucre immersed suborbicular; valves divided about half-way down, and conspicuously ciliated. *Hook. Syn. Fil.* p. 63. *Bedd. F. B. I. t.* 305.

Sikkim.

(Also in tropical America from Cuba and Mexico to Chili and South Brazil; Tropical Africa, Mauritius, Madagascar, Bourbon, Seychelles and Johanna Island; New Zealand.)

Mr. Clarke seems to doubt the Sikkim locality, but my specimens received from Dr. Jerdon quite agree with the type.

**** Margin of the frond serrulate.**

7. *HYMENOPHYLLUM SIMONSIANUM*. (*Hook.*) Stipe slender, wiry or winged, naked or with a few ferruginous hairs, 1–2 inches long; fronds linear-oblong, 2–6 inches long, up to 1½ inch broad, pinnatifid to the winged rachis; the primary segments fan-shaped, 1–6 lobate, ultimate segments broadly oblong, serrulate; sori 1–4 terminal on the divisions of the upper pinnæ, involucre broadly oblong, divided nearly to the base; valves serrulate. *Hook. Syn. Fil. p. 68. Bedd. F. B. I. t. 281.*

Sikkim, 4,000–10,000; common. Khasya Hills.

8. *HYMENOPHYLLUM SMITHII*. (*Hook.*) Stipe 1–2 inches long, wiry naked or slightly tomentose; fronds oblong-acuminate, 3–8 inches long, 2–3 inches broad, bipinnatifid; rachis winged above, ciliated below; pinnæ oblong-triangular, with a winged rachis and simple or 1–4 times forked linear-serrulate segments; sori 1–4 terminal on the segments of the pinnæ on both sides; involucre oblong, small, divided half down; valves entire. *Hook. Syn. Fil. p. 69. Bedd. F. B. I. t. 277.*

Malacca and Penang.

(Also in Philippines and Java.)

9. *HYMENOPHYLLUM DENTICULATUM*. (*Sw.*) Stipe 1–2 inches long, erect, ferruginously hairy or naked, often winged towards the apex; rachis winged throughout; fronds 2–3 inches long by 1 or more broad at the base, ovate to narrowly oblong, glabrous, or with scattered hairs, slightly crisped, pinnatifid to the rachis; primary segments pinnatifid, secondary ones oblong, sometimes again divided, serrulate on the margin, involucre ovate, glabrous or scabrous, or rarely slightly hairy; valves serrulate, or pectinate. *Swartz, Syn. Fil. t. 148 and 375. Hook. Syn. Fil. p. 70. Bedd. F. B. I. t. 278. Clarke, Ferns of North India, p. 438.*

Khasya hills, about Cherra Poonjee, 4,000–5,000 feet; Bhotan, Birma, Moulmein mountains, 7,000 feet.

(Also in Java.)

VAR. β FLACCIDUM. Main rachis often with much ferruginous hair beneath; fronds flaccid, hardly at all crisped, ultimate segments slightly serrulate in their upper portion only, the main rachis quite entire, valves of the involucre slightly serrulate at the summit. *H. Khasianum*, *Hook. Syn. Fil.* p. 464. *Bedd. F. B. I. t.* 276. *Clarke, l. c.*, p. 438.

Khasya Hills, growing with the type. Mr. Clarke, who has seen it growing, considers it only a variety of *denticulatum*.

10. HYMENOPHYLLUM

NEESII. (*Hook.*)

Stipe 2–1 inches long, naked, or slightly winged, with a crispate margin on both sides; frond ovate, about 2 inches long, $\frac{1}{4}$ –1 inch broad, tripinnatifid; rachis winged throughout, the wing and pinnæ much crisped; pinnæ with distant, narrow, simple or 1–3 times deeply forked, deeply toothed segments; sori small, usually single, supra-

axillary on the upper pinnæ; involucre subcylindrical below, divided more than half-way down, with two acute spinulose-dentate valves. *Hook. Syn. Fil.* p. 71. *Bedd. F. S. I. t.* 279. *Hymenophyllum Tunbridgense*, *Bedd. F. S. I. t.* 265.



HYMENOPHYLLUM NEESII.
(*Hook.*)

Ceylon, central provinces ; Tinnevely on the Chokampatty hills, 5,000 feet ; Penang.

(Also in Java, Borneo, Fiji, and Philippines.)

11. *HYMENOPHYLLUM LEVINGII*. (*Clarke*.) Very delicate in texture ; stipe 1 inch, with moniliform hairs ; frond 1-2 inches long, more or less covered with moniliform hairs, narrowly oblong, not crisped, pinnatifid to the winged rachis ; primary segments 1-4 lobate, ultimate segments oblong, remotely serrate, their midrib with many hairs and lanceolate scales of the same texture as the frond ; involucre usually 1-2 at the end of the segment, small glabrous subquadrate ; valves separating nearly to the base entire or slightly toothed at the apex. *Clarke, F. of North India, p. 439. t. xxix. fig. 3.*

Sikkim ; Yoksun and Neebay, 7,000 feet elevation.

GENUS XIII.—TRICHOMANES. (*Smitn.*)

(*Thrix*, a hair ; *manos*, soft.)

Sori marginal, always terminating a vein, more or less sunk in the frond, involucre monophyllous, tubular, closely corresponding with the frond in texture, the mouth truncated or winged, or slightly two-lipped ; receptacle filiform, elongated, often considerably exerted beyond the mouth of the involucre, capsuliferous principally at the base ; capsule sessile, depressed, surrounded by an entire broad nearly transverse ring, bursting vertically. Like the last genus in habit of growth and delicacy of texture.

1. *TRICHOMANES MOTLEYI*. (*I. D. B.*) Rhizome creeping, more or less tomentose ; fronds very small, sessile, or shortly stalked, about $\frac{1}{4}$ inch broad, cordate-orbicular, sometimes somewhat lobed, sterile ones nearly entire, fertile ones with a deep apical sinus ; veins in the sterile fronds radiating from the base without any distinct central costa, but a distinct costa is present in the fertile fronds ; spurious veins indistinct, a distinct marginal band ; involucre

stalked, proceeding from the base of the sinus, the mouth spreading and much dilated. *Hook. Syn. Fil. p. 73.* T. Henzaiense, *Bedd. F. B. I. t. 183.*

Tenasserim, Andamans, Ceylon.

(Also in Borneo and New Caledonia.)

2. TRICHOMANES EXIGUUM. (*Bedd. under Hymenophyllum.*)

Rhizome wide-creeping; stipe very short; frond $\frac{1}{4}$ to $\frac{1}{2}$ inch linear-oblong, narrowed at the base, blunt repand, rarely somewhat pinnatifid; veins pinnate from a central costa, simple or forked, spurious venules few but nearly as prominent as the veins, not reaching the costa, and often not touching the margin; involucre solitary, terminal, the base sunk in the frond; valves entire, large and spreading; receptacle exserted or included. *Bedd. F. B. I. t. 275.* *Hook. Syn. Fil. p. 464.*

Wynaad and Coorg, 3,000–4,000 feet; Ceylon.

3. TRICHOMANES NEILGHERRENSE. (*Bedd.*) Rhizome creeping more or less tomentose; stipe 3–6 lines long, glabrous or tomentose, fronds glabrous, very small up to 1 inch long, ovate to elliptic lanceolate or linear, entire or somewhat lobed; veins from a more or less distinct central costa, spurious venules numerous, a marginal band generally present; involucre terminal, 1–2 sunk in the frond, the mouth exserted, more or less two-lipped; receptacles exserted. *Bedd. F. S. I. t. 6.* *Hook. Syn. Fil. p. 74.*


Nilgiris, abundant about Walaghat, 3,000–4,000 feet; Anamallay hills, Tinnevely and Travancore hills; Ceylon, central provinces.

4. TRICHOMANES HENZAIANUM. (*Parish.*) Rhizome creeping, tomentose; stipe from obsolete up to $\frac{1}{2}$ inch long; frond $\frac{1}{4}$ – $\frac{1}{2}$ inch each way, very thin, orbicular to obovate-cuneate, more or less lobed, central costa present or often obsolete, except towards the base, above which it branches off into veins; veins prominent and distinct, spurious venules copious, marginal band absent or inconspicuous; involucre 1–6 sunk in the frond; the mouth two-lipped.


Parish in Hook. 2 Cent. Ferns, t. 1. Hook. Syn. Fil. p 75. Bedd. F. B. I. t. 303.

Moulmein and Pegue, on trees.

5. *TRICHOMANES MUSCOIDES*. (Sw.) Frond stalked, 1-3 inches long, by $\frac{1}{4}$ - $\frac{1}{2}$ inch broad, linear-oblong to suborbicular, entire or



variously lobed, spurious venules copious ; sori 1-8, terminal on the lobes, the tube sunk in the frond, the mouth free, dilated, but usually scarcely two-lipped. *Hook. Syn. Fil. p. 75. Bedd. F. B. I. t. 304.*



Nº 18.

TRICHOMANES PARVULUM.
(Poir.)



Northern India (received from Dyas) ; Khasya.

(Also in tropical America and Africa, Polynesia, &c.)

Mr. Clarke states that the typical form is

not found in Northern India, but only the variety *sublimbatum*, distinguished by having no trace of a marginal band, but in Griffith's specimen collected on the Khasya hills, this marginal band is more or less present in some fronds, and I do not think it can be separated as a variety. This species, as I understand it, is not


present in Ceylon or S. India, the Ceylon plant being identical with the Nilgiri *T. neilgherrense*.

6. *TRICHOMANES PARVULUM*. (*Poir.*) Rhizome wide-creeping, interlaced; stipe 1 inch long, wiry, slender, tomentose below; frond $\frac{1}{4}$ to $\frac{1}{2}$ inch each way, orbicular in general outline, cuneate or truncate at the base, flabellately cut about half-way down from the outer edge in the direction of the base into narrow irregular segments; veins close, prominent, so that the frond, when dry, appears channelled; dichotomous at a very small angle; spurious venules numerous; sori 4-6, terminal on the central segments, the tube quite sunk in the frond, the mouth dilated at the sides. *Hook. Syn. Fil. p. 75. Bedd. F. S. I. t. 179.*

Nilgiris.

(Also in Japan, China, Java, Polynesian Islands, Madagascar, Johanna Island, and the Moluccas.)

7. *TRICHOMANES PROLIFERUM*. (*Blume.*) Rhizome wide-creeping, interlaced; stipe 1-2 inches long, slender, bearing 1-3 deeply divided fronds about 1 inch broad each way, ovate-lanceolate to roundish; segments narrow dichotomously branched, divided down nearly to the central rachis; veins numerous, irregular; sori 4-12, terminal, the tube quite sunk, the mouth obscurely two-lipped; receptacles exserted. *Hook. Syn. Fil. p. 76. Bedd. F. S. I. t. 262.*



Nº 19.

TRICHOMANES DIGITATUM.

(*Sw.*)

Nilgiris, west slopes, 3,000 feet; Travancore Hills, Ceylon.

(Also in Java and the Philippine Islands.)

8. *TRICHOMANES DIGITATUM*. (*Swartz.*) Stipe very slender, naked, $\frac{1}{4}$ inch long; frond 1-2 inches long, $\frac{1}{4}$ -1 inch broad, irregular in general outline, divided down nearly to the base, or a broadly-

winged rachis into long broadly-linear dichotomous segments, sometimes ciliate, with only a central costa in each; sori 2-6, terminal on the segments; involucre cup-shaped, quite sunk, the mouth broadly dilated, slightly two-lipped. *Hook. Syn. Fil. p. 76. T. corticola, Bedd. F. S. I. t. 264.*

Ceylon, Ambagamwa district; also collected by Griffith either in North India or the Malay peninsula.

(Also in Mauritius, Bourbon, Java, and the Polynesian Islands.)

9. *TRICHOMANES KURZII. (Bedd.)* Rhizome slender, wide creeping; stipe very short; frond not 1 inch long, cuneate below,



TRICHOMANES KURZII. (Bedd.)

deeply pinnatifid, primary segments 1-3 lobate, ultimate segments linear, texture thick, surface somewhat undulated, margin bounded by a thickened line, a central vein only in each segment; sori 1-2 to a frond, subterminal, the involucre turbinate, sunk in the frond, the mouth dilated but without lips. *Bedd. F. B. I. t. 286. Trich. nanum, Hook. Syn. Fil. p. 77. Crepidomanes nanum, V. D. B. Hymen. Suppl. p. 122. Clarke, F. N. I. p. 440.*

Assam, Andaman Islands; Malabar, at Poodoopady, foot of the Tambacherry Ghat.

10. *TRICHOMANES VIRIDANS. (Mett.)* Rhizome creeping, tomentose; fronds with a broadly winged stipe $\frac{1}{2}$ -1 inch long, very irregular in outline, more or less deeply pinnatifid nearly down to the midrib; segments again more or less deeply divided, obtuse and rounded at the apex of the lobes; veins erect-patent from a central costa one to each lobe of the segments; spurious venules lineolate, numerous, very short and detached; sori numerous, terminal

at the apex of the lobes, tube almost entirely exerted, only the base being sunk in the margin of the frond, the mouth dilated. *Kuhn, Lin.* 35, p. 389. *T. pusillum, Bedd. F. B. I. t.* 302. (*Not of Swartz.*)

Moulmein.

11. *TRICHOMANES INTRAMARGINALE. (Hook and Grev.)* Rhizome slender, creeping, tomentose; stipe short; frond 1–2 inches long, $\frac{1}{4}$ – $\frac{1}{2}$ inch broad, tapering below, erect pinnatifid, central portion 1 line broad, segments few, erect, simple, forked or pinnate, subcoriaceous, opaque, the apices retuse, a central costa only in each segment, but the margin thickened; involucres 1–4 to a frond, subcylindric, tapering at the base, sunk entirely in the apex of the segments; receptacles protruding. *Hook. Syn. Fil. p.* 79. *Bedd. F. S. I. t.* 208.

Sivagiri Hills (Tinnevely district); Malabar plains, foot of the Bhagamandal Ghat, which descends from Coorg; Ceylon, in the Ambagamwa district.

12. *TRICHOMANES PALLIDUM. (Blume.)* Rhizome creeping, slender, tomentose; stipe 1–3 inches long, wiry, slender; fronds 2–6 inches long by 1–1 $\frac{1}{2}$ inch broad, ovate to lanceolate bipinnatifid, divided down to a narrowly winged rachis, glauco-fuscous when dry, primary divisions broad-lanceolate, the segments all acute-linear, simple or forked; veins broad and few; sori 1–4 to a pinna, superaxillary on short segments partially sunk, short cylindrical, attenuated at the base, the mouth broad-spreading, but scarcely two-lipped. *Blume Hook. Syn. Fil. p.* 80. *T. glauco-fuscum, Hook. Sp. Fil. 1, p.* 128. *Bedd. F. S. I. t.* 263.

Ceylon, Ambagamwa district; Penang.

(Also in the Pacific Islands, and the Philippines.)

13. *TRICHOMANES BIPUNCTATUM. (Poir.)* Rhizome wide-creeping, tomentose, rather slender, stipes 1–2 inches long, naked slightly winged above; frond 1–4 inches long, 1 $\frac{1}{2}$ –2 inches broad,

ovate, tripinnatifid, main rachis with a very narrow wing or free below; pinnæ ovate-rhomboidal, pinnatifid down to a narrowly-winged rachis; the lower pinnule again deeply pinnatifid; ultimate segment linear; texture firm, membranaceous; a central costa in each segment, spurious venules none or indistinct; sori 1-4 to a pinna, axillary or terminal on the segments; tube sunk or somewhat exserted; the mouth very distinctly two-lipped, the lips subtriangular with a rounded apex. *Poirot in Encyc. Méth. Bot.* viii. 69. *Trich. Filicula, Bory. Hook. Syn. Fil. p. 81. Bedd. F. B. I. t. 283.*

Himalayas and Khasya hills, up to 6,000 feet; all the western forests of the Madras and Bombay Presidencies, up to 8,000 feet; Ceylon; Birma.

(Also throughout the tropics of the whole world.)

VAR. β INSIGNE. (*V. D. B.*) A narrower more slender form. *Bedd. F. B. I. t. 284.*

N. W. Himalaya.

VAR. γ PLICATUM. (*V. D. B.*) Ultimate segments of the pinnules very narrow and acute; wing of the rachis somewhat crisped. *Bedd. F. B. I. t. 285.*

Birma.

14. TRICHOMANES PYXIDIFERUM. (*L.*) Rhizome wide-creeping, tomentose, rather slender; stipe 1-2 inches long, naked, winged above; frond 1-6 inches long, 1-1½ inch broad, ovato-oblong, tripinnatifid, main rachis with a very narrow wing; pinnæ ovate-rhomboidal, pinnatifid down to a narrowly-winged rachis, with the lower pinnule again deeply pinnatifid; ultimate segment linear, often emarginate at the apex; texture membranaceous; a central costa in each segment; spurious venules none or indistinct; sori 1-4 to a pinna, axillary, tube more or less winged, mouth broadly dilated but scarcely two-lipped; receptacle filiform, exserted. *Hook. Syn. Fil. p. 81. Bedd. F. S. I. t. 7. F. B. I. t. 301.* A small variety.

Scarcely distinguishable from No. 13, except by the involucre.

South Indian forests, Ceylon, Birma, Eastern Bengal.
(Also in the tropics throughout the world.)

VAR. β LIMBATURE. Fronds up to 8–10 inches long, by $1\frac{1}{2}$ –2 inches broad, flaccid, and larger and less cut than the type. *Bedd. F. B. I. t.* 348.

Khasya Hills, 6,000 feet.

15. TRICHOMANES BIRMANICUM. (*Bedd.*) Rhizome thick, wiry, wide-creeping, tomentose; stipe 1–3 inches long, winged throughout; frond 2–7 inches long up to $2\frac{1}{2}$ inches broad, ovate, rachis winged; pinnae very compound, very minutely furfuraceous (under the lens), the ultimate segments very narrow, $\frac{1}{4}$ line broad, a single costa to each ultimate segment; sori copious, supra-axillary, much exserted, the mouth truncated. *Bedd. F. B. I. Suppl. t.* 349.

Birma, common on the Mooleyit mountain, 5,000–6,000 feet.

16. TRICHOMANES RADICANS. (*Sw.*) Rhizome wiry, wide-creeping, tomentose; stipe strong, up to 6 inches long, naked or nearly so; fronds up to 12 inches long and 6 inches broad, 3–4-pinnatifid, main rachis naked or winged sometimes to the base of the stipe, lower pinnae 1–4 inches long, ovate-rhomboidal, ultimate segments oblong, one-nerved, texture firm, membranaceous; sori lateral, 1–4 to a pinnule, the tube small, subcoriaceous, more or less exserted, the mouth slightly lipped or altogether truncated, receptacle slender, elongated. *Swartz, Fl. Ind. Or.* 1736. *Hook. Syn. Fil. p.* 81. *Bedd. F. B. I. t.* 181. *T. umbrosum, Wallich.*

Himalayas from Nepal to Bhotan, 2,000–7,000 feet; common in Khasya, 2,000–5,500 feet; Mergui.

(Also scattered throughout warm, temperate regions of both hemispheres, and known as the Irish fern.)

Mr. Clarke says it often climbs to the height of 10 feet, and is 1–2 pinnate with finely divided fronds, in which state it is very distinct; but it also varies so as to be with difficulty distinguished from *pyxidiferum* on one hand and *auriculatum* on the other.

VAR. β ANCEPS. (*Wall.*) Frond smaller; stipe often winged to the base, primary segments pinnatifid or somewhat 2-pinnatifid; lips of the involucre slightly broader than the tube. *Clarke, F. N. I. p.* 441.

Sikkim and Khasya, frequent.

17. TRICHOMANES AURICULATUM. (*Blume.*) Rhizome strong, wide-creeping, tomentose; frond nearly sessile, 1-2-scarcely 3-pinnatifid, 6-12 inches long, 1-2 inches broad, rachis more or less winged, pinnæ shortly stalked, ovate-rhomboidal, obliquely cuneate at the base, irregularly pinnatifid half down or more, ultimate segments ovate entire with flabellate nerves, or narrowly oblong one-nerved, texture subcoriaceous; sori 2-12 to a pinna, the tube nearly or quite exserted, the mouth truncate. *Blume, Fl. Jav. Fil. p.* 225. *Hook. Syn. Fil. p.* 82. *Bedd. F. B. I. t.* 182.

Sikkim, Bhotan, 2,000-7,000 feet, frequent; Khasya, 3,000-5,500 feet, common; Cachar.


(Also in Japan, Formosa, Philippines, Java, and Guiana.)

18. TRICHOMANES JAVANICUM. (*Blume.*) Caudex tufted with strong wiry roots; stipe 1-4 inches long, wiry, erect, naked, or villous; fronds 2-8 inches long, 1-2 inches broad, ovate-lanceolate, acuminate, simply pinnate, pinnæ oblong serrate, or linear fimbriate, texture subcoriaceous; veins numerous, close, central costa not clearly marked or continuous to the apex; sori 1-4, placed in the axils of the linear segments on the upper side of the pinnæ; tube free, the mouth broadly dilated, but not two-lipped, receptacle much exserted. *Bl. En. Pl. Jav. Fil.* 224. *Hook. Syn. Fil. p.* 83. *Bedd. F. B. I. t.* 180.

Chittagong, up to 1,000 feet; Cachar; Malay peninsula.

(Also in Borneo, Java, Polynesian Islands, and Madagascar.)

19. TRICHOMANES RIGIDUM. (*Swartz.*) Stipe tufted, erect, wiry, 2-8 inches long, naked or very slightly winged above; fronds 2-8 inches long, 2-6 inches broad, deltoid or ovate, acuminate, 3-4 pinnatifid, the main rachis usually only slightly winged towards



A. CASTELLANOS

TRICHOMANES JAVANICUM. (Bl.)

the top ; lower pinnæ 2–3 inches long, erect-patent, ovate or lanceolate-rhomboidal cut down the rachis ; pinnules deeply pinnatifid with deeply toothed or pinnatifid linear lobes, texture subcoriaceous, colour dark green, nearly black when dry, nearly naked, often slightly crisped ; sori 2–16 to a pinnule, small axillary, the tube more or less exserted, the mouth slightly dilated. *Hook. Syn. Fil. p. 86. Bedd. F. S. I. t. 8.*

Common in the moist forests on the western mountains of the Madras Presidency, 2,000–4,000 feet elevation, and in Ceylon and the Malay Peninsula.

(Also in tropical America and the West Indies, Polynesian Islands, New Zealand, Japan, South China, South Africa, Angola, Fernando Pó, and the Mascareen Islands.)

TRIBE IV.—DAVALLIÆ.

Sori marginal or submarginal, roundish, covered by a reniform or suborbicular, squamiform indusium, which is open at the apex, fastened broadly at the base, open or free at the sides.

GENUS XIV.—HUMATA. (*Cav.*)

(*Humatos*, humid, in opposition to *Adiantum*.)

Sori intra- or sub-marginal, globose, indusium coriaceous suborbicular or reniform, attached by a broad base, the apex and sides free ; fronds articulated upon the rhizome, coriaceous, small usually deltoid, more or less dimorphous, the barren ones less cut ; veins always free, rhizome creeping, growing on trees or rocks.

1. HUMATA HETEROPHYLLA. (*Smith.*) Rhizome wide-creeping, scaly ; frond shortly stalked, 3–6 inches long, 1 inch broad, glabrous ; the sterile ones ovate-lanceolate entire or slightly lobed at the base, the fertile ones narrower, deeply sinuate-pinnatifid ; sori 2–10 to a lobe. *Hook. Syn. Fil. 88. Bedd. F. B. I. t. 100.*

Singapore, Penang, and the Malay Peninsula.

(Also in the Polynesian Islands.)

2. *HUMATA ANGUSTATA*. (*Wall.*) Rhizome wide-creeping, scaly; fronds subsessile or shortly stalked, 3-8 inches long, $\frac{1}{4}$ - $\frac{1}{2}$ inch broad, linear, slightly and irregularly crenate at the margin, rarely once forked; both surfaces naked; veins parallel, simple or forked thickened; sori in a row along the edges, indusium small, transversely oval. *Hook. Syn. Fil. p.* 88. *Bedd. F. B. I. t.* 237.

The Malay Peninsula, Malacca, &c.
(Also in the Malay Islands.)

3. *HUMATA PARALLELA*. (*Wall.*) Caudex creeping, paleaceous; fronds, 4-8 inches long, $1\frac{1}{2}$ -2 inches broad, stipitate, coriaceous, ovate-lanceolate, acuminate, deeply pinnatifid nearly to the rachis; segments close, parallel, horizontally patent, linear or linear-oblong subfalcate entire, the lowermost pair sometimes with a solitary obtuse lobe at the base beneath, rarely more; involucres semiorbicular copious marginal, but all pointing towards the apex of the segments in two close parallel lines (not pointing towards the margins); veins thickened sunk. *Hook. Syn. Fil.* 89. *Bedd. F. B. I.* t. 99.

Birma and the Malay Peninsula.
(Also in the Polynesian Islands.)



HUMATA PARALLELA. (*Wall.*)

4. HUMATA PEDATA. (*Smith.*) Rhizome creeping, scaly ; stipe 2–4 inches long, fronds 2–8 long, by $1\frac{1}{2}$ –4 inches broad, deltoid in outline, cut down nearly or quite to the rachis, the lower pair of segments or pinnules larger than the others, and more compound, deeply pinnatifid, with the segments crenated, or more or less pinnatifid ; sori in rows on the teeth on both sides of the lobes. *Smith, Tentam Gen. Fil.* 15. *Hook. Syn. Fil.* p. 89. *Bedd. F. S. I. t.* 12. *H. alpina*, *Blume*, is a smaller, more finely cut variety, which is found in the Himalayas as well as Mergui. *H. vestita* (Bl.), *Bedd. F. S. I. t.* 253, has a longer stipe, is more lanceolate in shape, and more finely cut, but is said to run into typical “pedata.”

The typical form is found on the western slopes of the Nilgiris, and on the Travancore Ghats, at 3,000–4,000 feet elevation ; Sikkim, Bhotan, Khasya, Jaintea, 4,000 feet ; Ceylon ; and the Malay Peninsula.

(Also in the Malay Islands, North Australia, South China, in Japan, and the Mascareen Islands.)

The variety *alpina* has been received from the north west Himalayas (Dyas), and Mergui, and also inhabits Java, Azores, and the Polynesian Islands.

The variety *vestita* is from Ceylon, and also occurs in Java.

GENUS XV.—LEUCOSTEGIA. (*Presl.*)

(From *leucos*, white ; *stegos*, a cover.)

Sori intra- or sub-marginal, indusium as in *Humata*, but smaller, narrower and thinner ; fronds articulated upon the rhizome (except in *nodosa*), generally membranaceous and flaccid, generally 3–4-pinnatifid or pinnate, rarely only bipinnatifid ; veins forked, venules free ; rhizome long, creeping, growing generally on trees and rocks.

1. LEUCOSTEGIA HYMENOPHYLLA (*Parish MS.*) Caudex creeping, furnished with numerous long wiry roots, stipe slender, 3–4 inches long ; fronds subglabrous, membranaceous, very flaccid and transparent, ovate, or deltoid-ovate pinnate ; lower pinnæ deeply pinnatifid with the lower basal segments much the largest, and again



48 C. VERA ACYRHOPEA.

HUMATA PEDATA. (Sim.)

pinnatifid with the segments entire or crenated, the remaining pinnæ pinnatifid with the segments crenated; sori at the apex of the superior veinlets of each segment, indusium as in the genus, veins terminating within the margin. *Hook. Syn. Fil. p. 90. Bedd. F. B. I. t. 96.*

This very delicate species was discovered near Moulmein, on limestone rocks, by the Rev. C. S. Parish.

Moulmein.

2. *LEUCOSTEGIA MEMBRANULOSA.* (*Wall.*) Caudex hispid, with very long, slender, subulate, rigid, membranaceous scales; fronds 9–15 inches long, thin and membranaceous ovate-lanceolate and as well as the slender stipe and rachis pubescent-hirsute, bipinnatifid; pinnæ lanceolate, their rachis winged; pinnules lanceolate pinnatifid, the segments ovate-lanceolate subfalcate, very acute entire or rarely toothed; involucre small, ovate-subrotund acute, very thin and membranaceous fixed by the broad base, the rest free. *Hook. Syn. Fil. p. 91. Bedd. F. B. I. t. 98.*

Very like *multidentata*, but differs in the scales of the rhizome, and in the fronds being much less compound; in this they are pinnate, with the pinnæ deeply pinnatifid, in *multidentata* bipinnate, with the pinnæ deeply pinnatifid.

Nepal, Kumaon, Moharguri Pass, 6,500 feet.

(Also in Yunan.)

3. *LEUCOSTEGIA MULTIDENTATA.* (*Wall. under Aspidium.*) Rhizome stout-creeping; scales ovate-acute; fronds with the stipe up to 2 feet long and 1 foot broad, deltoid, bipinnate, with the pinnules deeply pinnatifid, down to a winged rachis, texture herbaceous; rachises pubescent above, often furnished with a few distant large ovate scales, rachises and segments beneath, generally glandularly pubescent; sori 2–12 to a segment, placed at the base of its teeth, on the upper side of the central vein. *Wall. Cat. 346. Microlepia pteropus. Bedd. F. B. I. t. 313. Hook. Syn. Fil. p. 91.*

Himalayas, Nepal and Bhotan, 5,000–8,000 feet; Khasya, 4,000–5,000 feet, common.

4. *LEUCOSTEGIA ASSAMICA*. (*Bedd. under Acrophorus*.) Rhizome creeping, covered with numerous imbricated chaffy golden, ovate-acute scales; fronds glabrous, shining, 8–14 inches long (including the stipe, which is 2–3 inches), 2–3 inches broad, subcoriaceous lanceolate, pinnate with the primary pinnæ alternate or opposite, pinnatifid almost to the rachis, lower ones petiolate, upper ones decurrent so as to form a regular wing to the rachis, pinnules pinnatifid, segments toothed, each bearing one sorus, veins terminating within the margin; involucres as in the genus, as broad as deep. *Bedd. F. B. I. t. 94. Hook. Sp. Fil. p. 467.*

Bhotan, Mishmee, and by the Dihong.



LEUCOSTEGIA ASSAMICA. (*Bedd.*)

5. *LEUCOSTEGIA IMMERSA*. (*Wall.*) Rhizome creeping stout, fibrillose; stipe 4–8 inches long, strong, erect; fronds 12–18 inches long, 6–9 inches broad, deltoid, tripinnate, glabrous; lowest pinnule lanceolate and deltoid, 2–3 inches long, 1 inch broad, with broad segments, which are obliquely truncate at the base below, and roundly lobed with the lobes again crenate above; texture herbaceous; sori large, impressed, clearly visible from the upper side, 1–6 to a segment, oblique; involucre broader than long. *Hook. Syn. Fil. p. 91. Bedd. F. S. I. t. 11.*

Madras Presidency on the Western mountains; very abundant in Coorg, growing on trees. Himalayas, from Mussorrie to Bhotan, 3,000–6,000 feet. Sikkim, Khasya, Parasnath, and the Malay Peninsula.

(Also in Java.)

6. *LEUCOSTEGIA HOOKERI*. (*Moore, under Acrophorus.*) Rhizome stout ; scales dense, lanceolate, golden ; stipe up to 6 inches long slender often scaly at the base ; fronds deltoid up to 10 inches long, 3-4 pinnatifid ; lower pinnæ opposite or alternate, lanceolate or deltoid, 1-2 inches broad ; pinnules deltoid, ultimate segments ligulate, acute, 1-veined, $\frac{1}{2}$ -1 line broad texture ; membranaceous ; sori at the base of the ultimate lobes ; involucre persistent, membranaceous, broader than long. *Davallia Clarkii*, *Baker. Syn. Fil. p. 91.*

Himalayas, Lachen, and Sirmur, 8,000-12,000 feet elevation. Sundukphoo, near Darjeeling, 11,000 feet.

Very like *Polypodium dareæforme*, Hook., but with acute segments, and differing somewhat in the scales of the rhizome. Indian botanists, not having seen this plant, have quoted *Hookeri* of Moore (*Clarkii* of Baker) as a synonym of *dareæforme*, and united the two under the name of *Leucostegia dareæformis*. Mr. Levinge has lately gathered this species at Sundukphoo, but the Kew specimens from Lachen and Sirmur were gathered by Thomson in 1849. I have examined a great many specimens of *dareæforme* in the Kew Herbarium, and can find no trace of an indusium, so I retain it in *Polypodium* ; but it is very likely that there is a fugacious indusium, and that it is a *Leucostegia* nearly allied to this species.

7. *LEUCOSTEGIA PULCHRA*. (*Don. under Davallia.*) Rhizome wide-creeping, clothed with broad-obtuse, often peltately attached scales ; stipe 4-6 inches long, naked ; fronds 9-15 inches long, 4-8 inches broad, lanceolate-deltoid, 3-4 pinnatifid, lowest pinnule 1-2 inches long, 1 inch broad, cut down to a narrowly-winged rachis into deeply pinnatifid segments, ultimate lobes narrow lanceolate, not distant, not very acute ; texture thin, rather flaccid ; sori copious, usually as broad as the segment at the base of the teeth of which it is placed ; involucre prominent. *Don. Prod. Fl. Nep. 11. Bedd. F. S. I. t. 10* (under *Acrophorus*).

Madras Presidency, Western mountains, very common on rocks and trees ; Ceylon, central provinces, 3,000-5,000 feet ; Himalayas, Nepal, and Bhotan, 2,000-9,000 feet ; Khasya ; Moulmein.



LEUCOSTEGIA PULCHRA. (Don).

8. *LEUCOSTEGIA PSEUDO-CYSTOPTERIS*. (*Kunze.*) Very similar to *pulchra*, but the ultimate segments very acute, and the scales of the rhizome more spreading. *Kunze in Bot. Zeit.* 1850, p. 68. *Bedd. F. B. I. t.* 92.

Himalayas, Dalhousie to Nepal, 4,000–8,000 feet.

Mr. Clarke considers it is a variety only of *pulchra*.

9. *LEUCOSTEGIA HYMENOPHYLLOIDES*. (*Blume, under Davallia.*) Caudex creeping, thick, clothed with long narrow subulate scales; fronds ample, tall, 12–24 inches long, ovate-lanceolate, membranaceous, 3–4 pinnate or supra-decompound; primary pinnæ petiolate, ovate-lanceolate acuminate, secondary petiolate oblong-ovate; pinnules ovate, deeply pinnatifid; the segments ovate acute, subfalcate, entire or generally (the fertile ones) with a tooth on the inner margin; involucres small hemispherical or subreniform, placed near the centre of a segment below the sinus of the tooth; veins slender, black. *Davallia affinis*. *Hook. Syn. Fil. p.* 92. *Bedd. F. S. I. t.* 252.

Travancore Mountains, Ceylon, 3,000–4,000 feet.

(Also in Java and Polynesia.)


10. *LEUCOSTEGIA PARVULA*. (*Wallich.*) Caudex long, creeping, clothed with lax subulate scales; frond very small deltoid-tripinnatifid, glabrous, rigid (from the stout costa), segments linear throughout, slightly grooved above when dry, unequally forked and acute at the apices; sori at the sinus of the forks; involucres suborbicular, dilated above, and broader than the segments. *Davallia parvula*, *Hook. Syn. Fil. p.* 92. *Bedd. F. B. I. t.* 97 (under *Acrophorus*).

Placed by Mettenius in *Humata*.

Singapore.

(Also in Borneo and Java.)

11. *LEUCOSTEGIA NODOSA* (*Presl. under Acrophorus.*) Fronds tripinnate, membranaceous furfuraceous on both sides of the veins; pinnules sessile (chaffy beneath at their insertions), oblong-lanceolate, secondary ones sessile, oblong obtuse, pinnatifid; segments cuneiform obtuse, lowest ones inciso-serrate or at the base again subauriculate



LEUCOSTEGIA NODOSA. (Presl.)

sori solitary submarginal; rachis nodose above at the insertion of the pinnæ and ferrugineo-tomentose; stipe not articulate, slightly rough or glabrous, paleaceous below; rhizome short-creeping. *Fresl. Tent. Pter. t. 3. Hook. Syn. Fil. p. 92. Bedd. F. B. I. t. 93.*

Himalayas, from Nepal to Bhotan, 3,000–7,000 feet, abundant in Sikkim; Khasya, 3,000–6,000 feet; Malacca.

(Also in Java.)

This differs from the other *Leucostegias* in habit, and in the stipe not being articulate, and should be kept distinct as a genus under its original name of *Acrophorus*; it is very like *Diacalpe aspidioides* except in the indusium.

GENUS XVI.—PROSAPTIA. (*Fresl.*)

(Sori immersed in the margin of the frond.)

Fronds contiguous, linear-lanceolate, pectinate-pinnatifid; veins simple, free; involucre a marginal cyst formed of the substance of the frond subcylindric.



Nº27.

PROSAPTIA EMERSONI. (*Fresl.*)

1. PROSAPTIA EMERSONI.

(*Fresl.*) Fronds tufted, sessile, 6–12 inches long, 1 inch broad, linear-lanceolate, cut more than half-way down to the rachis into many linear-oblong, or at the lower part triangular lobes; texture coriaceous; sori 1–6 round the edge of the lobes. *Hook. Syn. Fil. p. 94. Bedd. F. S. I. t. 20.*

Anamallay Mountains, South India, 3,000–4,000 feet; Ceylon; Penang.

(Also in Java, Borneo, and the

Philippine Islands)

2. PROSAPTIA CONTIGUA. (*Swartz.*) Fronds tufted, sessile, or nearly so, 12–18 inches long, 1–1½ inch broad, linear-lanceolate, cut



PROSAPTIA CONTIGUA. (Sw.)

down nearly or quite to the rachis into numerous linear-acuminate or bluntish slightly-toothed lobes; texture coriaceous; sori 2-8 to a lobe, placed in the teeth on their upper part. *Sw. Syn. Fil. p. 130. Hook. Syn. Fil. p. 94. Bedd. F. S. I. t. 19.*


Anamallay Mountains and Travancore Hills, on the west side of the Madras Presidency, not common; Ceylon, central provinces, 5,000 feet.

(Also in Java and the Philippines.)

GENUS XVII.—DAVALLIA. (*Smith.*)

(In honour of Davall, a Swiss botanist.)

Fronds pinnate, bipinnate, or deltoid-multifid, firm and glossy; veins forked, venules free; sori intra- or submarginal; involucres ter-



N° 29.

DAVALLIA TRIPHYLLA. (*Hook.*)

minial on the veins, attached by the base and sides, and forming an urceolate tabular cyst impressed in the substance of the frond; rhizome creeping or suberect; stipe articulated upon the rhizome.

1. DAVALLIA TRIPHYLLA. (*Hook.*)

Caudex stout, creeping, covered with chaffy scales; fronds coriaceous small ternate, pinnæ oblong-lanceolate, obtuse cuneate at the base, in fertile plants more elongated, all of them entire, intermediate ones petiolate, lateral ones shorter, sessile, oblique at the base, veins horizontally patent, copious, crowded, parallel, forked, thickened, flat (not prominent); involucres semicylindrical, compressed,

Singapore.

2. *DAVALLIA SOLIDA*. (Swartz.) Tall, caudex stout, creeping, clothed with densely imbricated scales, fronds coriaceous trisubquadripinnate, pinnæ acuminate; pinnules trapeziform acuminate, pinnatifid, many-veined, terminal ones crenate-serrate coadunate into an acuminate point; involucres linear-oblong, sunk in a tooth or segment having a narrow wing on each side or entire. *Hook. Sp. Fil.* i 163. *Sw. Syn. Fil.* pp. 132 and 345. *Hook. Syn. Fil.* p. 95. *Bedd. F. B. I. t.* 104. The width of the pinnules and the amount of cutting differ considerably.

Davallia ornata (Wall.) is a variety with the pinnules very wide.

Birma; Penang; and Singapore.

(Also in Java and the Polynesian Islands.)

3. *DAVALLIA ELEGANS*. (Sw.) Rhizome stout, creeping, clothed with woolly fibres; stipe firm, erect, 4-8 inches long; fronds 1-2 feet long, 9-15 inches broad, deltoid-tripinnatifid; main rachis slightly winged towards the apex; pinnules of the lower pinnæ 2-3 inches long, 1 inch wide, deltoid-lanceolate, cut down quite to the rachis in the lower part, with oblong-deltoid segments, which are slightly toothed and obliquely truncate at the base on the lower side; texture coriaceous; venation close, prominent, irregular, many spurious venules between the veins proper; sori several to a segment, marginal, but the sharp teeth projecting beyond them at the edges; involucre half cup-shaped. *Hook. Syn. Fil.* p. 95. *Bedd. F. S. I. t.* 18.

Western mountains of Madras; Ceylon, 2,000-3,000 feet elevation; Malay Peninsula.



N° 30.

DAVALLIA SOLIDA. (Sw.)

(Also in China, Java, Borneo, Tropical Australia, Polynesian Islands, Madagascar, Ango'a, Fernando Po, and Johanna Island.)

4. *DAVALLIA EPIPHYLLA*. (*Blume.*) Rhizome thick, fibrillose; stipe 4–6 inches long, erect, firm; fronds 12–15 inches long, 6–9 inches broad, deltoid-lanceolate, tripinnatifid; main rachis hardly at all winged; pinnules of the lowest pinnæ lanceolate, $1\frac{1}{4}$ inch long, $\frac{1}{2}$ inch broad; segments narrow, mucronate, sharply toothed; texture coriaceous; veins not immersed, one or two carried into each tooth; sori small, submarginal, half cup-shaped, with the sharp mucro of the tooth extending beyond them. *Hook. Syn. Fil. p. 96. Bedd. Suppl. to Ferns, t. 350.*

Malay Peninsula.

(Also in Java.)

5. *DAVALLIA DIVARICATA* (*Blume.*) Rhizome creeping, clothed with long lanceolate-caudate chestnut scales; stipe firm, erect, 6–12 inches long; fronds 2–3 feet long, tripinnatifid; lower pinnæ often 12 inches long by 6 inches broad; segments deltoid, cut down to the rachis in the lower part, with linear-oblong, sharply-toothed lobes; texture coriaceous; veins uniform, not conspicuous; sori half cup-shaped, placed obliquely as regards the central veins in the teeth at some distance from the edge; involucres as long as broad. *Hook. Syn. Fil. p. 96. D. polyantha* (Hook), *Bedd. F. B. I. t. 107.*

Sikkim, Mishmee, Khasya; Ma'ay Peninsula.

(Also in South China and Java.)

6. *DAVALLIA GRIFFITHIANA*. (*Hook.*) Rhizome creeping, stout, clothed with long lanceolate-caudate white or yellowish scales; stipe erect, wiry, 4–6 inches long; fronds 9–12 inches long, 4–8 inches broad, deltoid 3–4-pinnatifid; pinnules of the lower pinnæ lanceolate-deltoid, 2–3 inches long, 1 inch or more broad; lower segments toothed on the barren fronds, cut down nearly to the rachis in the fertile; texture coriaceous; sori large, submarginal or marginal; involucres cup-shaped, very shortly attached on the sides. *Hook. Syn. Fil. p. 96.*

Bedd. F. B. I. t. 106. The texture and habit of *Davallia*, but the indusium is nearer that of *Leucostegia*.

Bhotan and Mishmee, Khasya and Jaintea, 3,000–5,000 feet.
(Also in South China.)

7. *DAVALLIA LORRAINEI*. (*Hance*.) Rhizome thick as a quill; scales linear-subulate, nearly black, densely grey ciliated; stipe 3–4 inches long, naked brownish; fronds $\frac{1}{2}$ –1 foot long, deltoid, 4-pinnatifid; pinnæ stalked deltoid, lowest largest produced on the lower side, their rachises winged to base; pinnules and segments subsessile, crowded deltoid much reduced on lower side; final lobes ligulate, $\frac{1}{4}$ – $\frac{1}{3}$ lines broad, with sorus at base of inner side; texture subcoriaceous; surfaces naked; barren lobes 1-veined; involucre subcylindrical, $\frac{1}{2}$ line long. *Hance. Ann. Sc. Nat. Ser. V. vol. v. p.* 254. *Hook. Syn. Fil. p.* 469. *Bedd. F. B. I. Suppt. p. 4. t.* 351.

Malay Peninsula.

8. *DAVALLIA BULLATA*. (*Wall.*) Rhizome creeping, stout, densely clothed with hair-pointed chestnut scales; stipe strong, erect, 3–4 inches long; fronds 8–12 inches long, 4–8 inches broad, deltoid, 4-pinnatifid; pinnules of the lower pinnæ lanceolate, 2–3 inches long, 1 inch broad, with deeply inciso-pinnatifid oblong rhomboidal segments; texture coriaceous; sori deeply half cup-shaped, occupying the greater part of the tooth in which they are placed, marginal, with usually a horn on the outside. *Hook. Syn. Fil. 97.* *Bedd. F. S. I. t.* 17.

All the Western Ghats of Madras and Bombay Presidencies; Himalayas, Nepal to Bhotan, 2,000–6,000 feet; Khasya; Ceylon; Birma, and Malay Peninsula.

(Also in Japan, South China, and the Malay Islands.)

9. *DAVALLIA SPECIOSA*. (*Mett.*) Rhizome wide-creeping, 2 lines thick, scales ferruginous dense linear-subulate; stipe 3–5 inches, naked, brown, stramineous; frond deltoid, 4-pinnatifid, 1–1 $\frac{1}{4}$ foot long; rachis winged in upper half, pinnæ lanceolate-deltoid, long-stalked, lowest largest 3–4 inches broad; pinnules and segments


deltoid, with rachises winged to base, cuneate-truncate on lower side, final lobes lanceolate-falcate under 1 line broad, with sorus on upper side at base; texture subcoriaceous; surfaces naked; involucre cup-shaped, broader than deep, $\frac{1}{2}$ line broad, oblique, close to final sinuses. *Hook. Syn. Fil.* 469. *Bedd. F. B. I. Suppl. t.* 352.

Moulmein.

GENUS XVIII.—MICROLEPIA. (*Presl.*)

(*Micros*, small; *lepis*, a scale, the small indusium.)

Fronds pinnate or variously compound; texture various; veins simple or pinnately forked; venules free; sori intra- or sub-marginal; involucre membranaceous, half cup-shaped, attached at the sides as well as the base; rhizome creeping; stipe continuous with the caudex.



Nº 32

MICROLEPIA HOOKERIANA (*Wall.*)

1. MICROLEPIA HOOKERIANA. (*Wall.*) Stipe a foot and more long, pubescent, at length glabrous; rachis hirsute-pubescent; fronds lanceolate pinnate; pinnae subpetiolate lanceolate from a broad hastate base, gradually acuminate submembranaceous, duplicate-crenate, sparingly hairy on the costa and veins beneath; veins parallel, dichotomous; sori approximate, forming a continued line at the base of the crenatures

of the margin; involucre half cup-shaped; stipes and rachis pubescent-hirsute. *Hook. Syn. Fil.* p. 97. *Bedd. F. B. I. t.* 101.

Sylhet, Upper Assam, Khasya and Mikir Hills.

(Also in Hong Kong.)



MONTÉILAC-ENGRAIS.

DAVALLIA BULLATA. (Wall.)

2. *MICROLEPIA PINNATA*. (*Car.*) Rhizome creeping, furnished with fibrillose scales; stipe strong, erect, 6–12 inches long, glossy; fronds 9–15 inches long, 4–8 inches broad, lanceolate-pinnate, glabrous; pinnæ slightly toothed, 6 inches long, $\frac{1}{4}$ inch broad, coriaceous, linear-lanceolate, gradually acuminate, obliquely acuminate at the base; sori one to each tooth, small, submarginal; veins sunk, inconspicuous, generally forked; involucres small, half-cup-shaped. *Hook. Syn. Fil. p.* 98. *Bedd. F. S. I. t.* 14.

Anamallay Mountains; Malay Peninsula.

(Also in Java and Polynesian Islands.)

3. *MICROLEPIA MARGINALIS*. (*Thunb. under Polypodium.*) Fronds broadly ovate-lanceolate, firm, membranaceous; pinnæ elongate, lanceolate, subfalcate, acuminate, pinnatifid-lobate, the acuminate apices serrated, pubescent-villous beneath, most so on the costa and prominent veins, unequally cuneate at the base, and subpetiolate; lobes acute crenate-dentate; veins pinnated; sori solitary in the axils of the smaller and upper lobes or serratures, and distant from the margin, marginal on the small teeth of the larger lobes; involucres broad half-cup-shaped, densely villous; rachis and stipe downy, the latter at length glabrous; rhizome creeping, villous; stipe 1–2 feet long, erect, strong; fronds 18–24 inches long, 9–15 inches broad, once pinnate. *Polyp. marginale, Thunb. Fl. Japan, p.* 337. *Microlepia scabra, Don. Bedd. F. B. I. t.* 102.

Nepal and Kumaon, Khasya, Mikir Hills.

(Also in Japan and Formosa.)

VAR. β *CALVESCENS*. (*Hook.*) Pinnæ narrower and more deeply pinnatifid, nearly glabrous beneath, except the strigose midrib. *Davallia calvescens. Hook. Sp. Fil. I. p.* 172, *t.* 48 B. *D. urophylla* (Wallich), *Bedd. F. B. I. t.* 103.

Kumaon.

4. *MICROLEPIA UROPHYLLA*. (*Hook.*) Rhizome creeping; stipe strong, erect, 2–3 feet long; fronds bipinnate, tripinnatifid, coriaceous, shining above and beneath, but pubescent on the rachises below;



LA CATTALACEAE

MICROLEPIA PLATYPHYLLA. (Don.)

secondary pinnæ lanceolate-linear, very finely caudate, cut down nearly to the rachis into ovate acuminate, unequal sided toothed lobes; sori submarginal, one to each of the lower sinuses of the lobes. *Hook. Syn. Fil. p. 99, not of Wallich or Bedd. Clarke, F. N. I. t. 50.*

Bhotan (allied to *M. platyphylla*).

5. *MICROLEPIA PLATYPHYLLA*. (*Don.*) Rhizome creeping, stout, scaly; stipes 2-3 feet long, firm erect; frond 3-4 feet long, tripinnatifid; lower pinnæ 12-15 inches long, 6-9 inches broad, lanceolate with distant linear-lanceolate pinnules, which are cut nearly to the wavy rachis below, into broad, bluntish toothed, oblong-deltoid lobes; texture coriaceous when adult, glabrous and shining on both surfaces, but when young more or less membranaceous and hairy; sori 2-12 to a segment, placed one in each tooth, a short distance from the edge, about a line across. *Don. Fl. Prod. Nep. 10. Hook. Syn. Fil. p. 99. Bedd. F. S. I. t. 13.*

Madras Presidency, throughout the Western mountains, up to nearly 6,000 feet; Ceylon; Himalayas, Nepal, Sikkim, Bhotan 3,000-5,500 feet elevation; Khasya.

6. *MICROLEPIA KURZII*. (*Clarke.*) Fronds large tripinnatifid; pinnæ as in *platyphylla*, with linear-lanceolate pinnules, which are cut down to its rachis (which is not wavy, or only slightly so towards the apex) below into triangular subobtuse or lanceolate-caudate lobes, which are bluntly serrate; texture not so coriaceous as in *platyphylla*, uniformly pubescent beneath, even when adult; venation much less prominent than in *platyphylla*; involucres finely ciliated round the margin. *Clarke F. N. I. p. 446.*

Birma.

This is very like *platyphylla*, and may turn out to be only a variety of it. I have only seen one specimen.

7. *MICROLEPIA MAJUSCULA*. (*Lowe.*) Rhizome creeping, tomentose; stipe 6-12 inches long, erect naked; fronds up to 3 feet long lanceolate-deltoid, tripinnatifid; lower pinnæ 9-12 inches long, 3-4 inches broad, lanceolate; pinnules lanceolate-acuminate, cut down

nearly or quite to the rachis into blunt slightly crenated oblong lobes, rachis and both surfaces slightly hairy, the upper bright green, shining ; texture subcoriaceous ; sori small submarginal, 2-12 to a segment. *Hook. Syn. Fil. p. 99.* *Microlepia proxima* (Thw.), *Bedd. F. S. I. t. 254.* Ceylon, Rangbodde, 3,500 feet elevation.

8. *MICROLEPIA STRIGOSA.* (*Swartz.*) Fronds tall, lanceolate, bipinnate ; stipes elongated ; rachis and veins pubescent-hispid, primary pinnæ petiolate, lanceolate-acuminate, secondary (or pinnules) mostly petiolate, subdimidiate-ovate, obtuse pinnatifid, chiefly on the upper edge, lower lobes obovate deep, the rest short, all of them angulate-dentate, veins pinnated, furnished with a few long scattered hairs both above and beneath (the remaining surface of the frond beneath being sometimes furnished with numerous small hairs, or sometimes glabrous as is the upper surface) ; involucre hairy, small, half cup-shaped. *Hook. Syn. Fil. p. 98.* *Bedd. F. S. I. t. 255.*

Tinnevelly and Travancore Mountains, South India ; Ceylon ; Himalayas ; and Malay Peninsula.

(Also in Japan, South China, Sandwich and Fiji Islands.)

Mr. Clarke considers this a variety, or rather only a young state of *speluncæ*, as he states it develops into this more compound form ; as far as the South Indian and Ceylon forms are concerned, this is never more than bipinnate, whereas *speluncæ* is 3-4 pinnate ; it has been for years in cultivation in ferneries, at Ootacamund, and is quite constant.

9. *MICROLEPIA SPELUNCÆ.* (*Linn.*) Rhizome creeping ; stipes strong, 1-1½ foot long ; fronds up to 6 feet long, rarely more, and 2 feet broad, ovate to deltoid, 3-4-pinnatifid, more or less hairy, strigose or villous, or with few or many long glistening scale-like flaccid hairs, rarely sub-glabrous ; texture membranaceous, or flaccid, pinnules from oblong or ovate to linear-lanceolate, ultimate segments entire or subentire and rhomboid, or irregularly inciso-obate or pinnatifid ; sori large 1-5 to the entire segments, more copious on the lobed segments ; involucre half cup-shaped, hispid or rarely glabrous ; veins more or less prominent beneath. *Polypodium speluncæ*, *Lin. Sp. Pl. 1555.*

The Himalayas from Kumaon eastwards ; Khasya ; Chittagong ; Madras, Western mountains ; Ceylon ; Malay Peninsula.

(Also in China, Japan, Malay Islands, Polynesia, and Tropical America.)

TYPE. Ultimate segments rhomboidal, subentire, or slightly crenated. *Davallia rhomboidea*, *Wall. Cat.* 257.

VAR. β HIRTA. Ultimate segments more or less deeply cut.


Rhomboidea (*Microlepia polypodioides*, *Bedd. F. S. I. t.* 15) can easily be picked out in the herbarium : it is very uniform in character, and does not run into *hirta* in cultivation ; it is very common throughout India and in Ceylon.

The variety *hirta* (*Bedd. F. S. I.* 256) is a much more variable plant, and I refer to this *Wall. Cat.* 262 *puberula* ; 263 *pilosula* ; 264 *virens* ; 2,218 *Roxburghii*, and 261 *pyramidata*. These are all so closely allied that it is impossible to keep them up as separate varieties, and they besides run one into the other. There are two varieties in Southern India and Ceylon : one very hairy and the other nearly quite glabrous, but differing in no way beyond the pubescence. The supposed species, *flaccida* (*Bedd. F. Sup. t.* 353) I now consider only a form of *hirta*, as I cannot distinguish it in the herbarium, it is said to be very flaccid, and to have fronds ten and twelve feet long. I have seen no forms anything like so large in Southern India or Ceylon, yet some Ceylon and South Indian specimens are referred to it. Wallich's *pyramidata* has the pinnules closer together and more elongated than any of the other forms, but otherwise it is quite like *hirta*.

GENUS XIX.—STENOLOMA. (*Fée.*)

(*Stenos*, narrow ; *loma*, border.)

Fronds bi-tripinnatifid, ultimate segments cuneiform, growing gradually wider from the base to the apex ; veins dichotomously forked, venules free ; indusium terminal on the segments, forming a compressed suborbicular, or cup-shaped pouch, only open at the top ; rhizome creeping ; stipes tufted, not articulated upon the rhizome.



8. CASTELLANERAS.

STENOLOMA CHINENSIS. (Sw.)

1. *STENOLOMA CHINENSIS*. (*Swartz.*) Rhizome stout, densely fibrillose; stipes strong, erect, polished, naked, dark brown, 6–12 inches long; fronds 12–18 inches long, 6–9 inches broad, ovate, 4-pinnatifid; lower pinnæ ovate-lanceolate, 4–6 inches long, 2–3 inches broad; pinnules lanceolate, their segments cut down to the rachis below, with toothed cuneate lobes, 1–1½ lines across at the apex; texture subcoriaceous, both surfaces naked, the upper shining; sori terminal, usually solitary, often rather broader than deep. *Swartz Syn. Fil.* 133 (under *Davallia*) *Davallia tenuifolia*, *Hook. Syn. Fil. p.* 102. *Bedd. F. S. I. t.* 16.

Madras Presidency, Western mountains, 3,000–6,000 feet: Himalayas, Kumaon to Bhotan, up to 1,000–4,000 feet; Khasya; Ceylon; Malay Peninsula.

(Also in China, Japan, Polynesia, and the East African Islands.)

GENUS XX.—CYSTOPTERIS. (*Bernh.*)

(*Cystos*, a cyst, in allusion to the inflated indusium.)


Fronds more or less compound, membranaceous; sori globose, placed on the back of the veins; indusium membranaceous, suborbicular inserted by its broad base, under the sorus, which, at the beginning it covers or partially covers like a hood; veins free.

1. *CYSTOPTERIS FRAGILIS*. (*Bernh.*) Stipes 2–12 inches long; fronds glabrous, weak, up to 1 foot, ovate-lanceolate, tripinnatifid; main rachis slightly winged above; pinnæ lanceolate-deltoid; pinnules oblong-rhomboidal, cut down to a broad central space into bluntly or sharply-toothed segments; sori 2–12 to a pinnule. *Hook, Syn. Fil.* 103. *Bedd. F. B. I. t.* 91.

North West Himalayas, from Kashmir to Kumaon, 10,000–15,000 feet elevation; Sikkim.

(Also in most parts of the world, in cold regions.)

2. *CYSTOPTERIS SETOSA*. (*Bedd.*) Caudex erect; stipes 1 foot long, sparsely covered with small flaccid subulate light-brown scales; fronds tufted 3–3½ feet long, tripinnate, pinnæ 8–10 inches long, lowest



A. J. CATTY-INGENIENSIS.

CYSTOPTERIS SETOSA. (Bedd.)

pair nearly as long as the central ones, gradually decreasing in size towards the apex, pinnules pinnatifid to nearly the rachis, but always more or less connected by a decurrent wing, very membranaceous and flaccid, and furnished on both sides sparsely (as is the rachis and costa) with long weak pellucid jointed setæ; segments of the pinnules generally as broad at the apex as at the base, more or less pinnatifid, with the lobes obtusely rounded or variously toothed; veins forked and simple reaching the margin; sori one to each segment, medial on the lower vein; involucre very membranaceous, small scale-like, ovate, roundish ovate, or oblong, from a broad base fornicate, not nearly covering the sorus, fimbriate or entire, fugacious. *Bedd. F. B. I. t. 312*; *also t. 262*, under *Lastrea*.

Moulmein Mountains; Sikkim 5,000–8000 feet.

TRIBE V.—LINDSAYEÆ.

Sori placed in a line, at or very near the edge of the frond, covered with an involucre, the inner valve of which is membranaceous, the outer formed of the margin of the frond.

GENUS XXI.—LINDSAYA. (*Dryand.*)

(In honour of Dr. Lindsay, a writer on Ferns.)

Sori marginal or submarginal, placed to the apex of, and uniting two or more veins; involucre double, opening outwardly, the inner valve membranaceous, the outer formed of the more or less changed margin of the frond; veins free; pinnæ unilateral or equilateral.

§ *Pinnæ unilateral.*

1. *LINDSAYA CULTRATA*. (*Swartz.*) Rhizome wiry, creeping, furnished with linear scales, stipes wiry flexuous, 3–6 inches long; fronds 6–12 inches long, about 1 inch broad, simply pinnate, pinnæ unequal-sided, lower edge nearly straight near the main nerve; upper edge slightly lobed or nearly entire; lower pinnæ stalked; texture rather coriaceous. *Swartz. Syn. Fil. 119. Hook. Syn. Fil. 105. Bedd. F. S. I. t. 23.* *Lindsaya Lobbiana* (Hook.), *Bedd. F. B. I. t. 28.* A variety with rounded pinnæ, and a 4-winged rachis.




AL. LINDSAY & CO. EDINBURGH

LINDSAYA CULTRATA. (Sw.)

Very common on the western side of the Madras Presidency, up to 6,000 feet, called the hay-scented fern, from its scent when drying. Ceylon, higher altitudes in the central provinces. Himalayas, Nepal, to Mishmee and Chittagong, up to 4,000 feet; Birma and the Malay Peninsula.

(Also in North Australia, Formosa, Japan, Malay Islands, and East African Islands.)

2. *LINDSAYA REPENS*. (*Thw.*) Caudex creeping, scaly; stipæ short; fronds rigid, membranaceous linear-lanceolate, 10–18 inches long, $\frac{1}{2}$ inch broad, pinnate attenuated at the base; pinnæ very numerous, 40 or more pair, half deltoid-ovate obtuse or sub-acute,



No 37.

LINDSAYA SCANDENS.
(*Hook.*)

base nearly parallel with the rachis, and with the lower margin quite entire, upper margin lobulate-crenate, costa parallel with and close to the margin; veins simple or forked, free; sori short, oblong, one near the margin of each lobule.

Thw. En. Pl. 388. *Bedd. F. S. I. t.* 209. *Odontoloma*, *Hook. Syn. Fil.* p. 93. *Lindsaya pectinata* (Blume), *Hook. Syn. Fil.* p. 106.

Ceylon, Singhe-Rajah Forest; Mishmee, Khasya, Sikkim-Terai, Dulkajhar 1,000 feet; Malay Peninsula.

Also in the Malay Islands, Polynesia and Mauritius.)

VAR. β MINOR. A smaller form, more membranaceous, and more deeply pinnatifid; sori much shorter. *Thw. l. c.* *Bedd. F. S. I. t.* 214.

3. *LINDSAYA SCANDENS*. (*Hook.*) Rhizome stout, wide-creeping, scandent, paleaceous; frond 9–12 inches long, $1\frac{1}{2}$ – $1\frac{3}{4}$ inch broad, simply pinnate; pinnæ $\frac{3}{4}$ inch long, $\frac{1}{2}$ inch broad, the lower line slightly decurved, the upper rounded, entire, the point broadly rounded, placed in a long row close together, but not imbricated; texture pellucid-herbaceous; costa marginal; sori in a continuous marginal line. *Hook. Sp. Fil. i. p.* 205, t. 63 B.; *Syn. Fil. p.* 106. *Bedd, F. B. I. t.* 298.

The Malay Peninsula.

(Also in the Philippine Islands.)

4. *LINDSAYA ORBICULATA*. (*Lam. under Adiantum.*) Rhizome creeping; stipes tufted, generally elongated, fronds linear-lanceolate and pinnate or deltoid and bipinnate; pinnules approximate shortly petiolate, rather rigid flabellate, and approaching to lunate or sub-rhomboid, with the sides unequal, the base obliquely cuneate, sometimes the upper ones are confluent, the superior margin crenulate or soriferous; no distinct midrib; sori continuous or interrupted; involucre toothed. *Hook. Syn. Fil.* i. 211. *L. flabellulata*, *Hook. Syn. Fil.* p. 107. *Bedd. F. B. I. t.* 216.

Davallia trichomanoides, *Bedd. F. B. I. t.* 178 (not Blume) an abnormally cut variety. *D. schizophylla*, *Hook. Syn. Fil.* p. 468.

VAR. β *TENERA*. Pinnules of thinner texture, and obtusely rounded in outline. *Bedd. F. S. I. t.* 24. This is the only form found in the Madras Presidency.

Western Forests of Madras; Ceylon; Khasya, Assam, Jaintea; Malay Peninsula.

(Also in South China and Australia.)

5. *LINDSAYA LANCEA*. (*L. under Adiantum.*) Rhizome creeping; stipes strong, up to 12 inches long; frond up to 15 inches long, bipinnate; pinnæ, several pairs erect-patent; pinnules about 1 inch long, by $\frac{1}{4}$ – $\frac{1}{2}$ inch deep, the lower line nearly straight, the upper rounded entire; texture pellucid-herbaceous; sori in a continuous line round the upper margin. *Adiantum Lanceum*. *L. Lindsæa trapeziformis* (Dry.), *Hook. Syn. Fil.* 107. *L. caudata* (Hook.), *Bedd. F. S. I. t.* 217. Mr. Wall sends from Ceylon a form with numerous pinnæ, which taper away into a long caudate apex, the lower pinnules being obovate from a longish stalk, and much smaller than in the type; the upper pinnules very small and tongue-shaped.

Ceylon; Malay Peninsula.

(Also in Tropical America, the West Indies, and the Malay Islands.)

6. *LINDSAYA RIGIDA*. (*J. Sm.*) Rhizome wide-creeping, stipes 4–6 inches long, rigid, erect, prickly towards the base; fronds with a long *unbranched central point*, and 1–4 pair of flexuose lateral


branches, 4-8 inches long; pinnules 3-4 lines broad, 2 lines deep, the lower edge often falcate, the upper 3-4 times bluntly, not deeply lobed, close together, but not imbricate; texture very thick, coriaceous; veins prominent; sori in a marginal line on the lobes.

Hook. Syn. Fil. p. 108. Bedd. F. B. I. t. 166.

Malacca, on Mount Ophir.

§§ *Pinnæ equilateral.*

7. *LINDSAYA WALKERÆ. (Hook.)* Rhizome creeping, thicker than a crow's quill, ferruginous with scale-like hairs. Stipes very long, often 1 foot, and as well as the rachis dark purple, glossy;



N°38.

LINDSAYA DIVERGENS.
(Wall.)

fronds lanceolate, pinnate, 6 inches to 1 long, pinnæ 6-9 pairs, with a terminal one, which is sometimes confluent with one or both of the upper pair, coriaceous, sub-opposite, remote, lanceolate or linear-lanceolate, equal, costa central, veins copious, almost parallel with the costa; sorus marginal, continuous on both sides.

Hook. Sp. Fil. i. 209. Syn. Fil. 109. Bedd. F. S. I. t. 215.

Ceylon, Hinidoon Corle in swamps.

(Also the island of Banca, east of Sumatra.)

8. *LINDSAYA DIVERGENS. (Wall.)* Rhizome creeping; stipes and rachis ebony-black, glossy; fronds lanceolate-pinnate; pinnæ approximate, crowded, horizontal-lanceolate, obtuse, subsemihastate at the base, glaucous beneath, the sides equal; costa central; veins oblique, once forked, distant, internal, obscure; sorus marginal and continuous on both edges and at the apex. *Hook. Sp. Fil. i. 210; Hook. et Grev. Ic. Fil. f. 226. Bedd. F. B. I. t. 250. Vittaria divergens, Herb. Roxb. Wall. Cat. n. 2191.*

Malacca.

(Also in Borneo.)

9. *LINDSAYA LANUGINOSA*. (*Wall.*) Rhizome creeping, stout, clothed with fibrillose scales; stipes stout, erect, 4–6 inches long; fronds 12–24 inches long, 3–4 inches broad, simply pinnate; pinnæ $1\frac{1}{2}$ –2 inches long, $\frac{1}{4}$ – $\frac{1}{2}$ inch broad, linear entire or very slightly toothed towards the point, which is acute in the fertile, bluntly rounded in the barren frond; texture coriaceous; rachis pubescent; sori in a continuous line along both edges. *Hook. Syn. Fil. p.* 110. *Bedd. F. B. I. t.* 140.

Birma and the Malay Peninsula.

(Also in Tropical Australia, Mauritius, and Tropical Africa.)

10. *LINDSAYA HETEROPHYLLA*. (*Bedd.*) Glabrous, fronds, 10–20 inches high; deltoid-bipinnate, in the lower portion, sometimes tripinnate, simply pinnate at the apex; lower pinnæ 4–6 inches long, upper ones gradually smaller and less compound, pinnules numerous, as many as 20, very variable in form, subrotund, rhomboidal or lanceolate, margin entire; veins simple or forked, all free; sori continuous round the whole margin except at the base, the larger pinnules have a distinct midrib. *Bedd. F. S. I. t.* 206.

This is more compound than any of the Indian forms of *Schizoloma heterophylla*; but there is a form of that plant from Hong-kong in the Kew Herbarium very like this in outline, but with the veins anastomosing. I have examined many fronds of this plant, and the veins never anastomose, and if it be lumped with *Schizoloma heterophylla*, it at once does away with that genus or subgenus. I am, however, inclined to consider it a distinct species more allied to *L. orbiculata* var. *tenera* than to *Schizoloma heterophylla*.


Shevaroy Hills (on the Green hills), Tinnevely Mountains.

GENUS XXII.—SCHIZOLOMA. (*Gaud.*)

(*Schizo*, I cut; *loma*, margin.)

As in *Lindsaya*, but veins more or less anastomosing.

1. *SCHIZOLOMA LOBATA*. (*Poir.*) Rhizome short-creeping; stipes 6–9 inches long, firm, erect; fronds pinnate; pinnæ 1–6 pairs, erect-patent, 3–6 inches long; pinnules about $\frac{1}{2}$ inch long, $\frac{1}{4}$ inch



SCHIZOLOMA LOBATA. (Poir.) VAR. MALABARICA.

broad, recurved, rounded, the upper margin with broad, shallow lobes, texture thin pellucid-herbaceous; veins anastomosing in the upper portion of the pinnules; sori marginal in the lobes, the inner valve of the involucre narrow and membranous, the edge of the frond produced beyond it, and scarcely altered. *Hook. Syn. Fil. p. 111.*
S. recurvata (Wall.), *Bedd. F. S. I. t. 27.* *L. nitens*, *Blume.*


Malabar and Travancore Mountains, Ceylon.

(Also in Queensland and the Polynesian Islands.)

VAR. β MALABARICA. (*Bedd.*) Fronds always simply pinnate; rachis tetragonous; pinnules erect, not recurved, upper margin more deeply lobed; veins less anastomosing. *Bedd. F. B. I. t. 268.*

Perhaps a distinct species.

Malabar and South Canara.



SCHIZOLOMA CORDATA.
 (*Gaud.*)

2. SCHIZOLOMA DAVALLIOIDES. (*Bl.*)

Rhizome short-creeping; stipes 6–12 inches long, firm, erect; fronds pinnate, pinnæ 2–3 pairs, erect-patent, 4–8 inches long, pinnules 4–6 lines long, 2–3 lines broad, the lower margin straight or slightly curved, the upper with 4–6 regular rounded but not deep lobes, texture pellucid-herbaceous; veins anastomosing at the base of the lobes; sori marginal in the lobes. *Hook. Syn. Fil. p. 111.* *Bedd. F. B. I. t. 141.*
 (Closely allied to lobata.)

Malacca.

(Also in the Malay Islands.)

3. SCHIZOLOMA CORDATA. (*Gaud.*) Rhizome short-creeping; stipe 3–6 inches long, slender, erect, wiry; barren frond 2–3 inches long, 1–1½ inch broad, cordate-oblong, quite entire, fertile one 3–6 inches long, linear entire or forked; texture coriaceous; sori in a continuous marginal line; veins anastomosing. *Hook. Sp. i. p. 219, t. 66 A.*; *Syn. Fil. p. 111.* *Bedd. F. B. I. t. 299.*

Malay Peninsula.

Malacca.

(Also in the Malay Islands.)


5. SCHIZOLOMA ENSIFOLIA. (*Swartz.*) Rhizomaleaceous; stipes 6–9 inches long, wiry, flexuous; fronds 6–9 inches long, 3–4 inches broad, with a linear-lanceolate pinnatifid apex, simply pinnate below, pinnæ here 10–12 pairs, rarely reduced to one, all stalked, $1\frac{1}{2}$ –2 inches broad, varying from linear-acuminate to lanceolate, only a little toothed; veins copiously anastomosing, with a continuous marginal line. *Hook. Syn. Fil. p. 112. B.*

(*Griffithiana* (Hook.), *Bedd. F. B. I. t. 29* is a simple

Western mountains of Madras; Ceylon; Himalayas; Munnepore and Chittagong up to 4,000 feet; Birm

(Also in North Australia, Tropical Africa and in Polynesia.)

6. SCHIZOLOMA HETEROPHYLLA. (*Dry.*) Rhizomaleaceous; stipes 4–8 inches long, firm, naked, erect; fronds 4–8 inches long, 3–6 inches broad, lanceolate or oblong, deltoid



SCHIZOLOMA ENSIFOLIA. (Sw.)

TRIBE VI.—PTERIDEÆ.


Sori marginal, oblong, or linear; indusium of the same shape as the sorus, formed of a more or less changed and reflexed portion of the frond, opening inwardly.

GENUS XXIII.—ADIANTUM.

(*Adiantos*, dry; water will not lie on the fronds.)

Sori marginal, varying in shape from globose to linear, usual numerous and distinct, sometimes confluent and continuous; indusium of the same shape as the sorus, formed of the reflexed margin of the fronds, bearing the capsules on its under side; veins free.

1. ADIANTUM PARISHII. (*Hook.*) Small, caespitose; no distinct



ADIANTUM PARISHII. (*Hook.*)

caudex; roots fibrous, tomentose; fronds about 1 inch each way, orbiculate flabellate, membranous, pellucid, sterile ones crenate dentate, fertile on few (3-5) lobate sinuses deep soriferous; veins originating from the base, flabellately-

convergent, repeatedly dichotomous; stipes slender, filiform, ebeneous black, shining, articulated at the summit. *Hook. Sp. Fil.* ii. 237, at *Fil. Exot.* i. pl. 51; *Syn. Fil.* p. 114. *Bedd. F. B. I.* t. 16.

Moulmein, on limestone rocks.

2. ADIANTUM LUNULATUM. (*Burm.*) Stipes 4-6 inches long, tufted, wiry, naked, polished dark chestnut-brown; fronds 6-1

inches long and 3 inches broad, simply pinnate, often elongated and rooting at the apex; pinnæ subdimidiate, the lower edge nearly in a line or oblique with the petiole, the upper edge rounded and like the bluntly-rounded sides usually more or less lobed; petioles of the lower ones spreading $\frac{1}{4}$ – $\frac{1}{2}$ inch long, texture herbaceous; the rachis and both surfaces naked; sori in continuous lines along the edge *Burm. Fl. Ind. p. 235. Hook. Syn. Fil. p. 114. Bedd. F. S. I. t. 1.*

Throughout Northern India in moist places; South India, very general on the western side in the plains and lower slopes of the hills; Ceylon; Birma.

(Also in the tropics of nearly the whole world.)

VAR. β METTENII. (*Kuhn.*)

Stipes and rachis winged, not polished, petioles winged, very short, otherwise as in *lunulatum* (type). *Kuhn, Fil. Af. p. 65. Bedd. Ferns Suppl. t. 354. A. pteropus, R. Br.*


Travancore Mountains, in dense evergreen forests between Courtallum and Quilon, 1,000–2,000 feet. Perhaps a distinct species.

(Also in Tropical Africa.)

3. ADIANTUM CAUDATUM. (*L.*)

Stipes 2–4 inches long, tufted, wiry, spreading, dark chestnut-brown, tomentose; fronds 6–12 inches long, simply pinnate, often elongated and

rooting at the extremity, pinnæ $\frac{1}{4}$ – $\frac{3}{4}$ inch long, $\frac{1}{2}$ inch deep, dimidiate, nearly sessile, the lower line straight and horizontal, the upper rounded, more or less cut, often deeply and repeatedly, the point usually blunt, the lower ones slightly stalked; texture coriaceous; the veins prominent; the rachis and both surfaces of the frond villose; sori roundish or transversely oblong on the edge of the lobes. *Hook. Syn. Fil. p. 115. Bedd. F. B. I. t. 2.*



No. 43.
ADIANTUM LUNULATUM.
VAR. METTENII. (*Kuhn.*)

Throughout India, Ceylon, and the Malay Peninsula, in the plains and on lower slopes of the hills.

(Also in South China, Tropical Africa, the Malay Islands, Java, Mauritius, and Cape Verde Islands.)

VAR. β EDGEWORTHII. Pinnæ and rachis glabrous; pinnæ less cut on the upper margin. *Adiantum Edgeworthii*. *Hook. Sp. Fil.* ii. 14. *Bedd. F. B. I. t.* 17.

Nepal, Mooltan, Gurwhal. Perhaps a distinct species.

Clarke refers this to *rhizophorum* Wall. Cat. 82, but that plant has the rachis scabrous above, and is one of the forms of typical *caudatum*.

VAR. γ SOBOLIFERUM. Stipes petioles and rachis winged. *Hook. Sp. Fil.* ii. 13. *Bedd. F. B. I. t.* 19.


Birma.

4. *ADIANTUM CAPILLUS VENERIS*. (L.) Stipes suberect, rather slender, 4–9 inches long, polished, blackish, naked; fronds bipinnate, with a short terminal pinna and numerous erect-patent lateral ones on each side, the lowest slightly branched again; segments $\frac{1}{2}$ –1 inch broad, the base cuneate, the outer edge rounded, deeply lobed from the circumference in the direction of the centre, and the lobes again bluntly crenated, lowest petioles $\frac{1}{4}$ inch long, texture pellucid-herbaceous, thin; rachis and both surfaces naked; sori roundish or obreniform, placed in the roundish sinuses of the crenations. *Hook. Syn. Fil.* p. 123. *Bedd. F. S. I. t.* 4.

Madras Presidency, west side, common on banks of rivers in the plains, and up to 5,000 feet on the mountains; Ceylon; North India, in many localities.

(Also in Europe, Africa, America, and Australia.)

5. *ADIANTUM ÆTHIOPICUM* (L.) Stipe 6–9 inches long, rather slender, erect, dark chestnut-brown, polished, naked; fronds up to 18 inches long, 6–9 inches broad, deltoid in outline, 3–4-pinnate; lower pinnules 3–4 inches long, 2–3 inches broad, deltoid; ultimate segments $\frac{1}{4}$ – $\frac{1}{2}$ inch across, suborbicular, straight or subcuneate or rounded at the base, the upper part broadly not deeply lobed; texture thinly pellucid-herbaceous; rachis and surfaces naked; sori in several roundish or transversely oblong patches in rounded hollows



ADIANTUM CAUDATUM. (Linn.)

of the outer edge. *Hook. Syn. Fil. p. 123. Bedd. F. S. I. t. 5.*
Ad. emarginatum, Bedd. F. B. I. t. 18.

Nilgiri and Pulney Mountains at the higher elevations. Ceylon.

(Also in Australia, New Zealand, America, Africa and the East African Islands.)

6. *ADIANTUM VENUSTUM. (Don.)* Fronds 3-4-pinnate; pinnules firm, membranaceous-chartaceous, glabrous, and slightly glaucous beneath, shortly petiolulate obovate-cuneate, rarely subrhomboid-acuminate, striated, the superior margin rounded, scarcely ever or but slightly 2 or 3 lobed, finely dentate-serrate, fertile lobes with 2, rarely 3 notches, each notch bearing a rather large sorus at the bottom; involucre reniform-cordate, submembranaceous; stipes and slender rachis everywhere ebeneous-glossy, glabrous. *Hook. Sp. Fil. ii. 40.*
A. venustum, Don Prodr. Fl. Nep. 16. Bedd. F. B. I. t. 20.

N. E. Himalayas, very common, 3,000-10,000 feet elevation.
 (Also in Cabul.)

7. *ADIANTUM PEDATUM. (L.)* Stipes 6-12 inches long, polished, dark chestnut-brown, glabrous; fronds dichotomous, with the main divisions flabellately branched; central pinnæ 6-9 inches long, 1-1½ inch broad; pinnules ½-¾ inch long, ¼ inch deep, dimidiate, broadest on the side nearest the stem, the upper and outer margin lobed, sometimes one-third down, the lowest on short slender stalks; texture pellucid-herbaceous; rachises and surfaces naked; sori roundish or transversely oblong, 1-2 lines broad. *Linn. Sp. Pl. 1557.*
Hook. Syn. Fil. p. 125. Bedd. F. B. I. t. 167.

N. W. Himalayas, from Gurwhal to Sikkim, 6,000-9,000 feet elevation.

(Also in Japan and North America.)

8. *ADIANTUM HISPIDULUM. (Swartz.)* Stipes up to 15 inches long; strong, erect, polished, dark chestnut-brown, scabrous; fronds dichotomous, with the main divisions flabellately branched; central pinnæ 6-9 inches long, ½-1 inch broad, dimidiate, sub-rhomboidal, rather broader on the side nearest the stem, hispid on both sides, the *outer edge bluntly rounded or oblique*, upper and outer margin



ADIANTUM VENUSTUM. (Don.)

finely toothed, lower ones slightly stalked; texture subcoriaceous; sori roundish, numerous, contiguous round the upper and outer edge. *Hook. Syn. Fil. p. 126. Bedd. F. S. I. t. 3.*

Common in the Western Hills of the Madras Presidency, 3,000–5,000 feet; Ceylon up to 4,000 feet.

(Also in Australia, New Zealand, Fiji, Africa and its Eastern Islands.)

9. *ADIANTUM FLABELLULATUM*. (*L.*) Scales on the rhizome long, linear, lax, chestnut coloured; fronds flabellate, bipartite-pedately divided, tripinnate; secondary pinnæ lanceolate-acuminated; pinnules glabrous, subcoriaceous-chartaceous, obliquely cuneate or semi-orbicular-cuneate, superior base truncate, superior margin 2–4 lobed and serrate-dentate in the sterile one; lobes soriferous; involucres large, the breadth of the lobe, oblong, straight, rarely a little curved, hard coriaceous; stipes elongated ebeneous-scabrous below; the rest, as well as the slender rachis, glossy and glabrous. *Hook. Sp. Fil. ii. 30. Linn. Sp. Fil. p. 1558. Hook. Syn. Fil. p. 126. Bedd. F. S. I. t. 218.*

Nepal, Assam, Khasya, Sylhet; Ceylon, in the Ouvah district; the Malay Peninsula.

(Also in the Malay Islands, South China and Japan.)

GENUS XXIV.—CHEILANTHES. (*Swartz.*)

(*Cheilos*, lip or margin; *anthos*, flower; the fructification on the margin).

Sori terminal, or nearly so, on the veins, at first small. subglobose, afterwards more or less confluent; indusium formed of the changed reflexed margin, roundish and distinct, or more or less confluent, but not quite continuous; fronds subcoriaceous in texture, mostly under 12 inches, often under 6 inches long, 3–4-pinnatifid; veins free.

1. *CHEILANTHES FRAGRANS*. (*Swartz.*) Stipes cæspitose, wiry, 1–3 inches long, densely clothed with reddish-brown linear scales; fronds 2–3 inches long, about 1 inch broad, ovate-acuminate, bi- or tripinnatifid; pinnæ opposite, $\frac{1}{4}$ – $\frac{3}{4}$ inch long, $\frac{1}{4}$ – $\frac{3}{8}$ inch broad, deltoid, cut down to the rachis below into several sinuate-pinnatifid linear-

oblong lobes; texture subcoriaceous; rachis polished but slightly scaly, both surfaces green and naked; sori small, copious; involucre light brown, membranaceous, toothed. *Hook. Syn. Fil. p. 134; Sp. Fil. ii. 81.* Polypodium fragrans (Linn.), *Bedd. F. B. I. t. 338.*

Murree, 4,000–5,000 feet; Mountains of Kashmir; Kishtwar, 3,500–5,000 feet.

(Also in Cabul, and all round the Mediterranean, Canaries, Madeira.)

2. CHEILANTHES SZOVITZII. (*Fisch and Meyer.*) Stipes densely tufted, erect, wiry, polished, brown, thinly coated, as is the rachis, with spreading woolly hairs and linear scales; fronds 3–6 inches long, 1–1½ inch broad, ovate-lanceolate, bipinnate to tripinnate; pinnæ in opposite pairs, the lower ones deltoid; pinnules linear-oblong, contiguous, cut down to the rachis below into small round bead-like segments, ⅙ inch in diameter, subcoriaceous, above green slightly tomentose, below covered with white woolly hairs, which arise from the sori, the margins much incurved; sori copious, marginal. *Fisch and Meyer in Bull. Soc. Mosc. 1838, p. 241. Hook. Syn. Fil. p. 139. Bedd. F. B. I. t. 145.*

Kashmir and Baltistone, 5,000–7,000 feet, common; Kulu.

(Also in Cabul, Asia Minor, Persia, and South Europe.)

3. CHEILANTHES MYSORENSIS. (*Wallich.*) Roots densely caespitose, the fibres very woolly, stipes slightly scaly below, short, 1–2 inches, and as well as the main rachises, deep glossy-ebeneous, rigid; fronds a span or more long, in outline narrow oblong, acute, tapering below by the diminishing of the pinnæ, glabrous, membranaceous but firm, bipinnate; lower pinnæ very small, all of them oblong-ovate, sessile, frequently opposite pinnate below, the upper half pinnatifid; pinnules or segments linear-oblong, plane (much incurved if dried without pressure), toothed or lobate-pinnatifid, each tooth or lobe bearing one or two subconfluent, small, whitish, suborbicular sori. *Hook. Sp. Fil. ii. p. 94; Syn. Fil. p. 135. Bedd. F. S. I. t. 190.*

South India, common in dry, rocky places in the plains and lower slopes of the hills; Ceylon, at low elevations.

(Also in China and Japan.)



A. J. GAY FILL. IN CYPRINAE.

CHEILANTHES MYSORENSIS. (Wall.)

4. CHEILANTHES FRAGILIS. (*Hook.*) Caudex none except the copious tufted wiry fibres, from which the stipes arise in tufts; main rachis and slightly scaly stipes (3-5 inches long) clothed with a greyish glandulose tomentum; fronds of an opaque brownish-green colour, submembranaceous, 8-15 inches long, erect, rather stiff but fragile, oblong-lanceolate, pinnate with the pinnæ pinnatifid; pinnæ alternate distant, horizontal up to 2 inches long, pinnatifid nearly to the rachis (which is furnished with subulate scales); segments crenated; lobes obtuse and partially ciliated at the margin; veins pinnated, simple or forked; sori approximate roundish; involucre ciliated, formed of the slightly changed lobes of the margin. *Hook. Fil. Exot. t. 96; Syn. Fil. p. 135. Bedd. F. B. I. t. 223.*

Perhaps only a form of No. 3.
Moulmein, on limestone rocks.



CHEILANTHES VARIANS. (*Wall.*)

5. CHEILANTHES VARIANS. (*Wall. under Pteris.*) Root tufted; stipes 4-6 inches long, slender, ebeneous-glossy, plane and margined on the upper side, obsoletely setose-paleaceous; fronds sub-membranaceous, glabrous, about a span long, long-lanceolate, the pinnatifid apex acuminate, pinnated above, bipinnate below; primary pinnæ distant, spreading or a little curved upwards, sessile, superior ones lanceolate, acuminate sinuate-pinnatifid at their base, and somewhat auricled at the upper base, lower ones deltoid acuminate pinnate at their base, pinnatifid acuminate (caudate) in the upper half; pinnules lanceolate acuminate or acute pinnatifid below, the lowest inferior pinnæ the longest. *Hook. Sp. Fil. ii. p. 89; Syn. Fil. p. 136. Bedd. F. S. I. t. 189.*

... narrow-lanceolate scales ; fronds narrow-lanceolate, 2 to 3 inches long by 3 inches broad, bipinnate or tripinnate or nearly so ; pinnae distant, sessile, deltoid, lobes distant, pinnules blunt, upper adnate entire, lower free lobed, texture very thin ; surfaces naked ; dull green beneath on the young fronds ; involucre narrow.

9. 245. *Cheilanthes Thwaitesii*, *Mett.* *Hook. Syn.* Ceylon, Kurunagalla and Kallupahane.

7. *CHEILANTHES TENUIFOLIA*. (*Sw.*) Annual, creeping, scaly ; stipes elongated, rarely scaly ; fronds glaucous, glabrous, 3-4 inches to a span and more long, bipinnate or more or less deltoid, subtripinnate, ultimate lobes distant, primary and secondary divisions the largest, more or less pinnatifid, elliptic, oblong or oblong-lanceolate subpinnatifid, margins with broad blunt teeth ; involucre mostly elongated, persistent, more or less crenated or denticulate, some wrinkled ; stipes and rachis purple-black, main rachis with secondary and tertiary rachises all with a narrow wing. *Fil.* ii. *p.* 82, *t.* 87c. *Syn. Fil.* *p.* 138. *Bedd. F. S.* Madras Presidency, common in the plains and

pinnatifid, acuminate, pinnæ mostly lanceolate pinnatifid, the one or two lowermost pair more or less half deltoid bipinnatifid below; involucre brown, scariose, rounded, sometimes confluent, and then waved or lobed, the margin entire or toothed and jagged. *Hook. Sp. Fil.* ii. p. 77; *Syn. Fil.* p. 142. *Bedd. F. S. I. t.* 191 and 192 (*bullosa*).

Throughout North India, in the hills up to 5,000 feet; Madras Presidency, in the plains and up to 8,000 feet on the hills; Ceylon; Birma.

(Also in Tropical America, Java, Philippines, East Africa and its Islands, and Arabia.)

VAR. β DALHOUSIÆ. (*Hook.*) Pinnæ (even when young) without hairs, scales or powder underneath; involucre deeply crenulate toothed or lacerate on the margin. *Clarke, F. N. I. p.* 459, and *t.* 51. *P. Dalhousiæ*, *Hook. Sp. Fil.* ii. p. 10; *Syn. Fil.* p. 137.

West Himalaya, from Kashmir to Kumaon, 6,000–9,000 feet, Sikkim, Lachen, 10,000 feet.

VAR. γ FLACCIDA. Ultimate pinnules very flaccid, and generally broader than in the type, the white powder sparse on young fronds, altogether absent on mature ones; stipes and rachis weak and often wavy. *Cheilanthes Dalhousiæ*, *Bedd. F. S. I. t.* 192.

Nilgiris, in woods near Makoorty Peak, 7,000 feet, Anamallays, open grassy places on Ponachy Hill, 6,000 feet.

VAR. δ CHRYSOPHYLLA. Powder beneath of a bright golden colour; fronds with quite the outline of typical *farinosa*, but smaller. *Hook. Fil. Exot. t.* 95, *fig.* 1. *Cheil. argentea* var. *chrysophylla*, *Hook. Syn. Fil.* p. 142.

Khasya, 5,000 feet; *Cheilanthes argentea* var. *sulphurea*, *Clarke, F. N. I. p.* 458, is only the young undeveloped frond of this.

9. CHEILANTHES SUBVILLOSA. (*Hook.*) Stipes densely tufted, 2–4 inches long, polished, naked, bright chestnut coloured, rather brittle, clothed with lanceolate acuminate scales below, main rachis glabrous beneath, the partial rachises with crisped woolly salmon-coloured hairs beneath; fronds 6–12 inches long, 2–3 inches broad,

Mr. Clarke says that the involucre is that of *L.* *farinosa*, thus this might be referred.

N. W. Himalayas, Palur Valley, and Kitghur, near

10. CHEILANTHES ALBO-MARGINATA. (*Clarke*.)
Stipes of hair-pointed scales, stipes up to 10 inches
longer than the frond; glabrous, reddish-brown, shining
particularly below and when young, with lanceolate
scales; fronds deltoid to deltoid-lanceolate, w
completely covered beneath with lanceolate brown
powdery or whitish powder, in age glabrous except
the rachis and costa, which are scaly; lowest pair of pinnae
small with their lower pinnules much more developed
than the upper (as in *farinosa*); involucre lacerate on the margins
N. I. p. 456, t. 52. Perhaps only a form of *farinosa*.
N. W. Himalayas, Kashmir, Basaoli, 5,000 feet;
Simla, 7,000 feet; Gurwhal, 2,000–9,000


11. CHEILANTHES RUFA. (*Don*.) Stipes tufted,
glabrous, densely clothed with rusty brown, woolly tomentum
on the lower surface of the scales.

being present, but very sparse ; the difference between the two is only a question of the tomentum, and both may well be varieties of *farinosa*.

Khasya, 4,000 feet, plentiful on limestone ; Sikkim, 5,000 feet ; Gurwhal, 2,000-4,000 feet ; Dalhousie.

12. *CHEILANTHES ARGENTEA*. (*Kunze*.) Stipes densely tufted, 3-6 inches long, thick, dark brown, polished, clothed at the very base with linear scales ; fronds 3-4 inches long by 2 inches broad, triangular or deltoid, tripinnatifid, lowest pinnæ much the largest but not cut down to the rachis, tripinnatifid ; rachis and costa polished like the stipe, upper surface naked, green, lower covered with white powder ; involucres crenate or fimbriate. *Kunze. Linnaea*, 1850, p. 242. *Hook. Syn. Fil.* p. 142. *Bedd. F. B. I. t.* 143. (The lowest pair of pinnæ is rarely almost quite free, the decurrent wing on the rachis from the next pair being very narrow ; the pinnæ are generally broadly decurrent, so that the frond is not cut down nearly to the rachis.)

Birma ; Khasya, 3,000-5,000 feet.
(Also in Siberia, Japan, and China.)



Nº 48.

CHEILANTHES ARGENTEA
(*Kunze*.)

GENUS XXV.—ONYCHIUM. (*Kaulf.*)

(*Onychion*, a little nail ; resemblance to the fertile segments of the frond.)

Sori placed upon a continuous linear receptacle, which connects the apices of several veins ; indusium parallel with the margin of the segments, linear, opposite, pressed down over the sori, the edge

nearly or quite reaching the midrib. Closely allied to *Pteris* and referred to that genus by Mettenius.

1. *ONYCHIUM AURATUM*. (*Kaulf.*) Stipes 6–12 inches long, stout, erect, naked, straw-coloured or pale brown; fronds 12–18 inches long, about 6 inches broad, ovate, 4-pinnatifid; lower pinnæ subdeltoid, erect-patent; pinnules and segments numerous, usually deltoid; ultimate divisions of the sterile frond often obovate-cuneate, trifid at the apex, about 1 line long, $\frac{1}{3}$ – $\frac{1}{4}$ line broad when entire, coriaceous in texture, fertile segments pod-like, $\frac{1}{3}$ –1 inch long, $\frac{1}{8}$ inch broad; rachis and both surfaces naked, the membranous involucres and copious sori a rich golden yellow. *Kaulf. En. Fil. p. 144.* *Hook. Syn. Fil. p. 143.* *Bedd. F. S. I. t. 30.*

Plains of East Bengal, from Nepal to Assam, up to 4,000 feet; Birma; (once received from the Paulghaut Hills in Southern India, but as it has not been detected since, it was probably from a cultivated specimen).


(Also in New Guinea and the Malay Islands.)

2. *ONYCHIUM JAPONICUM*. (*Kunze.*) Stipes tufted, 6–12 inches long, stout, erect, straw-coloured or pale brown, scaly at the base; fronds subcoriaceous when mature, shining on both surfaces, 12–18 inches long, ovate, 4-pinnate, lower pinnæ lanceolate-deltoid; pinnules and segments numerous, usually deltoid, the copious linear-mucronate ultimate divisions $1\frac{1}{2}$ –2 lines long, nearly uniform in the barren and fertile segments; rachis and both surfaces naked; indusium pale, membranaceous, ripe capsules deep brown. *Kunze in Sch. Fil. Suppl. p. 11.* *Hook. Syn. Fil. p. 143.* *Bedd. F. B. I. t. 21.*

Himalayas, Gurwhal to Mishmee and Khasya, 3,000–6,000 feet; Birma.

(Also in Japan and China.)

Mr. Clarke makes two varieties besides the type: 1st. *multisecta*, fertile frond, very finely cut, herbaceous, hardly shining, often 5-pinnate, ripe capsules straw-coloured, not numerous; involucre remaining closed over the ripe capsules. 2nd, *intermedia*; fronds lax, *more coarsely cut*; involucres often $\frac{1}{4}$ inch; to this 2nd variety he



A. J. P. T. E. I. C. H. E. T. H. O. M. S.

ONYCHIUM AURATUM. (Kaulf.)

refers, *Bedd. F. B. I. t.* 21, but it was taken from *multisecta*; the two varieties, however, differ very slightly, if at all.

GENUS XXVI.—CRYPTOGRAMME. (*R. Br.*)

(*Kryptos*, hidden; *gramme*, a line.)

Sterile and fertile fronds usually different from the same root; sori terminal on the veins, at first separate, subglobose, afterwards confluent, the continuous indusium formed of the changed margin of the frond, rolled over them till full maturity. (Differs from *Pellaea* rather in the dimorphic fronds than in anything else.)

I. CRYPTOGRAMME CRISPA. (*R. Br.*) Glabrous, tufted, scales at base of stipe, lanceolate, acute, pale-brown; fronds 2–4 inches long, $1\frac{1}{2}$ –2 inches broad, oblong, 3–4-pinnatifid; ultimate segments of the barren frond obovate-cuneate, deeply pinnatifid, those of the fertile frond pod-shaped, $\frac{1}{4}$ – $\frac{3}{8}$ inch long; texture thickly herbaceous, both surfaces naked. *R. Br. in Richardson's Appen. to Franklin's 1st Journal*, p. 54. *Hook. Syn. Fil.* p. 144. *C. Brunoniana* (Wall.), *Bedd. F. B. I. t.* 164. *Allosorus crispus*, *Bernh.*

Himalayas, Kumaon to Kashmir, 10,000–15,000 feet; Sikkim, 10,000–14,000 feet, head of Lachen Valley.

(Also in Arctic and Alpine Europe, Asia, and North America.)

GENUS XXVII.—PELLÆA. (*Link.*)

(*Pellos*, dark coloured, the colour of the fronds.)

Sori intramarginal, terminal on the veins, at first dot-like or decurrent on the veins, but soon running into a line; involucre or indusium formed of the more or less changed edge of the frond, quite continuous, sometimes very narrow; veins free. (Differs from *Cheilanthes* in the continuous indusium.)

§ *Veins clearly visible; indusium broad.*



A. S. CATTELL & SONS

CRYPTOGRAMME CRISTA. (R. Br.)



N° 51

PELLÆA STELLERI.
(Gmel.)

gracilis (Rup.), Bedd. *F. R. I. t.* 73.

1. PELLÆA STELLERI. (*Gmelin under Pteris*.) Rhizome wiry, creeping; stipes scattered, slender, 1-3 inches long, naked, straw-coloured or pale-brown, polished, furnished about the base with few linear pale brown scales; fronds 2-4 inches long, 1-2 inches broad, ovate bi-tripinnatifid, often with the lowest pinnæ barren, the upper fertile; pinnæ lanceolate-deltoid, 1-2 inches long, cut down to the rachis; lower pinnule again sometimes slightly divided; texture thinly herbaceous or flaccid; rachis and both surfaces naked; veins visible; young sori subterminal on the veins, clustered, not forming a marginal line till old; indusium broad, continuous, membranous. *Pellæa gracilis*, Hook. *Syn. Fil.* p. 145. Clarke, *F. N. I.* p. 460. *Allosorus*



N° 52.

PELLÆA CONCOLOR. (*Langs and Fisch.*)

Himalayas, Baltic, 9,000 feet, Kumaon, Champua, 10,000 feet, Kashmir, Pir Punju 11,000 feet; Karakorum 11,000-13,000 feet.

(Also in Canada and North United States.)

2. PELLÆA CONCOLOR. (*Langs and Fisch. under Pteris*.) Stipes 6-9 inches long, naked, erect, wiry, brownish-black, polished, slightly scaly towards the base; fronds 2-4 inches each way deltoid, cut down nearly to the rachis into 3-4 pinnæ on each side, of which the lowest pair is much the largest, with the

pinnules on the lower side much larger than the others, and deeply lobed with linear-oblong segments; rachis and costa dark coloured and polished like the stipe; texture herbaceous, lateral veins once forked; sori in broad marginal lines. *Langs and Fisch. Ic. t. 21.* *Pteris geraniifolia, Bedd. F. S. I. t. 37.* *Pellaea geraniifolia, Hook. Syn. Fil. p. 146.*

Western forests of the Madras Presidency, and in Ceylon up to 4,000 feet, very common.

(Also in Tropical America, Polynesian Islands, North China, Cape Colony, and the Mascareen Islands.)

3. *PELLÆA TAMBURII. (Hook.)* Stipes 6–9 inches long, erect, naked, chestnut-brown, polished; fronds about 6 inches each way; subcoriaceous, whitened beneath, deltoid, with three principal lobes, the terminal one cut down nearly to the rachis into several broad opposite lanceolate segments, of which the upper ones are entire and the lower ones larger and sinuated; lateral lobes with the segments on the upper side nearly entire, but those on the lower side prolonged and again deeply lobed, the largest entire divisions about 1 inch deep, $\frac{1}{2}$ inch broad at the base; rachis polished; indusium brownish, continuous, but regularly crenated along the outer edge. *Hook. Syn. Fil. p. 146; Sp. Fil. ii. t. 129A.*

Tambur Valley, East Nepal.

§§ *Veins not perceptible; indusium broad, conspicuous.*

4. *PELLÆA NITIDULA. (Wall. under Pteris.)* Caudex short-creeping, stout; stipes 2–5 inches long, very numerous and crowded, hispid, with subulate deciduous chaffy dark brown scales, and as well as the rachis (which is downy on one side) ebenous; fronds 3–4 or rarely 5 inches long, subdeltoid-oblong acuminate (sterile ones broader), coriaceous, shining beneath, glabrous, pinnate-pinnatifid, below subbipinnate; pinnae approximate, nearly opposite, broad-lanceolate dimidiate (the inferior half broadest), deeply pinnatifid nearly to the rachis, lowest pair again subpinnate and the secondary pinnae pinnatifid; lobes all oblong entire or sinuate, gradually coming to a sharp point, the lower base decurrent, the lowest inferior lobes

the longest; involucre subintramarginal, continuous or here and there interrupted, broad, flat, membranaceous, brown, close-pressed, frequently lobed and crenated and transversely wrinkled. *Wall. Cat. p. 89.* *Cheilanthes nitidula*, *Hook. Sp. Fil. ii. p. 112*; *Syn. Fil. p. 149.* *Bedd. F. B. I. t. 222.* (The involucre is sometimes continuous as in *Pteris*, sometimes interrupted and quite like *Cheilanthes*.)

Kashmir, 3,000–6,000 feet, frequent and plentiful to Chumba, scarce eastwards to Kumaon.


5. *PELLÆA BOIVINI.* (*Hook.*) Stipes 4–8 inches long, erect, naked, black, polished, very brittle; fronds 6–9 inches long, 4–6 inches broad, deltoid to lanceolate, pinnate to tripinnate; pinnæ erect-patent or spreading gradually smaller upwards; ultimate pinnales broadly ovate in the more simple forms and linear-lanceolate in the more compound, cordate or rounded at the base, very coriaceous, pale and dull beneath; both surfaces naked; veins sunk in the frond; sori in a continuous line round the segments; involucre membranous, $\frac{1}{2}$ a line broad. *Hook. Syn. Fil. p. 149.* *Bedd. F. S. I. t. 36* (under *Pteris*).

Nilgiris, Sispara and on the ghat, 4,000–5,600 feet; Anamallays, 4,000 feet, dry rocky places; Travancore and Tinnevely Mountains; Ceylon. The Nilgiri and Anamallay plant is the more simple form (pinnate or bipinnate), as represented in the figure; the Travancore plant is much larger and more compound, being tripinnate.

(Also in Madagascar and the Mauritius.)

§§§ *Veins usually hidden; indusium so narrow as to be soon hidden by the sori.*

6. *PELLÆA FALCATA.* (*Fée.*) Rhizome wide creeping; stipes 3–6 inches long, strong, erect, more or less pubescent and scaly; fronds 6–18 inches long, 1–2 inches broad, linear-oblong, simply pinnate; pinnæ 10–20 on each side, nearly sessile, $\frac{1}{4}$ –1 inch long, $\frac{1}{4}$ – $\frac{1}{2}$ inch broad, lanceolate or lanceolate-oblong, usually mucronate, slightly falcate, cuneate, or dilated and cordate at the base;



A. & C. PELLERIN


PELLAEA BOIVINI. (Hook.)

texture coriaceous; rachis densely scaly and tomentose, with surfaces nearly naked; veins not visible; sori in broad marginal lines, soon hiding the narrow involucre. *Hook. Syn. Fil. p. 151. Bedd. F. S. I. t. 22* (under *Platyloma*).

Nilgiris, Anamallays, and elsewhere on the Western mountains of the Madras Presidency, 2,000–4,000 feet elevation, Malay Peninsula; Ceylon, Telgamma.

(Also in Australia and New Zealand.)

7. *PELLÆA CALOMELANOS*. (*Link.*) Glabrous, caudex short, thick, very scaly; fronds cæspitose, subcoriaceous, oblong-triangular, bipinnate, pinnules all petiolulate, cordate- (rarely subhastate-) triangular, very obtuse, entire, sometimes subtrilobate or sinuate at the margin; the sinus deep and narrow; veins dichotomously radiated; sori continuous; involucres membranaceous; stipes short, scaly at the base, and as well as the slender rachises and petioles black-ebeneous, very glossy. *Hook. Sp. Fil. ii. p. 140. Link. Fil. Hort. Berol. p. 61. Hook. Syn. Fil. p. 152. Pteris calomelanos, Bedd. F. B. I. t. 22.*



PELLÆA FALCATA. (*Fée.*)
Base of frond.


N. W. Himalayas, below Almora, 4,000 feet, Tikri in Sirmow, 5,000–6,000 feet.

(Also in Africa, from Abyssinia to Cape of Good Hope.)

GENUS XXVIII.—PTERIS. (*Linn.*)

(From *pteryx*, a wing; the supposed likeness to wings in the branches of the common *Pteris aquilina* or Bracken.)

Sori marginal, linear, continuous, occupying a slender filiform receptacle in the axis of the indusium; indusium the same shape as the sorus, usually membranous, at first quite covering it, at length more or less spreading.



PTERIS LONGIFOLIA. (L.)

1. *PTERIS LONGIFOLIA*. (L.) Stipe 6–12 inches long, stout, erect, pale, clothed more or less below with pale brown linear scales; fronds 1–2 feet long, 4–9 inches broad, oblong-lanceolate, attenuated below; pinnæ sessile, 20–30 on each side, 3–6 inches long, $\frac{1}{8}$ – $\frac{3}{8}$ inch broad, linear, entire, truncate or cordate or slightly auricled at the base; texture subcoriaceous; rachis naked or slightly scaly below or throughout; both surfaces naked; veins close and fine, usually once-branched; indusium membranous, yellowish-brown. *Linn. Sp. Pl.* 1531. *Hook. Syn. Fil. p.* 153. *Bedd. F. S. I. t.* 33.

Madras Presidency, in many localities; throughout Bengal in the plains and up to 5,000 feet; Ceylon; Birma.

(Also widely distributed all round the world.)

2. *PTERIS CRETICA*. (L.) Stipes 6–12 inches long, erect, wiry, naked, straw-coloured, polished; fronds 6–12 inches long, 4–8 inches broad; lateral pinnæ usually in 2–6 opposite sessile pairs, of which the upper one is sometimes a little decurrent, 3–6 inches long $\frac{1}{8}$ – $\frac{3}{4}$ inch broad, the sterile ones generally the broadest, spinulose-serrated, the lowest or even 1–4 other pairs often cleft down nearly to the base into 2–3 linear pinnules; margins, where barren, spinulose-serrate; rachis and both surfaces naked; texture coriaceous; veins fine, close, parallel or wide apart in the narrow-segmented forms, simple or once-forked; involucre membranous. *Linn. Mant. p.* 130. *Hook. Syn. Fil. p.* 154. *Bedd. F. S. I. t.* 39.

Madras Presidency, very common from sea-level up to 8,000 feet; Bengal Presidency, very general up to 9,000 feet; Ceylon; Birma.

(Also widely distributed in Africa, Tropical America, South Europe, and elsewhere.)

VAR. β *HETEROMORPHA*. Pinnæ abnormally pinnatifid. *Pteris heteromorpha*, *Hook. Syn. Fil. p.* 156. *Bedd. F. B. I. t.* 89. This is, I think, only a variety of *cretica*, not a distinct species.

Birma.

(Also in the Philippine Islands.)

3. *PTERIS PELLUCIDA*. (*Presl.*) Stipes 1 foot or more long,

erect, naked ; fronds 12-18 inches long, ovate-lanceolate, pinnate ; pinnæ acute, sometimes many, sometimes 7, 5, 3, or 1, all simple, or the lowest pair 2-fid, often 1 inch broad or rather more ; margins where barren, entire, undulate, crisped or crenulate ; rachis naked ; texture coriaceous, surfaces bright green often glossy ; veins fine close, simple or forked at right angles from the midrib ; indusium membranous. *Presl. Rel. Hænk.* 55. *Hook. Syn. Fil.* p. 154. *Clarke, F. N. I.* p. 462. *Bedd. F. S. I. t.* 38.

Plains and hills of Bengal up to 3,000 feet elevation, Western forests of Madras Presidency ; Birma.

(Also in Malay Islands, and Guinea Coast.)

VAR. β STENOPHYLLA. Pinnæ 3-4 or sometimes 2-1 subdigitate, fertile pinnæ very long and narrow. *Hook. and Grev. Ic. Fil. t.* 130. Gurwhal and Nepal, 3,000-4,000 feet.

4. PTERIS HOOKERIANA. (*Agardh.*) Stipes 6-12 inches long, naked, erect, pale ; fronds coriaceous, opaque, pinnate or bipinnate, 9-12 inches long, 6-9 inches broad, with 2-6 pairs of sessile or stalked alternate or opposite entire pinnæ, the largest of which is about 6 inches long, $\frac{3}{4}$ -1 $\frac{1}{4}$ inch broad, the 2-4 lower pairs more or less pinnatifid ; rachis and both surfaces naked ; veins about 1 line apart at the base, simple or once-forked, conspicuous above, but scarcely visible beneath ; sori slightly intermarginal ; indusium narrow. *Hook. Syn. Fil.* p. 155. *Bedd. F. S. I. t.* 40.

Ceylon, Adam's Peak, about 2,000 feet elevation, and the Southern provinces at low elevations.

5. PTERIS DACTYLINA. (*Hook.*) Stipes 6-9 inches long, slender, wiry, naked ; fronds digitate, coriaceous, consisting of 3-7 (usually 5) linear glabrous pinnæ, 2-6 inches long by 1 $\frac{1}{2}$ -2 lines broad, the barren margins serrate ; veins wide apart, thick and conspicuous, simple or forked ; indusium broad. *Hook. Syn. Fil.* p. 155. *Bedd. F. B. I. t.* 23.

Khasya, 4,200 feet elevation ; Sikkim.

6. PTERIS ENSIFORMIS. (*Burm.*) Stipes up to 10 inches long,

naked; fronds subcoriaceous, more or less dimorphous, up to 12 inches long by 3-6 inches broad, fertile ones with a long terminal pinna, and 2-4 pairs of lateral ones which are simple, forked, trifid or even more compound, the lateral lobes being pinnatifid, the segments very narrow, $\frac{1}{4}$ inch or less broad, entire; sterile fronds generally smaller, more compound, the lateral pinnæ being pinnate, sharply toothed, and much broader than in the fertile; veins forked or simple, diverging at about two-thirds of a right angle from the midrib, very close together. *Burm. Fl. Ind. p. 230. Hook. Syn.*

Fil. p. 155. Pteris crenata (Sw.), Bidd. F. S. I. t. 35.

Malabar, rare; Hills of Vizagapatam district, common; plains of East Bengal, Bhotan to Chittagong; Ceylon; Birma; and Malay Peninsula.

(Also in Tropical Australia, China, and Polynesia.)



PTERIS DACTYLINA. (Hook.)

lower pinnæ; it has more the habit of crenata, I think, than of ensiformis.

Sylhet, Mishmee, Khasya, Chittagong Hills.

7. PTERIS GRIFFITHII. (Hook.) A foot and more high, slender graceful; caudex none; roots of tufted fibres; frond ovate-lanceolate, submembranaceous pinnate, upper pinnæ simple, 2 or 3 of the lower pairs again pinnated simple; pinnæ and pinnules exactly linear, $\frac{1}{2}$ -1 $\frac{1}{2}$ inch long, obtuse, mostly opposite, everywhere

entire sessile, with the base adnate and decurrent, so as to form a narrow wing to the rachis; lowest pinnules of the inferior pinnæ sometimes again divided, terminal pinnule always elongated; veins distant, once-forked, veinlets divaricating; sori continuous, involucre sub-intramarginal, membranaceous, not reaching to the apex; stipes slender, longer than the fronds, stramineous-brown and a little scaly at the base. *Hook. Sp. Fil.* ii. p. 170, t. 123A. *Bedd. F. B. I. t.* 24. Mishmee.

8. *PTERIS SEMIPINNATA*. (L.)

Stipes strong, erect, dark brown, polished, hispid at the base, and then furnished with linear scales; fronds 12–18 inches long, 6–9 inches broad, ovate-lanceolate, the upper part cut down nearly to the rachis into numerous close entire linear lobes, the largest of which are $1\frac{1}{4}$ –3 inches long, $\frac{1}{4}$ – $\frac{1}{2}$ inch broad, the lower two-thirds with 6–8 pairs of opposite distantly placed pinnæ, with long linear entire points and a broad

entire wing on the upper side of the rachis, the lower side with several linear pinnules 1–2 inches long, $\frac{1}{4}$ inch broad; texture scarcely coriaceous; barren segment finely serrated; rachis and surfaces naked; veins simple or once-forked, $\frac{1}{4}$ inch apart at the base; involucre membranous. *Hook Syn. Fil.* p. 157. *Linn. Sp. Pl.* 1534. *Bedd. F. S. I. t.* 34.



NO 57

PTERIS GRIFFITHII. (Hook.)

East Bengal, Assam to Chittagong, up to 1,000 feet ; Travancore Hills ; Birma.

(Also in Malay Islands, China, and Japan.)

In Indian examples the pinnæ are always semipinnate, but in Chinese and Japanese forms the upper margin of the pinnæ is also pinnatifid.



Nº 58.

PTERIS SEMIPINNATA. (L.)

9. *PTERIS DALHOUSIÆ.* (Hook.) Stipes strong, erect, about 1 foot high, polished, naked ; fronds 2-3 feet long, 12-18 inches broad, 3-4 pinnatifid ; upper pinnæ linear, unbranched decurrent down to the next pair, lower ones sometimes 1 foot long, deltoid ; pinnules with simple or branched linear segments, the longest undivided ones 6 inches long, $\frac{1}{4}$ - $\frac{3}{8}$ inch broad, those of the pinnæ with usually about 1 inch between them ; the margins very slightly serrated ; texture subcoriaceous ; rachis and surfaces naked ; veins fine, simple, or once-forked ; indusium narrow-membranous. *Hook. Syn. Fil. p. 157. Bedd. F. B. I. t. 191.*

Penang.

10. *PTERIS QUADRIAURITA.* (Retz.) Stipe glabrous or scabrid ; fronds very variable in size and texture, lanceolate or ovate-lanceolate, with 3-11 subopposite pairs of pinnæ ; in the typical form all except the lowest pair are narrowly oblong acuminate or caudate, and cut down nearly to the rachis into many uniform segments, which are obtuse or acute entire or serrated, and the lowest pair bipartite ; but in some forms several or nearly all the pinnæ are bipartite, and the lowest or two lower pairs have several pinnæ descending from the lower side, and in one form the lowest pair is completely bipinnate with five pinnæ on each side of the rachis ; veins conspicuous,

usually once-forked ; lowest veins reaching the margin above the sinus when very deeply cut, at the sinus when less deeply cut ; sori generally partial on the segments, rarely continued to the apex. *Retz. Obs.* vi. 38. *Hook. Syn. Fil.* p. 158. *Bedd. F. S. I. t.* 31.

Throughout India, Ceylon, and the Malay Peninsula, from the plains up to 8,000 feet, very common.

(Also all round the world throughout the tropics and a little beyond them.)

The following varieties are tolerably constant even in cultivation :

VAR. β ARGENTEA. Like the type, but with broad white bands down the centre of the pinnæ.

Nilgiris and Western mountains of the Madras Presidency only at high altitudes ; much in cultivation even in England.

VAR. γ ASPERICAULIS. Rachis and nerves red ; stipe and rachis asperous, generally very rigid, coriaceous, and shining. (Often called rubro-nerva.)

Wynaad, Coorg, and elsewhere on the western side of Madras Presidency, 3,000–4,000 feet elevation.

VAR. δ SETIGERA. Rachises and costules setigerous, very membranaceous in texture, nearly all the pinnæ bipartite, the lowest pair with sometimes several pinnæ descending along the lower side. *Bedd. F. B. I. t.* 202.

Coorg and Malabar : Moulmein on limestone rocks.

VAR. ϵ LUDENS. Pinnules very abnormally cut, sometimes only with a waved margin for the greater length, then developing normal segments, then again narrowing into only the waved margin. *Bedd. F. S. I. tabs.* 41 and 219 (under Otaria.)

Malabar, Quilon ; Ceylon.

Mr. Clarke gives the following varieties for Northern India :

VAR. ζ KHASIANA. The lowest pinnæ with 5 secondary pinnæ

on each side of the rachis (*i. e.* completely bipinnate), those of the upper margin little smaller than those of the lower.

Khasya, 3,000 feet elevation.

Mr. Clarke says it graduates into the type.

VAR. η BLUMEANA. Lateral pinnæ with 50–60 segments, which are 1 inch long by $\frac{1}{8}$ inch broad, equally wide throughout or broader at apex, not falcate, obtuse, rounded, entire or crenulate at the barren apex. A very large fern. *Clarke, F. N. I. t. 55.*

Chittagong ; Tenasserim ; Singapore.

Much the habit of longipinnula.

VAR. θ SUBINDIVISA. (*Sp. Clarke.*) Small, with only 3 pinnæ, terminal one 6 inches long, with many segments, lateral pair hardly longer than the segments of the terminal pinna, or subobsolete.

Bhotan, Sikkim bed of the Teesta, 1,000 feet elevation.

Mr. Clarke has constituted this a species in his work on the "Ferns of Northern India," but he now agrees that it is only a variety of quadriaurita. *Clarke, F. N. I. t. 56.*

(Mr. Clarke's variety major has arched veins and is *Campteria biaurita*.)

11. PTERIS GREVILLEANA. (*Wall.*) Dimorphic, barren stipe shorter, winged towards the apex ; barren frond pedately 5-fid, scarcely pinnate, margin spinulose-serrate ; fertile frond with 5 pinnæ, the lower pair bipartite, veins exceedingly obscure. *Wall. Cat. 2,680. Agardh, Recens. Gen. Pter. 23. Clarke, t. 54.*

Sylhet, Cachar, Shapoor.

Suspiciously like quadriaurita, except that the fronds are dimorphic. Mr. Clarke thinks it is more nearly allied to ensiformis.

12. PTERIS LONGIPINNULA. (*Wall.*) Stipes erect, naked, 2–3 feet ; fronds up to 3 feet long, and 18 inches broad, pinnate ; pinnæ about 5 pairs, large, rather distant, deeply pinnatifid, segments entire, blunt, falcate, not distant, $1\frac{1}{2}$ – $2\frac{1}{4}$ inches long, $\frac{1}{4}$ – $\frac{1}{2}$ inch broad, apex usually barren, crenate, or subentire, the lowest pair undivided, or rarely bi-partite ; texture subcoriaceous ; rachis and surfaces naked ;




Fig. 1000

PTERIS PATENS. (Hook.)

veins once forked; sori continuous nearly to the apex of the segments. *Hook. Syn. Fil. p.* 158. *Bedd. F. S. I. t.* 43.

Sikkim; Khasya; Cachar; Sylhet up to 3,000 feet elevation; Travancore and Tinnevely Mountains; Malay Peninsula.

(Also in the Malay Islands.)

13. *PTERIS EXCELSA*. (*Gaud.*) Fronds ample, 5–6 feet long, submembranaceous, light green, pinnate; pinnæ large, remote, 6 inches to a foot long, numerous, ovate-lanceolate caudate sessile, lower ones long-petiolate, lowest pair of pinnæ entire, or sometimes bipartite, all deeply pinnatifid nearly to the costa; segments (2–4 inches long) from a broad base, linear-lanceolate obtuse serrated subfalcate, lower base decurrent, veins free, forked at or near the middle or 3–4 branched; very rarely a few veinlets anastomose; involucre continuous from the base almost to the apex, rather broad; stipes very stout and flexuose; rachises bright castaneous glossy. *Hook. Sp. Fil. ii.* 183. *Gaud. in Fréyc. Voy. Bot. p.* 388. *Bedd. F. B. I. t.* 218.

Kumaon, Gurwhal; Dalhousie, 4,000–8,000 feet elevation; Sikkim, 8,000–9,000 feet; Nynee Tal.

(Also in the Sandwich Islands.)

14. *PTERIS PATENS*. (*Hook.*) Caudex erect; stipes numerous, stout, scaly at their base, rachis stout, glabrous; fronds ample, erect, strict, membranaceous, pinnate; lower pinnæ bi-tripartite; pinnæ 8–16 inches long, strict horizontally patent, numerous approximate, elongate oblong, acuminate, rarely subcaudate, sessile or lower ones shortly petiolate, deeply (almost to the rachis) pinnatifid; segments horizontal, very uniform, linear, scarcely acuminate, serrulate where sterile, the base dilated decurrently-confluent on both sides, especially at the lower base, sometimes apart; veins all free, simple or once or twice forked; sori continuous nearly to the apex. *Hook. Syn. Fil. ii.* 177. *Bedd. F. S. I. t.* 205. *Campteria patens*, *Baker. Hook. Syn. Fil. p.* 165. (The supposed narrow costal arches are only a wing to the rachis.)

Madras Presidency, Carcoor ghat, Wynaad and Travancore Hills,

2,000–2,500 feet elevation; Ceylon, Dodawilla, Matele and Moorowa forests, above 2,000 feet elevation.

(Also in Borneo, Philippines and Society Islands.)

15. *PTERIS LONGIPES*. (*Don.*) Fronds membranaceous, furnished above with a few subulate scales on the rachis of the pinnæ, and with hairs on the rachis of the segments, ternately divided, or subpedately divided into 5–7 segments, lateral divisions multipinnate; pinnæ short, petiolate or sessile lanceolate acuminate, deeply pinnatifid to the apex, or with the apex long caudate, segments oblong obtuse subtruncate, serrated at the sterile portions of the apex; sori lateral on the segments, never extending to the apex or base; veins rather remote, all forked, basal ones arising from the axil of the costule, and extending to the margin much above the sinus. *G. Don. Prod. Fl. Nep.* 15. *Pteris pellucens*, *Agardh, Sp. Gen. Pter.* 43. *Hook. Sp. Fil.* ii. 191. *Bedd. F. S. I. t.* 32. (South Indian examples are always ternately divided, but those from N. India have often 5, rarely 7, pinnæ.)

All the western hills of the Madras Presidency, 2,000–5,000 feet elevation; Ceylon; Birma; Sikkim, Bhotan; and Khasya 1,000–5,000 feet elevation.

(Also Guinea coast and Guatemala.)

§ *Stipes distant from a long creeping rhizome, indusium double.*
(*Genus Pæsia and Ornithopteris of authors.*)

16. *PTERIS AQUILINA*. (*L.*) Rhizome stout, creeping underground; stipes about 1 foot long, strong, erect, naked; fronds 2–6 feet long, 1–2 feet broad, subdeltoid in outline, only the uppermost pinnæ simple, the next lanceolate cut down nearly or quite to the rachis into short triangular or linear pinnules, the lowest long-stalked, 1 foot or more long, with ample lanceolate pinnules which are cut down to the rachis into numerous lanceolate segments, which are again fully pinnate; largest entire ultimate segments 1 inch long, 2 lines broad; texture thin or subcoriaceous; rachis and surfaces naked or pubescent; veins close, conspicuous, often twice forked; involucre double, or the

inner obsolete. *Hook. Syn. Fil. p.* 163. *Linn. Sp. pl.* 1533. *Bedd. F. S. I. t.* 42.

Throughout India, Ceylon, and the Malay Peninsula on the mountains, 2,000–8,000 feet elevation.

(Also throughout the whole world, except the Arctic zones and temperate South America.)

VAR. β ESCULENTA. Ultimate segments distant, linear and elongated. Moulmein and the Malay Peninsula southwards.

GENUS XXIX.—CAMPTERIA. (*Presl.*)

(*Kamptos*, arched; the lowest veins meeting and forming an arch.)


As in *Pteris*, but the lowest veinlets from the base of the vein or costule of each segment united with the veinlet from the next costule, and forming long arches along the costa of the pinnæ; upper veinlets free.

1. CAMPTERIA BIAURITA. (*Linn. under Pteris.*) Quite similar to *Pteris quadriaurita*, except that the costules of the segments are connected by a long arching veinlet; the pinnæ are generally less deeply divided, and the plant is generally larger and coarser, but these two latter differences cannot be relied on. *Linn. Sp. Pl.* 1534, under *Pteris*. *Hook. Syn. Fil. p.* 164. *Bedd. F. S. I. t.* 44.

Western parts of Madras Presidency up to 6,000 feet; not common. Pretty general throughout Northern India from the plains up to 6,000 feet. Ceylon; Malay Peninsula.

(Also in Malay Islands, China, tropical Africa, and Australia.)

2. CAMPTERIA KLEINIANA. (*Presl.*) 4–6 feet high; fronds very large, sub-membranaceous, pedately ovate, pinnate; pinnæ sub-sessile lanceolate with a long acumination; lowest pair with two large pinnæ descending from the lower margin of the costa; all the pinnæ deeply pinnatifid (leaving a broad margin on each side of the costa); segments falcate oblong, sterile ones broadly crenated, and fertile ones



L. CAMPILLO & P. B. R. C.

CAMPTERIA KLEINIANA. (Presl.)


crenated at their sterile apices; basal veins uniting in pairs into an arch below the sinus (one generally proceeding from the costa, the other from the costule), and bearing 3-4 free veinlets; other veinlets forked, terminating in dots within the margin; indusium never reaching the base or apex of the segments. *Klein. Tent. Pter.* 147. *t.* 5. *Pteris geminata*, *Wall. Cat.* 2180. *Hook. Syn. Fil.* 164. *C. Anamallayensis*, *Bedd. F. S. I. t.* 45.

Western mountains of Madras Presidency 2,000-4,500 feet; rather rare. United by Hooker and by Clarke with *biaurita* but really quite distinct; its thin texture and very wavy appearance are very unlike any form of *biaurita*; its margin is, besides, always crenated, and the veins never reach the margin, which they always do in *biaurita*. It is one of the most beautiful ferns in the Madras forests, and has been for years in cultivation at Ootacamund. Wight's specimens from Courtallum are named *geminata* in the Kew Herbarium, and there are no specimens from N. India; there are specimens from Johanna Island; also from the Isle of Mayotta, (collected by Lady Barkly), named by Mr. Baker *Campteria maxima*. *Sp. Fil. p.* 480.

3. CAMPTERIA WALLICHIANA. (*Ag.*) Fronds ample, glabrous, sub-membranaceous, 3-partite, lateral branches compoundly divided, terminal one elongated, pinnated; pinnæ numerous, sessile, 4-6 inches long, linear-lanceolate acuminate, deeply pinnatifid; segments linear-lanceolate obtuse, sterile ones serrulate, all approximate, about $\frac{1}{4}$ an inch long; basal veins monarculate, the rest free and forked, rarely a solitary areole at the base of the costule; stipes very long, 5-6 feet; stout, rich castaneous, and, as well as the paler-coloured rachises, very smooth and glossy. *Ag. Sp. Gen. Pter. p.* 69. *Hook. Sp. Fil. ii.* 206. *Syn. Fil. p.* 165. *Bedd. F. B. I. t.* 25 and 217. *Var. quadripinnata* (*Clarke*) is a very compound form with the pinnæ tripinnate; it is probably an abnormal form or monstrosity which would not be constant.

Himalayas from Chumba to Bhotan 3,000-8,000 feet, abundant; Khasya.

(Also in the Philippines, Java, and Samoa.)



A. D. CASTELL. DORYPTERIS.

DORYOPTERIS LUDENS (Vall.)

FERNS OF BRITISH INDIA AND CEYLON.

GENUS XXX.—DORYOPTERIS. (*J. Smith.*)

(*Dory*, spear ; *pteris*, form of the fronds.)

Fronds small, sub-pedate or sagittate, in texture and colour like *Pellaea* ; veins copiously anastomosing, without free included veinlets ; the rest as in *Pteris*.

1. DORYOPTERIS LUDENS. (*Wall.*) Rhizome creeping, furnished with linear adpressed brown scales which have white margins ; stipes solitary distant, polished, sometimes with a few scales, and often with dusky sub-tomentose pubescence at base and apex ; barren frond on a stipe 3-4 inches long, triangular with two slightly deflexed basal lobes, to hastate with two basal and two large spreading lateral lobes, the margin entire ; fertile frond, on a stipe often 12 inches long, 4-6 inches each way, cut down into five linear-lanceolate or lanceolate lobes, one erect, two spreading and two deflexed, of which all except the last are sometimes again forked ; texture coriaceous, costa polished, veins hidden ; sori continuous all round the margin. *Wall. Cat.* 88. *Hook. Syn. Fil.* 166. *Clarke, F. N. I.* 470. *Litobrochia ludens* and *pedata*, *Bedd. F. N. I. t.* 26 and 27.

Chittagong Hills up to 1,000 feet elevation ; Orissa, on the Balasore Hills ; Birma. (A specimen in Wight's herbarium of this or an allied species is supposed to be from the Dindigul mountains in the Madras Presidency ; but it has never been found there of late.)

(Also in the Philippine islands.)

GENUS XXXI.—LITOBROCHIA. (*Presl.*)

(*Lithos*, a stone ; *brocha*, spots ; the areoles of the net-like veins resembling pavement.)

Veins copiously anastomosing with some free included veinlets ; otherwise as in *Pteris*.

1. LITOBROCHIA INCISA. (*Thunb.*) Caudex long creeping, subterraneous ; stipes and rachis castaneous glossy ; fronds ample, sub-



LITOBROCHIA INCISA. (Thunb.)

membranaceous, distant ovate, long stipitate, glaucous beneath, tripinnate; pinnae all sessile, subadnate mostly opposite horizontal; pinnules opposite sessile, lanceolate obtuse, more or less deeply lobed and pinnatifid; segments ovate or oblong (sterile ones subsinuate) or triangular, lowest one often remote and forming auricles on the rachis, superior ones confluent; veins all anastomosing, areoles next the costa and costules the largest and most elongated; involucres continuous or interrupted, membranaceous entire at the edge. *Thunb. Pr. Fl. Cap.* 171. *Hook. Syn. Fil. p.* 172. *Litobrochia aurita*, *Hook. Sp. Fil.* ii. 231. *Bedd. F. S. I. t.* 221.

Ceylon, central and southern provinces, above 3,000 feet. Travancore Hills 3,000–4,000 feet. Sispara ghat on the Nilgiris, 4,500 feet. Sikkim, Bhotan, and Khasya, 3,000–6,500 feet.

(Also widely distributed in the tropics and sub-tropics of America, Africa, Asia, Australia, and Polynesia.)

2. LITOBROCHIA MARGINATA. (*Bory, under Pteris.*) Stipes elongated, stout, and with the rachis sub-castaneous; fronds ample, tripartite, lateral branches pinnate, spreading, long petiolate, sub-membranaceous, glabrous; pinnules 4–6 inches or more long, linear-oblong or linear-lanceolate sub-sessile, acuminate, rather deeply pinnatifid; segments approximate, linear-oblong falcate (upwards) obtuse or acute, entire or scarcely serrated and only at the apex, nearly half an inch long; sinuses obtuse; basal veins forming a single arc parallel with the costa; veins forming a series of two or more areoles (parallel with the costule), from which proceed free veinlets which do not extend to the margin but terminate with a thickened apex just within it; involucres continuous on the lower two-thirds of the margin of the pinnules not extending to the apex. *Bory. Voy.* 2. p. 192. *Hook. Syn. Fil. p.* 172. *Pteris tripartita*, *Hook. Sp. Fil.* ii. 225. *Litobrochia tripartita*, *Bedd. F. S. I. t.* 220.

Ceylon, common about Calle and the central provinces; Madras Presidency, much cultivated, and said to be found wild in the Cochin forests; Malacca.

(Also in West tropical Africa, the East African Islands, Queensland, the Philippines, and Polynesia.)

GENUS XXXII.—CERATOPTERIS. (*Brong.*)

(*Keras*, *keratos*, a horn ; *pteris*, a fern: the horned fern.)

Sori placed on two or three distantly anastomosing veins which run down the frond longitudinally, and are parallel both with the edge and midrib ; veins of the sterile fronds articulated in oblique oblong hexagonal areoles. Capsules scattered on the receptacles, sessile, subglobose, with a ring which is either complete or more or less partial or obsolete. Indusium formed of the reflexed margin of the fronds, those of the two sides meeting against the midrib. A very anomalous genus, regarded by some as a distinct order ; it is very unlike Pterideæ, and should be placed in a distinct tribe.

1. CERATOPTERIS THALICTROIDES. (*Lin.*) Stipes tufted, thick inflated, filled with large air-cells ; fronds succulent in texture, the barren ones floating or erect, simple or slightly divided when young, bi-tripinnate with narrow linear segments when mature, fertile ones bi-tripinnate ; ultimate segments podlike. *Hook. Syn. Fil. p.* 174. *Bedd. F. S. I. t.* 75. *Acrostichum thalictroides*, *L. Sp. Pl.* 1527.

Throughout India, Ceylon, and the Malay Peninsula up to 3,000 feet elevation ; common in tanks, ditches, and swampy places, or even dry ground during the rains. Mr. J. Smith says it is an annual, but I do not think it is so in cultivation, if kept in water or very moist, as I had the same plant growing for some years at Ootacamund.

(Also in the tropics of the whole world.)

GENUS XXXIII.—LOMARIA. (*Willd.*)


(*Loma*, fringe or border ; relating to the indusium.)

Sori linear continuous, parallel with the midrib and occupying the whole, or nearly the whole, of the space between it and the edge ; indusium membranous, formed of the revolute edge of the frond. Fronds dimorphous, usually once or twice pinnatifid or pinnate, rarely simple or bi-pinnate ; veins free ; ring of capsule vertical.

1. LOMARIA PATERSONI. (*Spreng.*) Rhizome short-creeping ; stipes 2–3 inches long, wiry, erect, rather scaly below ; sterile frond simple, about 1 foot long and 1 inch broad, narrowed at both ends, or pinnatifid, 2 feet or more long, cut down nearly to the rachis into 6–12 segments on each side, which are often 6–9 inches long and nearly 1 inch broad, and suddenly decurrent at the base ; texture coriaceous ; fertile fronds simple and only $\frac{1}{8}$ inch broad, or pinnatifid with numerous segments on each side, 6 inches long by $\frac{1}{8}$ inch broad ; or the fronds are sometimes in part sterile, in part fertile ; veins prominent in the young sterile fronds, inconspicuous in mature fronds, forked, thickened at the apex close to the margin ; sori covering the whole space between midrib and margin. *Spr. Sys. Veg.* iv. 62. *Hook. Syn. Fil.* 174. *L. elongata* (Blume), *Bedd. F. S. I. t.* 28, 28A.

Nilgiris and Anamallays 5,000–8,000 feet elevation. Ceylon, 4,000 feet and upwards. Nilgiris examples have the fertile fronds always pinnatifid as far as I have observed, and the sterile ones generally so. From Ceylon I have examples with the sterile and fertile both quite simple.

(Also in the Philippines, Fiji, New Zealand, S. Australia, and Tasmania.)



A. S. CHITALL & CO. INC. - BOSTON

LOMARIA PATERSONI. (Spr.)

GENUS XXXIV.—PLAGIOGYRIA. (*Kunze.*)

(*Plagios*, oblique ; *gyros*, a circle ; the oblique ring of the capsule.)


As in *Lomaria*, except that the capsules have an oblique ring, and the base of the stipe is suddenly dilated, fleshy, triquetrous, and furnished with large spongy glands.

1. *PLAGIOGYRIA ADNATA*. (*Blume.*) Caudex stout short; stipes caespitose, elongated, brown, triquetrous; a span to 1½ foot long; slender, naked at the base, dilated and subcarnose, furnished with orbicular spongy glands; fronds a foot and more long, ovate-lanceolate, chartaceous-membranaceous, sterile ones pinnatifid almost to the rachis, below pinnated, but all the pinnæ united by a narrow wing; segments mostly opposite distant lanceolate, much acuminate, more or less falcate, the inferior base round, superior base extended upwards, the margins entire or obscurely toothed, the apex strongly serrated; fertile fronds pinnated; pinnæ alternate distant linear sessile, rather obtuse; sori covering the whole under side between the costa and the brown membranaceous very distinct involucre. *Bl. En. Fil. Jav. p. 205.* *Hook. Sp. Fil. iii. 19.* *Bedd. F. B. I. t. 51.*

Khasya, 4,000–5,000 feet elevation.

(Also in Japan and Java) Mr. Clarke thinks it possible that this is not the same as Blume's Java plant, as the only example of that is a portion of a fertile frond.

2. *PLAGIOGYRIA GLAUCA*. (*Bl.*) Caudex stout erect; stipes elongated, subcompressed tetragonous with two furrows in front, the base dilated triquetrous on the anterior face bearing spongy depressed glands, fronds ovate-oblong 1–2 feet long, pinnated; sterile pinnæ numerous, 3–5 inches long, chartaceous, horizontally spreading, sessile, or very nearly so, lanceolate, acuminate at the base, truncated beneath, having a gland as much attached to the rachis as to the very short petiole; the margins finely dentate-serrulate, rather



more strongly at the apex, white and almost powdery beneath, green above with a tinge of red; veins copious, compact, simple and forked; ultimate short pinnæ confluent into a pinnatifid and lobed caudate apex, fertile pinnæ narrow-linear obtuse erect-patent on very short petioles, which sometimes bear a gland beneath; sori and involucres as in other Plagiogyriæ. *Bl. En. Fil. I. p. 204.* Lomaria (Plagiogyria) glauca, *Hook. Sp. Fil. iii. 22.* *Bedd. F. B. I. t. 90.*

Khasya, 4,000–5,000 feet elevation. (*Probably only a variety of pycnophylla.*)

(Also in Java.)

3. *PLAGIOGYRIA PYCNOPHYLLA.* (*Kunze.*) Caudex stout, erect, very woody, and having many wiry roots; fronds fascicled; stipes sub-quadrangular 2-furrowed in front (as is the rachis), at the base dilated fleshy, triquetrous, bearing spongy glands; fronds 1–2 and more feet long, sterile ones, sub-chartaceous opaque, brown when dry, oblong-ovate pinnated for its whole length, except that the small terminal ones are confluent into a lobed (or pinnatifid) and finely acuminate and serrated apex, pinnæ horizontally patent sessile or very nearly so, numerous, approximate, narrow oblong-lanceolate very finely almost caudately acuminate, the base very obtuse and truncate (neither dilated nor contracted), at the base beneath furnished with one or two prominent glands (sometimes wanting); the margin very minutely serrulate, the caudate apex strongly serrated; veins numerous, closely placed, simple and forked; fertile fronds narrower pinnated; pinnæ linear obtuse on very short petioles with a gland on the under side; involucre at first vaulted, conspicuous, membranaceous, brown, at length forced back by the capsules. *Kze. in Bot. Zeit. iv. p. 143.* *Hook. Sp. Fil. iii. 21.* *Bedd. F. B. I. t. 52.*

Nepal and Bhotan, 6,000–11,000 feet; very common at 7,000–8,000 feet. Khasya, 4,000–6,000 feet; Malay Peninsula.

(Also in Java.)

4. *PLAGIOGYRIA EUPHLEBIA.* (*Kunze.*) Caudex stout woody ligated; stipes 1 foot or more long, naked erect, triquetrous below,

fronds ovate-lanceolate, 1-2 feet long, 6-12 inches broad; pinnæ 1 inch or more apart at the base, all stalked except towards the apex, erect-patent, linear, 4-6 inches long, $\frac{1}{4}$ - $\frac{3}{4}$ inch broad, narrowed at the base on both sides, and at the apex, where it is toothed; texture subcoriaceous; veins prominent; fertile fronds similar, but the pinnæ more distant and narrowly linear; rachis erect naked. *Kze. in Bot. Zeit.* 1848, p. 521. *Hook. Syn. Fil.* p. 183. *Bedd. F. B. I. t.* 165.

Nepal, Khasya, 4,000-6,000 feet.

(Also in Japan and N. Australia.)

TRIBE VII.—BLECHNEÆ.

Sori linear or oblong dorsal, parallel with the midrib and edge of the segments, not close to the latter; indusium the same shape as the sorus, superior, opening towards the midrib.


GENUS XXXV.—BLECHNUM. (L.)

(One of the Greek names for a fern.)

Sori linear, continuous, or nearly so, parallel with and usually contiguous to the midrib; indusium membranous, distinct from the edge of the frond; fronds uniform or only slightly dimorphous, generally pinnate or pinnatifid; veins free.

1. BLECHNUM CARTILAGINEUM. (*Swartz.*) Caudex oblique, densely clothed at the crown with blackish fibrillose scales; stipes strong erect scaly and muricated in the lower part; fronds ovate-oblong, 1-2 feet long, 6-9 inches broad, pinnatifid and pinnate towards the base; pinnæ numerous erect-patent linear, 4-6 inches long, $\frac{1}{4}$ - $\frac{1}{2}$ inch broad, narrowed gradually upwards, margin serrulate; the lowest pinnæ linear distant, narrower than those above, but not reduced to mere auricles. *Sw. Syn. Fil.* 114 and 312. *Clarke, F. N. I.* p. 473. *Hook. Syn. Fil.* 184.

Mishmee.—Mr. Clarke refers Griffith's solitary specimen to this species, which is Australian. Hooker referred it to nitidum, an



BLECHNUM ORIENTALE. (L.)

American plant. The two species appear to me so very closely allied as to be distinguished only with difficulty, if at all; the rachis is slightly puberulous in both. The figure *t.* 49, *Ferns Brit. Ind.*, given to represent the Mishmee plant, is a copy of Hooker's figure of the American species.

2. BLECHNUM SERRULATUM. (*Rich.*) Caudex elongated stout ascending; stipes 6–12 inches long, strong, erect, smooth, nearly naked; fronds oblong acuminate, 1–2 feet long, 6–9 inches broad, with 12–24 pairs of quite distinct articulated linear-oblong pinnæ on each side, the largest of which are 4–5 inches long, $\frac{1}{2}$ – $\frac{3}{4}$ inch broad, narrowed gradually towards the point and downwards to a narrow base; the margin finely incised; texture coriaceous; rachis rigid naked; veins very fine and close, not conspicuous; fertile pinnæ narrower; sori in a continuous line close to the midrib. *Rich. in Act. Soc. Nat. Par. i.* 114. *Hook. Syn. Fil.* 186.

Malacca.

(Also in tropical America, Australia, Borneo, and New Caledonia.)

3. BLECHNUM ORIENTALE. (*Linn.*) Caudex stout erect, clothed at the crown with fibrillose dark brown scales; stipes 4–8 inches long, strong, erect, scaly below; fronds 1–5 feet long, 6–36 inches broad, ovate, with very numerous nearly contiguous linear pinnæ on each side, which are about $\frac{1}{4}$ inch broad, narrowed to a long point, the bases quite distinct, the upper ones decurrent, a few of the lowest mere auricles; texture subcoriaceous; rachis and both surfaces naked; veins fine and close; sori in a long continuous line close to the midrib. *Lin. Sp. Pl.* 1535. *Hook. Syn. Fil.* 186. *Bedd. F. S. I. t.* 29.

Throughout India, Ceylon, and the Malay Peninsula, up to 6,000 feet in the south, and 4,000 feet on the Himalayas.

(Also in the Malay Islands, China, Polynesia and Australia.)

4. BLECHNUM FINDLAYSONIANUM. (*Wall.*) Caudex very small; stipes short; frond ample, 2–4 feet, ovate-lanceolate, sub-

coriaceous, pinnated; pinnæ erect-patent, rather distant, 6-12-14 inches long, 1-1½ wide, oblong or elongate-oblong, sessile, obliquely cuneate at the base, entire, upper ones much decurrent, but (except the two upper ones) not coadunate; the apex suddenly and sharply acuminate; several of the lowest pairs abortive, reduced to small hard scales; the surface glossy; veins obscure, compact, parallel; sori close to the costa continuous, narrow-linear; involucre also very narrow and indistinct. *Wall. Cat. n. 2172. Hook. Sp. Fil. iii. 53, and Syn. Fil. p. 187. Bedd. F. B. I. t. 249.*

Penang and the Malay Peninsula. (*Perhaps only a very broad-leaved form of orientale.*)


GENUS XXXVI.—BLECHNIDIUM. (*Moore.*)

(Resemblance to *Blechnum*.)

Like *Blechnum*, but the veins anastomosing and forming large arched areoles.

1. *BLECHNIDIUM MELANOPUS.* (*Hook. under Blechnum.*) Stipes black erect, 6 inches long, naked; fronds broadly falcate-lanceolate, glossy, sub-coriaceous, pinnate; pinnæ very approximate from a rather broad adnate base, linear-oblong falcate, somewhat acute, the margin entire, narrowly revolute, uppermost ones gradually confluent into a lanceolate elongated lobe, the lowest 3-4 pair suddenly diminished, short, very obtuse, sub-auriculate at the superior base; veins internal reticulated, forming large angled areoles next the costa, narrower oblique ones often extending to the margin; ultimate veinlets free and clavate at their apex just within the margin; sori quite close to the costa, continuous, not reaching to the apex of the pinnæ, lower half of the rachis beneath black. *Hook. Sp. Fil. iii. 64. Syn. Fil. p. 186. Bedd. F. B. I. t. 50.*

Khasya Hills.



GENUS XXXVII.—WOODWARDIA. (*Smith.*)

(In honour of T. J. Woodward, an English Botanist.)

Sori linear or linear-oblong, sunk in cavities of the frond, placed in single rows parallel with and contiguous to the midribs of the pinnæ and pinnules; indusium sub-coriaceous, the same shape as the sorus, closing over the cavity like a lid; veins forming a series of costal arches; the rest free or anastomosing; fronds ample and bipinnatifid.

1. WOODWARDIA RADICANS. (*Smith.*) Stipes elongate, with large lanceolate acuminate paleaceous scales at the base; fronds ample, pinnate, subcoriaceous; pinnæ distant, often a foot or more long, petiolate, deeply pinnatifid broad-lanceolate acuminate; laciniae ovate-lanceolate, acuminate, subfalcate, spinulose-serrate towards the apex; veins reticulate towards the main costa (of the pinnæ), a single series of areoles outside of the sori thence free to the margin; sori parallel with and near to the costule, short oblong, sunk in the cavity of the sorus, which has an elevated margin; involucre vaulted coriaceous; rachis here and there producing large scaly germinating gemmæ at the base of some of the pinnæ. *Smith, Act. Taur.* v. p. 4-12. *Hook. Sp. Fil.* iii. 66. *Syn. Fil.* 138. *Bedd. F. B. I. t.* 88.

Himalayas, from Kashmir to Bhotan, 3,000-8,000 feet elevation; common in Khasya, 4,000-5,000 feet.

(Also in Java, S. Europe, Macaronesia, California and Mexico.)

GENUS XXXVIII.—DOODIA. (*R. Br.*)

(In honour of Samuel Doody, a British cryptogamist.)

Sori oblong or sub-lunulate, in one or more series, arranged transversely with the veins forming areoles which are superficial, not sunk; involucre membranaceous, of the same form as the sori, opening towards the costa and lying parallel with it; veins uniform, the



lower ones arcuately anastomosing, forming elongated costal areoles (one or more series), the marginal venules free; fronds pinnate or pinnatifid; segments or pinnæ sharply serrated, rhizome sub-globose.

1. *DOODIA DIVES*. (*Kunze.*) Rhizome obliquely ascending, paleaceous, rootlets clothed with black fibre; stipes angled brown, very paleaceous near the base; fronds coriaceous, glabrous ovate-oblong, long caudate at the apex, pinnate at the base, with some of the lower pinnæ auricled, pinnatifid above; pinnæ and segments patently divergent, oblong, margins pungently serrated; pinnæ or segments of the fertile fronds often distant; sori in one or two series on each side of the costa of the pinnæ and segments, and also on the wings of the rachis, lunate, often more or less confluent. *Kze. in Schk. Fil. Suppl.* ii. p. 12. t. 185. *Hook. Sp. Fil.* iii. 73. *Syn. Fil.* 189. *Bedd. F. S. I.* t. 222.

Ceylon, Hackgalla and the central provinces above 4,000 feet elevation.

(Also in Java.)

TRIBE VIII.—ASPLENIEÆ.


Sori attached to the veins, oblique with regard to the costa or occasionally sub-parallel with it, linear or oblong or horseshoe-shaped; indusium the same shape as the sorus, when single opening over the midrib, sometimes double.

GENUS XXXIX.—THAMNOPTERIS. (*Presl.*)

(*Thamnos*, a thicket; *pteris*, fern.)

Sori unilateral linear; indusium linear, straight, single, free; veins forked; venules parallel; their apices combined by a dense continuous marginal vein; fronds undivided, simple, lanceolate to broad-lanceolate.

THAMNOPTERIS NIDUS. (*Linn.*) Fronds very variable in breadth, lanceolate acute or acuminate at the apex, taper-



ing gradually into a short stem, the edge entire, the midrib rounded on the back ; texture coriaceous ; veins fine and parallel, half a line apart ; sori reaching about half-way towards the margin. *Lin. Sp. Pl.* 1537. *Hook. Syn. Fil.* 190. *Bedd. F. B. I. t.* 197.

North India ; the Malay Peninsula ; Ceylon.

(Also pretty general throughout tropical Asia, Mauritius, Madagascar, &c.)

Typical *Nidus* has the fronds up to 6 feet long and 5-6 inches wide, and has not been found in the Madras Presidency.

VAR. β MUSÆFOLIA. (*Mett.*) Fronds up to 6 feet long and 1 foot broad ; sori extending nearly to the margin. *Mett. Asp. p.* 85. *Hook. Sp. Fil.* iii. 78.

Malacca, Singapore.

(Also in the Philippines.)

VAR. γ PHYLLITIDIS. (*Don.*) Fronds up to 2 feet long and 2, rarely 3 inches broad. *Don. Prod. Nep.* 7. *Hook. Sp. Fil.* iii. 80. *Bedd. F. S. I. t.* 123.

Western forests of the Madras Presidency, very common ; Himalayas ; Khasya ; Malay Peninsula.

(Also in Java Luzon, and Samar.)

The above have been considered three different species by many botanists, they are now generally all united under *Nidus* ; there appears to be no real difference except as to size, and it is doubtful if this is constant, or that the supposed varieties do not run into each other.

2. THAMNOPTERIS GREVILLEI. (*Wall.*) Fronds 12-18 inches long by 2-3 broad, lanceolate-spathulate, narrowed to an acute apex and suddenly below to a broad wing to the stipe, which grows very gradually narrower downwards, the margin entire ; texture coriaceous ; midrib keeled below ; veins nearly horizontal, those in the body of the frond about $\frac{1}{2}$ line apart ; sori on most of them extending within a short distance of the edge. *Wall. Cat.* 1036. *Hook. Sp. Fil.* iii. 80. *Bedd. F. B. I. t.* 66.



Birma and Mishmee.

The spatulate fronds with the winged stipe give this fern a very different appearance to any form of *Nidus*.

3. *THAMNOPTERIS SIMONSIANA*. (*Hook.*) Fronds 12–18 inches long, 1–1½ inch broad, tapering above into an acuminate point, and very gradually below into a short stem, the margin nearly entire; midrib only rounded and prominent below; texture coriaceous; veins erect-patent, about ½ line apart; sori copious, reaching from the midrib to within a short distance of the edge. *Hook. Syn. Fil.* 191. *Bedd. F. B. I. t.* 247.

Khasya and Jaintea hills up to 4,000 feet; Chittagong; Madras Presidency, on the hills west of Vizagapatam, 3,000 feet.

Mr. Clarke is in error in saying that it is found in Malabar; he unites it with Nidus, but it would hardly be safe to do so unless it be proved to vary in cultivation.

GENUS XL.—ASPLENIUM. (*Linn.*)

(*A*, privative; *splen*, spleen; in allusion to its medical properties.)

Sori dorsal or sub-marginal, linear or oblong; indusium similar in shape, straight, single, plane or tumid, bursting along the outer edge; veins free.

* *Fronds quite entire.*

1. *ASPLENIUM ENSIFORME*. (*Wallich.*) Stipes tufted 1–3 inches long, firm, erect, scaly below; fronds 8–20 inches long, ¾–1 inch broad, much acuminate; margin entire or nearly so, the lower part of frond very gradually narrowed; texture coriaceous; veins immersed, inconspicuous, erect-patent, usually once forked; sori broad, reaching nearly to the midrib and margin. *Wall. Cat.* 200. *Hook. Syn. Fil. p.* 191. *Bedd. F. S. I. t.* 125. *A. stenophyllum*, *Bedd. F. B. I. t.* 147.

Himalayas; Gurwhal to Bhotan, 4,000–9,000 feet elevation Ceylon; Birma; Western forests of Madras Presidency. The South Indian, Ceylon, and Birma plant has fronds generally about 8 inches long and nearly 1 inch wide. Some Himalayan examples have very long and very narrow fronds, but they cannot be separated

... long below, very
 undulate, crenate, texture subcoriaceous;
 usually once forked; sori reaching from the
 the way to the margin. *Hook. Syn. Fil.* 193



Sikkim, b
 feet elevation
 Khasya, 4,000
 Mergui and Ta

*There is als
 a fern from Per
 with this, excep
 stipe 6-9 inches.*

*** * Fronds**

3. ASPLENIUM

Caudex short, c
 rooting, squamos
 as is the very s
 long) stipe, and
 beneath; fronds
 span long, charta
 pale rusty green

Nº 71
ASPLENIUM ENSIFORME.
(Willd.)

lanceolate

extending to west frontier of Kashmir, and eastward to Sikkim, where it is rare.


(Also in Abyssinia.)

* * * *Fronds pinnate.*

4. *ASPLENIUM VIRIDE.*
(Hudson.) Stipes densely tufted; often brown or blackish, 2-4 inches long, naked; fronds 4-6 inches long, $\frac{1}{2}$ inch broad with 12-20 sub-sessile pinnæ on each side, which are ovate or elliptic, the upper edge narrowed suddenly at the base, the lower obliquely truncate, the outer part deeply crenated; texture herbaceous; rachis naked green; veins sub-flabellate; sori copious linear-oblong oblique. *Huds. Fl. Ang.* 453. *Hook. Syn. Fil.* 195. *Bedd. F. B. I.* 64.

Kashmir, 12,000-13,000 feet; Gulmurg, Tilail, Kumaon.

(Also in Europe, Asia, and N. America in Arctic and Alpine regions.)



Nº72.

ASPLENIUM ALTERNANS. (Wall.)

5. *ASPLENIUM TRICHOMANES.* (Linn.) Stipes densely tufted, 1-4 inches long, naked glossy brown or black; fronds 6-12 inches long, about $\frac{1}{2}$ inch broad, with 15-30 opposite pairs of sessile horizontal pinnæ, which are $\frac{1}{4}$ - $\frac{3}{8}$ inch broad, $1\frac{1}{2}$ -2 lines deep, the edge slightly crenate, the two sides unequal, the upper one the

3,000–6,000 feet elevation.

(Also in all four quarters of the world,

6. ASPLENIUM NORMALE. (*Don.*)

tufted, wiry, dark purple, polished; fronds 1
broad, with very numerous close placed pinnæ
are $\frac{3}{4}$ –1 inch long, $\frac{1}{4}$ inch broad, deflexed;
incised-crenate, the upper side auricled at
the base, the lower truncate in a straight line
rachis glossy; veins mostly once forked; sori
in direction, some at right angles with the
them, or in one row, or only 1–2 to each
Nep. p. 7. Ho. k. Syn. Fil. p. 197. Asplen
Bedd. F. S. I. t. 133.

Madras Presidency, in the forests of
3,000–6,000 feet elevation; Ceylon, above
Himalayas from Nepal to Bhotan, 4,000–8,000
5,000 feet. Mr. Clarke says some forms are
Trichomanes, others to monanthemum, and
Malayan examples are much more variable than

(Also in Canton.)

VAR. β . Pinnæ more numerous and

numerous, approximate, horizontal sessile, oblong obtuse, upper half, obscurely serrated, cuneate at the base, superior base truncated subauriculate, inferior margin cut off as it were by a straight line; terminal pinna elongated, pinnatifid below; costa and veins sunk, obsolete; sori 2-6, oblong oblique, occupying the disc of the frond between the middle and the apex; involucre firm, membranaceous; rachis ebony-black, terete, rigid, sparsely setose. *Hook. Sp. Fil.* iii.

143. *Syn. Fil.* p. 198. *Bedd. F. B. I. t.* 288.

Penang.


8. *ASPLENIUM SEPTENTRIONALE*. (*Lin. under Acrostichum*). Small caudex creeping, densely radiculose scarcely paleaceous; stipes numerous, tufted, 3-6 inches high, erect, flexuose, green brown at the base; fronds 1-2 inches long, coriaceous, glabrous, pinnated; pinnæ $\frac{3}{4}$ -1 inch long; long-petioled, linear or linear-lanceolate, subunguiculate, often very acute or acuminate, rarely solitary, generally 2-3, alternate entire or forked; lateral segments small subulate, rarely laciniated (2-3 acuminate segments); veins forked parallel; sori very long; involucre also much elongated, attached near the margin. *Hook. Sp. Fil.* iii. 174. *Syn. Fil.* p. 198. *Bedd. F. B. I. t.* 60. *Acrostichum septentrionale*, *Linn. Sp. Pl.* 1524. *Acropteris*, *Link.*

Northern India, Kashmir, 9,000-12,000 feet elevation; Gurwhal 8,000 feet.

(Also in Arctic and Alpine Europe, Asia, and America.)

9. *ASPLENIUM LONGISSIMUM*. (*Blume*.) Rhizome short-creeping; stipes tufted 3-12 inches long, strong erect, blackish, nearly naked; fronds 2-8 feet long, linear or lanceolate elongate, proliferous and rooting at the apex; pinnæ horizontal, very numerous, 2-4 $\frac{1}{2}$ inches long $\frac{1}{4}$ - $\frac{3}{8}$ inch broad, acuminate, the two sides nearly equal, with a distinct central midrib, the edge slightly toothed, the base on both sides often auricled; texture herbaceous; rachis dark-coloured strong, slightly scaly; veins forked oblique; sori numerous in two regular rows on each side the midribs, and reaching nearly to the edge. *Blume, En. Plant. Jav. Fil.* 178. *Hook. Syn. Fil.* 199.

sori opposite
 on each side, distinctly stalked, the lower
 with a long acumination, $\frac{1}{2}$ 1 inch by
 truncate-cuneate, often a little unequal, the
 colour pale green; texture coriaceous
 simple, or once forked; sori very re



N°73

reac
 sium
 2215
Beda
 Vulc
 Syn.

rarely
 pinna
 serrai
 (moun
 lays,
 4,000
 Penir

bitid ; fronds about 10 inches long by 2 inches broad. *Bedd. F. S. I. t. 127.*

Anamallays ; banks of Toracadu and elsewhere. I believe this to be only a form of *Wightianum*, though Mr. Baker is inclined to regard it as a distinct species.

11. *ASPLENIUM TENERUM*. (*Forst.*) Stipes tufted, firm, erect, greyish, naked, 4–11 inches long ; fronds 8–15 inches long, 3–4 inches broad, oblong-lanceolate ; pinnæ numerous on each side, stalked, horizontal, $\frac{1}{4}$ – $\frac{3}{8}$ inch broad, more or less bluntly rounded at the apex, the edge rather deeply crenated all round, the two sides unequal, the upper one narrowed, almost at a right angle, sometimes auricled, the lower one obliquely truncate ; texture herbaceous ; rachis compressed ; veins all simple except the lowest superior one, which is always forked except in the uppermost pinnæ ; sori numerous, regular, parallel, not reaching the margin or midrib. *Forst. Prod. p. 80. Hook. Syn. Fil. p. 201. Asp. elongatum* (Sw.), *Bedd. F. S. I. t. 224.*

Ceylon ; at Oodawella and Matale East, not common. *Mr. Wall sends a specimen with most of the pinnæ pinnatifid in their lower half nearly to the rachis.* Singapore, Penang.


(Also in the Philippines, Borneo, Java, Marianne Islands, Tahiti, and New Guinea.)

12. *ASPLENIUM LUNULATUM*. (*Sw.*) Stipes tufted, 2–4 inches long, firm, nearly naked, grey or ebeneous ; fronds 8–18 inches long, 1½–3 inches broad, narrowly lanceolate-oblong, with many horizontal pinnæ on each side, which are $\frac{1}{4}$ – $\frac{1}{2}$ inch broad, bluntish or acute at the apex, more or less deeply crenate throughout, the two sides unequal, the upper one at the base narrowed suddenly at about



ASPLENIUM TENERUM.
(*Forst.*)

winged pinnæ 30-40 pair, alternate or
 long, rounded at the apex, lower ones ofte
plenium camptorachis, Kze. *Asplenium bi*
S. I. t. 135.



1
 r
 C

 F
 lo
 pi
 th
 in
 ed
 (R
B.
 lay

rally proliferous at the apex ; pinnæ distant, alternate, 15–20 pair, stalked, ligulate-lanceolate acuminate, 3–5 inches long, 1–1½ inch broad, crenate, broadly rounded on the upper, cuneate-truncate on the lower side at the base ; lowest rather reduced ; texture firm ; surfaces naked, dark green ; veins erect-patent, fine distant, forked ; sori regular, parallel, 3–4 lines long, scarcely produced more than half-way to the margin ; indusium firm, broad, *Kze. in Lin. xxiv. p. 259. Hook. Syn. Fil. 484. Asp. persicifolium* (Hook.), *Bedd. F. S. I. t. 127.*

Nilgiris and Pulneys at the higher elevation ; Ceylon at Newera Elya, &c., above 5,000 feet elevation.

14. *ASPLENIUM HIRTUM*. (*Kaulf.*) Stipes tufted, 2–4 inches long, strong, erect, brownish, deciduously scaly ; fronds 12–24 inches long ; pinnæ very numerous, horizontal, 1–3 inches long, ¼–¾ inch broad ; the apex rounded or acute ; the margin broadly not deeply toothed, the base on the upper side conspicuously auricled, on the lower side rather suddenly but obliquely narrowed ; texture coriaceous ; rachis dark-coloured, strong, more or less fibrillose ; veins oblique, inconspicuous ; sori in two regular rows which do not reach the margin. *Kaulf. En. Fil. 169. Hook. Syn. Fil. p. 205. Bedd. F. B. I. t. 194.*

Sikkim at Lingtam near Darjeeling ; Moulmein.

(Also in the Mascareen Islands, Hong Kong, Philippines, and Solomon Islands.)

15. *ASPLENIUM AURITUM*. (*Sw.*) Stipes tufted, 4–8 inches long, firm, erect, grey, naked ; fronds 3–12 inches long ; pinnæ 10–20 distinctly stalked, horizontal, 2–3 inches long, the apex acute or blunt, either deeply pinnatifid or pinnate throughout, or only sharply toothed or variously lobed, especially on the upper side towards the base ; texture very coriaceous ; rachis firm, erect, naked ; veins close, oblique, inconspicuous ; sori in two broad rather oblique rows. *Sw. Syn. Fil. 78. Hook. Syn. Fil. 208. Bedd. F. S. I. t. 137.*

South India, rare ; Conoor, on rocks 5,600 feet elevation ; Anamallays, bed of the Toracadu river, 4,000 feet.

(Also throughout Tropical America and the West Indies, and in the Mascareen Islands.)

near apex, the margin
upper base and sometimes both the upper
auricled, giving the pinna a hastate shape
obliquely truncate; texture coriaceous;
the stipes; midrib rather indistinct; veins
oblique. *Hance, in Ann. Sc. Nat. Ser. v*
208. Asp. Beddomei, Mett. in Linnæa,
Bedd. F. S. I. t. 141.

Madras Presidency, on the Anamall
4,000 feet elevation; Sikkim, Jaintea hills
allied to the next species, and probably only
(Also in China)

17. *ASPLENIUM FALCATUM. (Lam.)*
long, erect, greyish, glabrous, or more or less
to 2 feet long or more, 4-8 inches broad; pinnae
alternate, sub-opposite or opposite, lanceolate
inch broad, the edges serrated or lobed, with
two sides unequal, and the lower one at the
texture coriaceous; rachis glabrous or fibrillose
sori in long irregular lines reaching nearly to
306. *Hook. Syn. Fil. 208. Bedd. F. S. I.*

Madras Presidency. Western mountains.

Localities much the same as *falcatum*. Mr. Clarke says not in Ceylon, but I have gathered many specimens there.

19. *ASPLENIUM CAUDATUM*. (*Forst.*) As in *falcatum*, but the sori generally confined to two parallel rows close to the midrib, or with a few other short oblique sori. *Forst. Prod. p.* 80. *Hook. Syn. Fil. p.* 209. *Aspl. contiguum, Bedd. F. S. I. t.* 140. *Probably only a variety of falcatum.*

S. India, on the Western mountains; Ceylon; Malay Peninsula. (Also in Australia, Africa, and Brazil.)

20. *ASPLENIUM GARDNERI*. (*Baker.*) Rhizome short, creeping; stipes grey, 2-6 inches with a few scattered small scales; fronds lanceolate, $\frac{1}{2}$ -1 foot long, 3-5 inches broad, often gemmiparous; rachis not winged; pinnæ 8-10 pair, distinct, ascending, lanceolate, short-stalked, acute, central ones 2-3 inches long $\frac{1}{2}$ - $\frac{5}{8}$ inch broad, toothed, square on the upper side, cuneate on the lower side at the base; veins erect-patent, forked, moderately close; sori regular, parallel, $\frac{1}{4}$ - $\frac{1}{2}$ inch long. *Hook. Syn. Fil. p.* 485. *Bedd. Suppl. Ferns. t.* 355.

Closely allied to macrophyllum, and probably only a variety of it. I have specimens from North India (not gemmiparous), which I do not know whether to refer to this or to macrophyllum. My Ceylon specimens are all gemmiparous, and the texture is generally thinner and the veins further apart.


Ceylon; Khasya Mountains, Umwai, 3,000 feet elevation.

21. *ASPLENIUM PARADOXUM*. (*Blume.*) Stipe firm, erect, grey, slightly scaly; fronds 1-2 feet long, with a large oblong-rhomboidal terminal pinna, and 8-12 lateral ones on each side, the lowest of which are 5-6 inches long, 1-1 $\frac{1}{2}$ inches broad, the apex acuminate and slightly toothed, the two sides unequal, the upper one narrowed suddenly almost at a right angle, the lower obliquely cuneate; texture coriaceous; rachis firm, erect, compressed, rather scaly; veins obscure, very oblique, usually once forked; sori reaching nearly from the midrib to the margin. *Bl. Fil. Jav. p.* 179. *Hook. Syn. Fil. p.* 209. *Aspl. zamioides, Hook. Sp. Fil. iii. p.* 114. *Bedd. F. B. I. t.* 193. Penang. The compressed rachis and short broader almost entire

pinnæ give this a very different look to *Gardneri* or *macrophyllum*. I have only seen two or three specimens and do not know how far it varies, but should not be surprised if it turned out to be only a variety of *macrophyllum*.

(Also in Java and Sumatra.)

22. *ASPLENIUM FORMOSUM*. (*Willd.*) Stipes tufted, very short, naked, polished, dark brown; fronds 12-16 inches long; pinnæ 20-31 pair (lower ones generally reduced), sessile, horizontal, $\frac{1}{4}$ inch long, by $1\frac{1}{2}$ -2 lines deep, the upper edge deeply cut, the



ASPLENIUM PARADOXUM. (*Blume.*)

(Also in Tropical America, the West Indies, and Africa, Angola, and Congo.)

23. *ASPLENIUM UNILATERALE*. (*Lam.*) Stipes scattered slender, naked, dark brown, polished or grey, and opaque, 4-8 inches long; fronds 6-15 inches long, lanceolate-oblong; pinnæ 10-30 pairs sub-sessile horizontal 1-3 inches long, by $\frac{1}{4}$ -1 inch broad, in shape almost dimidiate, two-thirds of the lower side being entirely cut away, the apex acute or bluntish, all except the truncate part incised, the upper half of the base narrowed nearly at a right angle; texture thinly-herbaceous; rachis usually polished like the stem, or green and

apex rather obtuse, the lower edge truncate in a straight line, so that about half the lower side is cut away; texture herbaceous; veins obscure, simple or forked; sori linear-oblong, short, oblique, confined to the lower side, or 1-4 on each side of the midrib. *Willd. Sp. Pl.* v. 329. *Hook. Syn. Fil. p.* 210. *Bedd. F. S. I. t.* 136.

South India, abundant in moist woods at the higher elevations on the Nilgiris, Pulneys and Anamallays; Ceylon, rare.

opaque ; veins once or twice forked ; sori not reaching either midrib or margin, often quite allantodioid in South Indian examples. *Lam. Enc.* ii. 305. *Hook. Syn. Fil.* p. 210, under resectum. *Bedd. F. S. I. t.* 132, under resectum.

Madras Presidency, in all the Western forests up to 5,000 feet ; Himalayas from Chumba to Bhotan 1,000–5,000 feet ; common eastward of Nepal ; Chittagong ; Ceylon ; Malay Peninsula.

(Also in the Malay Islands, Polynesia, Japan and Tropical Africa.)

VAR. β RIVALE. (*Bedd.*) Pinnæ 30–40 pairs, close set and much smaller than in the type, being only $\frac{1}{4}$ inch broad, less cut and almost a parallelogram in shape ; sori confined to the apex ; in habit resembling normale, var. β .

Pulney Hills and Anamallays, only in beds of rivers.

VAR. γ . UDUM. (*Atkinson.*) Pinnæ very transparent and membranaceous, smaller than in the type, very oblique, and the cutting often fimbriate. (*Bedd. Supp. Ferns. t.* 357, as var. decurrens.)

Himalayas ; Dalhousie, 5,500 feet elevation, and other localities, pendant from dripping rocks.

VAR. δ DELICATULUM. (*Parish.*) Fronds and pinnæ, much smaller than in the type, the latter less obliquely cut away on the lower side, and the main vein further from the margin. *Bedd. Supp. Ferns. t.* 358.

Tenasserim, on trees, 3,000 feet elevation. I have only seen one specimen sent by Mr. Parish.

24. ASPLENIUM HETEROCARPUM. (*Wall.*) Stipes scattered, firm, erect chestnut-coloured, polished ; fronds up to 2 feet, of which the stipe is 6–8 inches, membranaceous, glabrous, linear in outline, pinnæ numerous (40 or more), dimidiate, nearly the whole of the lower margin being cut away, about 1 inch long by $\frac{1}{2}$ inch broad ; upper margin and apex deeply cut, the segments being again bifid ; lower margin entire ; sori confined to the segments and directed upwards. *Wall. Cat.* 218. *Hook. Syn. Fil.* p. 210. *Bedd. F. S. I. t.* 131.

Madras Presidency, not uncommon in the Western forests, 4,000–7,000 feet elevation ; Himalayas from Nepal to Bhotan,

4,000–8,000 feet; Khasya, 3,000–5,000 feet; Malay Peninsula; Ceylon, central provinces, above 3,000 feet.

(Also in south-east China and Borneo.)

25. *ASPLENIUM LACINIATUM*. (*Don.*) Stipes and rachis with lanceolate or linear scales; fronds linear, up to nearly 2 feet long; pinnae numerous, alternate or subopposite, linear-oblong or half ovate, very much cut away, and unequal at the base, slightly pinnatifid, with the segments serrated, or more or less deeply pinnatifid nearly to the rachis and the segments distant; texture subcoriaceous; veins very oblique; sori copious. *Don. Prod. Fl. Nep. p. 8. Hook. Syn. Fil. p. 211. Bedd. F. S. I. t. 145. Aspl. planicaule* (Wallich), *Hook. Syn. Fil. 211. Bedd. F. S. I. t. 139.*

The typical *laciniatum* of North India has the pinnae elongated and much narrowed towards the apex, and deeply cut into distant segments; typical *planicaule* of S. India has very short, half ovate pinnae, scarcely at all cut; there are however many connecting links and it is impossible, I think, to separate them even as constant varieties. *Var. depauperatum* (*Clarke*) is also found in S. India but is, I believe, only a starved form.

Madras Presidency, all the Western mountains, 3,000–8,000 feet; Himalayas, Gurwhal to Bhotan; Khasya; Ceylon.


(Also in Japan.)

VAR. CRINIGERUM. (*Bedd.*) Pinnae all more deeply and finely cut than any typical *laciniatum*; stipe very short ($\frac{1}{2}$ inch), and very scaly; rachis densely scaly; pinnae with long hair-like scales on the nerves beneath.

South India, on the Balarangam Mountains (Mysore), and Myhendra Mountains near Ganjam, 3,000–4,000 feet elevation.

26. *ASPLENIUM PUMILUM*. (*Sw.*) Stipes tufted, 3–4 inches long, erect, glabrous or sparingly pubescent; fronds 2–6 inches long, broadly deltoid, the upper part sinuated only, the lower cut into distinct pinnae, of which the lowest pair is much the largest and often pinnatifid nearly to the midrib; texture herbaceous; sori very oblique, the lower ones sometimes $\frac{3}{4}$ –1 inch long. *Sw. Fl. Ind. Ccc. iii. 161.*

(West Indies and Central America.)



A. LATHROPYCNIS

ASPLENIUM HETEROCARPUM. (Walt.)

VAR. HYMENOPHYLLOIDES. (*Fee.*) Fronds smaller, pubescent or ciliate on both surfaces ; texture exceedingly thin, with prominent venation ; pinnæ obtuse. *Fee, 7th. Mem. 54. t. 15. f. 1. Clarke, N. India, p. 482.*

Mt. Aboo, Rajaputana, 4,000 feet elevation.

(Also in Abyssinia.)

* * * * *Fronds bi-quadripinnate.*

27. ASPLENIUM RUTA-MURARIA. (*Linn*) Stipes tufted, 2-4 inches long, slender, wiry, naked, ebeneous towards the base ; fronds 1-2 inches long, about 1 inch broad, glabrous, deltoid, cut down to the rachis into a few pinnæ on each side, the lower ones again cut down into spathulate cuneate pinnules, which are serrated round the outer edge ; texture coriaceous ; rachis firm, green, naked ; veins flabellate ; sori copious ; margin of indusium fimbriate. *Linn. Sp. Pl. 1541. Hook. Syn. Fil. 213. Bedd. F. B. I. t. 61.*

Kashmir.

(Also in Europe, North Africa, Thibet, Siberia, and the United States.)

28. ASPLENIUM PEKINENSE. (*Hance.*) Stipes tufted, 1-3 inches long, with a few linear scales near the base ; fronds 2-6 inches long, narrowly oblong, shining, glabrous, with several lanceolate-deltoid pinnæ on each side, which are cut down to a compressed rachis into several ovate rhomboidal pinnules on each side, which are again pinnatifid (hence it is bipinnate sub-tripinnate), ultimate segments linear-cuneate, sharply forked or toothed at the apex, entire below ; sori 2-6 to a pinnule, ultimately confluent ; involucre entire. *Hance in Seem. Journ. Bot. v. 262. Hook. Syn. Fil. p. 213. Clarke, F. N. I. t. 56. f. 2.*

Kashmir ; Jhelum Valley, 2,000-2,500 feet elevation.

(Also in China and Japan.)

29. ASPLENIUM ADIANTUM-NIGRUM. (*Linn.*) Stipes tufted, 6-9 inches long, nearly glabrous, polished ; fronds bi-tripinnate, deltoid to lanceolate, 6-12 inches long, 4-6 inches broad at the base ; pinnæ numerous, the lower ones deltoid, their lower pinnules again stalked and completely pinnate and lanceolate deltoid in

shape, the ultimate segments ovate or oblong, acutely serrate, texture coriaceous; veins obscure, oblique; sori copious, involucre with an entire margin. *Linn. Sp. Pl.* 1541. *Hook. Syn. Fil. p.* 214. *Bedd. F. B. I. t.* 62.

Kashmir, 5,000–8,000 feet, extending to Dalhousie and Chumba.

(Also in Europe, North Asia, North Africa, South Africa and its islands, and Sandwich Islands.)

30. *ASPLENIUM FURCATUM*. (*Thunb.*) Stipes tufted, 4–8 inches long, clothed with deciduous woolly hairs, as is the rachis; fronds coriaceous, 6–18 inches long, 4–6 inches broad; pinnæ 12–20 pair, lanceolate-deltoid, $\frac{3}{4}$ –1 inch broad, cut down throughout nearly to the rachis into linear-cuneate pinnules, which are sharply serrated, on the outer edge; veins flabellate, deeply channelled; sori linear. *Thunb. Prod. Fl. P.* 172. *Hook. Syn. Fil. p.* 215. *Bedd. F. S. I. t.* 144.

South India, common on the higher Western mountains, Ootakamund, &c.; Ceylon, 5,000–7,000 feet.

(Also in Tropical America, Polynesian Islands, Australia, Cape Colony, Abyssinia, Mascareen Islands, and Canaries.)

31. *ASPLENIUM AFFINE*. (*Swartz.*) Stipes 6–12 inches long, firm, erect, grey, nearly naked; fronds 12–18 inches long, 6–12 inches broad; pinnæ numerous on each side, the lower ones lanceolate-rhomboidal, cut down to the rachis into numerous distinct rhomboidal pinnules, 4–6 inches long, $1\frac{1}{2}$ –2 inches broad, which are incised or deeply lobed; texture subcoriaceous; rachis firm, naked; veins subflabellate; sori copious, linear. *Sw. in Schr. Jour.* 1,800. 11. 56. *Hook. Syn. Fil.* 215. *Asp. spathulinum* (J. Sm.), *Bedd. F. S. I. t.* 226 (probably taken from a specimen of *furcatum* and not from *affine* as represented at Kew.)

Ceylon.

(Also in Mascareen Islands, Philippines, Borneo, Fiji, and Sandwich Islands.)

32. *ASPLENIUM NITIDUM*. (*Swartz.*) Stipes 1 foot long, firm, erect, greyish, naked; fronds 2–3 feet long, 1 foot broad; pinnæ numerous, lanceolate-deltoid, the lowest 6–9 inches long, 2–3 inches broad, cut down to the rachis into numerous stalked subdeltoid

pinnules, which are again cut down to the rachis into broad flabellate cuneate segments, serrated round the outer edge; texture subcoriaceous; rachis firm, grey, naked; veins channelled, flabellate; sori short, radiant. *Sw. Syn. Fil. p. 280. Hook. Syn. Fil. p. 215. Bedd. F. S. I. t. 148, 149. Aspl. laserpitiifolium, Bedd. F. S. I. t. 225.*

Sikkim, Bhotan, Assam; Khasya, Cachar, 1,000–4,000 feet elevation; Ceylon; Madras Presidency in the Western forests; Malay Peninsula.

(Also in Malay Islands and S. Africa.)

33. *ASPLENIUM FONTANUM*. (*Bernh.*) Stipes tufted, 2–4 inches long, wiry, slender, glabrous; fronds 3–6 inches long, 1–1½ inches broad, oblong-lanceolate; pinnæ numerous, the lower ones reduced, often somewhat remote, pinnate, pinnatifid or only toothed; texture herbaceous, rachis and surfaces bright green; veins oblique; sori copious, covering nearly the whole segment. *Bernh. in Schrad. Neu. Jour. 1 pt. ii. 26. Hook. Syn. Fil. p. 216. Bedd. F. B. I. t. 146.*

Himalayas, from Kashmir to Gurwhal, 5,000–9,000 feet.

(Also in Europe.)

VAR. β *EXIGUUM*. Frondsless divided and more narrow, and an ebeneous rachis. *Asp. exiguum, Bedd. F. S. I. t. 146.*

Nilgiris, Kulhatty; Himalayas.

34. *ASPLENIUM VARIANS*. (*Hook and Grev.*) Stipes tufted, 1–3 inches long, slender, naked, greenish; fronds 4–6 inches long, 1 inch broad, oblong-lanceolate; pinnæ 8–12 pair, the lower ones subdeltoid ½–¾ inch long, ¼–⅜ inch broad, cut down to the rachis into a few cuneate flabellate pinnules, the lowest two lines across, sharply toothed on the outer edge; texture thin, herbaceous; rachis, slender, naked, green; veins, one carried into each tooth; sori copious. *Hook. Syn. Fil. 216. Bedd. F. S. I. t. 129.*

South India, common on the Nilgiris and the mountains on the west side, 3,000 feet upwards; Himalayas, 6,000–10,000 feet, rarer eastwards; Ceylon, 3,000 feet and upwards.

A form found at Kulhatty on the Nilgiris, quite runs into the European lanceolatum. Another form on the Tinnevely hills west of

Courtallum has short deltoid fronds fully 2 inches across at the base, and broader pinnules.

(Also in South Africa, North China and Japan.)

35. *ASPLENIUM BULBIFERUM*. (*Forst.*) Stipes tufted, 6–12 inches long, firm, erect, scaly near the base, sometimes hairy throughout; fronds 1–4 feet long, oblong to ovate-lanceolate, 2–3 pinnate; pinnæ many pair, horizontal, often proliferous from the upper surface, the largest eight inches long, and two inches broad, cut down to a compressed winged rachis, into many lanceolate-deltoid pinnules which are cut into slightly toothed linear oblong flaccid segments; texture herbaceous; veins firm, pinnate; sori oblong, large, often filling the whole segment. *Forst. Fl. Ins. Aust. Prod.* 80. *Hook. Syn. Fil. p.* 218. *Bedd. F. B. I. t.* 65, as *bullatum* (Wall.)

Nepal, Sikkim, Bhotan, Khasya, 2,000–6,000 feet; Penang.

(Also in Australia, New Zealand, Mexico, New Caledonia, Natal, and the East African Islands.)

36. *ASPLENIUM TENUIFOLIUM*. (*Don.*) Stipes tufted, 3–6 inches long, slender, glabrous, or with a few hairs; fronds green, thinly herbaceous, up to 20 inches long, lanceolate, 3–4 pinnate; pinnæ many pair, the lower ones 3 inches long by $\frac{3}{4}$ inch broad, lower pinnule subdeltoid, cut down to a narrowly winged rachis into spatulate segments, quarter inch or less broad, sharply cut on the outer edge; veins solitary; sori 1–2 to a segment. *Don. Fl. Nep.* 8. *Hook. Syn. Fil. p.* 220. *Bedd. F. S. I. t.* 130.

South India, higher ranges of the Nilgiris, Pulneys, and Anaimallays; Ceylon at Newera Elya; Himalayas, Nepal to Bhotan 5,000–9,000 feet; Khasya, 4,000–5,500 feet; Birma.

***** *Ultimate division of the frond, narrow, linear; sori linear or linear-oblong, marginal or submarginal. Darea of authors.*

37. *ASPLENIUM RUTÆFOLIUM*. (*Kunze.*) Stipes tufted, 6–9 inches long, compressed, greenish, glabrous; fronds 6–15 inches long, narrowly oblong, glabrous, bipinnate, the rachis often prolonged, naked, and rooting at the apex; pinnæ 12–20 pair, the lowest sub-deltoid, two inches or more long, cut down to the rachis many erect-patent distant pinnules on each side, the lowest on



42 C. T. P. & C. P. 1913.

ASPLENIUM TENUIFOLIUM. (Dois)

the upper side again cut into erect-patent linear segment
on the under side suppressed ; texture coriaceous ; rachis
sori linear marginal. *Kze. in Linnæa*, x. 521. *Hook.*

222. *Bedd. F. S. I. t.* 138.

South India, on the Tinnevely Mountains ; Ceylon, S
Forest ; Himalayas, Bhotan, and Mishmee ; Khasya.

(Also in South Africa, Japan and Fiji.)

38. ASPLENIUM BELANGERI. (*Kze.*) Caudex
erect ; stipes densely tufted, 4 inches to a span long,
upwards, greenish ; fronds 1-1½ feet long, narrow-oblong
almost forming a parallelogram), acuminate only at
sparingly stellate-squamulose beneath, here and there
subchartaceous, pale green, bipinnate, primary pinna
inches long, horizontally patent, singularly uniform in
length, linear obtuse, nearly sessile, again pinnated (or
said to be deeply pinnatifid), pinnules or segments 2-
linear-clavate, or subspathulate, obtuse, straight, or a
decurrent, with the rachis simple monosorous, or the lower
one is bifid or subpinnatifid ; veins solitary central ;
conspicuous, occupying nearly the length of the pinnule
involucres of the same colour and texture almost of the
rachis is compressed, partial ones distinctly winged. *Hook.*

209. *Syn. Fil. p.* 223. *Bedd. F. B. I. t.* 287.

The Malay Peninsula.

(Also in Java, Sumatra, and Borneo.)

GENUS XLI.—ATHYRIUM. (*Roth.*)

(*Derivation unknown.*)

As in Asplenium, but the involucres, or at least main
more or less curved, often horseshoe-shaped, rarely quite
in *Isoetes*.

1. ATHYRIUM SPINULOSUM. (*Max.*) Stipes 6-12
scattered, firm, erect, straw-coloured, furnished more
pale brown lanceolate scales ; fronds 6-12 inches, deltoid
long, 3-pinnate ; pinnæ 9-12 pairs, lowest the largest,



Asplenium rutæfolium

ASPENIUM RUTÆFOLIUM. (Lam.)

Broad, ovate-lanceolate ; pinnules lanceolate, cut down to the rachis on each side into 6–9 oblong rhomboidal segments two lines long, one line broad, which are sharply toothed or crenated ; texture herbaceous ; rachis naked or nearly so ; sori 2–10 to a segment ; indusium slightly curved, or horseshoe-shaped, very small and thin. *Cystopteris*, *Max. Fl. Amur.* 340. *Baker in Hook. Syn. Fil.* 225, under *Asplenium*. *Athyrium subtriangulare* (Hook), *Bedd. F. B. I.* t. 242. *The supposed species, or variety, subtriangulare, does not differ from spinulosum, except that the teeth are less spinulose ; it cannot, I think, be considered even a variety.*

Sikkim 10,000–12,000 feet ; Lachen, Samding, Yeumtong.

(Also in Manchuria, Amurland and Korea.)

2. *ATHYRIUM ATKINSONI*. (*Bedd.*) Stipes 6–12 inches long, erect, straw-coloured, furnished towards the base with a few long narrow pale-coloured scales ; fronds 10–12 inches long, triangular, 10–12 inches broad at basal pinnæ, gradually narrowing upwards, tripinnate, with the tertiary pinnæ deeply pinnatifid ; pinnæ 7–8 pairs, the lowest much the largest, about three inches broad ; texture herbaceous, naked, secondary pinnæ with the rachis winged, but the pinnules (tertiary pinnæ) often distinctly petioled, at least in the lower portion of the pinnæ, pinnatifid $\frac{1}{2}$ – $\frac{2}{3}$ down, the segments being more or less crenated ; sori one to each segment on the basal vein, linear, curved or almost reniform, never double ; veins terminating just within the margin. *Bedd. Fern. Sup. p.* 11, t. 359.

Sikkim at high levels.

(*Clarke's variety Andersoni* is *Athyrium fimbriatum*, var. *sphaeropteroides*.)

3. *ATHYRIUM HOHENACKERIANUM*. (*Kze.*) Stipes tufted, 2–4 inches long, clothed throughout rather closely with linear-subulate scales ; fronds 6–18 inches long, $1\frac{1}{2}$ –3 inches broad, ovate-lanceolate ; pinnæ in numerous pairs, lanceolate, 1 – $2\frac{3}{4}$ inches long, cut nearly or quite to the rachis into ovate or oblong serrated pinnules ; texture herbaceous ; rachis naked upwards, scaly below ; sori copious ; involucre conspicuous, linear-oblong, curved or horseshoe-shaped. *Hook. Syn. Fil. p.* 225. *Bedd. F. S. I. t.* 151.

Madras Presidency in the Western Forests, from the plains up to 4,000 feet; Ceylon, Scinde.

4. *ATHYRIUM FALCATUM*. (Bedd.) Stipes 1-9 inches long with many linear golden scales at the base; fronds 6-14 inches long, with the rachis above dilated or winged, linear-lanceolate, narrowed at both ends, pinnae 15-20 on each side, sessile, alternate, 1-1½ inch long, by ¼-⅓ inch broad, falcate-ovate, deflexed, obtuse or acuminate, generally with a large obtuse auricle at both the superior and



ATHYRIUM FALCATUM. (Bedd.)

inferior base, above pinnatifid almost half down to the costa; segments obtuse, crenated; sori numerous on each side of the costa, at length confluent and covering the whole or nearly the whole of the under surface. *Bedd. F. S. I. t. 151. Asplenium drepanophyllum, Baker, Syn. Fil. 226.*

South India, Anamallay Hills, dry grassy places 5,000 feet. Myhenda Hill near Berhampore, 4,500 feet; Mahableshwar; Parasnath 4,000 feet.

5. *ATHYRIUM THELYPTEROIDES*. (Michx.) Stipes 6-8 inches or more, furnished at the base with large lanceolate thin light coloured scales; fronds generally about 3 feet long and 8 inches broad, lanceolate, gradually narrowed towards the base; pinnae numerous, erect-patent, ¾-1 inch broad, cut three-quarters of the way to the rachis or more into oblong blunt crenated segments; sori in close regular rows, short, oblong; indusium persistent. *Michx. Fl. Bor.-Am. 11. 265. Hook. Syn. Fil. p. 226. A. allantodioides, Bedd.-F. B. I. t. 221, not F. B. I. t. 68, which is a form of Diplazium sorzogonense.*

Himalayas from Kashmir to Bhotan, common in many places.
(Also in North America, Canada, Amur land.)

6. *ATHYRIUM MACROCARPUM*. (*Blume.*) Stipes 6–9 inches long, firm, erect, straw coloured, scaly below ; fronds up to three feet long, generally lanceolate in outline ; pinnæ 20–30 pair, very variable in size and cutting, sometimes less than one inch long, and only pinnatifid, usually about four inches, and sometimes 9–10 inches long, again pinnate, with the pinnules deeply pinnatifid, and as large or much larger than the whole pinnæ in the less compound forms ; texture herbaceous, shining, striate beneath the lowest lobe on the upper side often larger ; margin toothed ; involucres very large, often reniform (as in *Lastrea*), but always mixed with some that are linear (asplenioid), and horseshoe-shaped, margin more or less fimbriate. *Aspidium macrocarpum*, *Bl. En. Fil. Jav.* 162. *Asplenium*, *Hook. Syn. Fil. p.* 227. *Bedd. F. S. I. t.* 152 and 153. *A. squarrosum*, *Wall. Cat.* 356.

South India, very common on the Western Mountains, above 3,000 feet ; Ceylon ; Himalayas, Gurwhal and Bhotan 2,000–9,000 feet ; Khasya ; Birma and the Malay Peninsula.

Clarke's variety, *Atkinsoni*, is one of the simplest forms, and is very common on the Nilgiris, mixed with the more compound examples and connected by intermediate forms. *Beddome's macrocarpum*, var. β , *F. S. I. t.* 153, is at first sight very distinct looking, and is often collected as a *Lastrea* ; it is, however, only a state in which the fructification is less and the lamina of the frond more developed than usual.

(Also in Malay Islands, China and Japan.)

VAR. β . 1-PINNATA. (*Clarke.*) Fronds linear ; pinnæ short, slightly crenate, in shape much like some forms of the *Polystichum auriculatum*.

Khasya, 3,000–4,000 feet. I have only seen this in Mr. Clarke's collection, and it looks distinct from any of the simpler forms of *macrocarpum* from Southern India, the pinnæ being much less cut.

7. *ATHYRIUM NIGRIPES*. (*Mett.*) Caudex stout, often 2-3 inches out of ground; stipes tufted, firm, round, glabrous; fronds 8-30 inches long, quite deltoid, with the lowest pinnæ largest, to oblong, lanceolate, herbaceous, not plicate-striate beneath; pinnæ about 8 inches long, by $2\frac{1}{2}$ broad, (but variable in size) pinnules often deeply pinnatifid to only a winged rachis; segments crenate or serrate, not spinulose; sori linear-oblong, only slightly curved in two oblique rows towards the midrib of the pinna; involucre persistent. *Mett. Farng. Aspl.* 195. *Hook. Syn. Fil.* 227, not *Bedd. F. S. I. t.* 157, which is *selenopteris*.


Himalayas, from Kashmir to Bhotan, 3,000-11,000 feet elevation, very common about Darjeeling; Khasya 2,000-6,000 feet. Not in Southern India or Ceylon, the former plant being *selenopteris*, (Kunze), the latter *gymnogrammoides*.

VAR. β *CLARKEI*. (*Bedd.*) Fronds linear-lanceolate, up to 3 feet long, and 3-4, rarely 6 inches, broad, gradually narrowed at both ends, generally rooting from a bud on the upper side of the rachis; rachis of pinnæ, costa, and veins, sometimes furnished with a few weak setæ. *Athyrium Clarkei*, *Bedd. Sup. Ferns, t.* 360.

Nepal, Sikkim, common at Surail near Darjeeling, 6,000 feet; Kulhoit Valley, 6,000-7,000 feet. I feel quite certain now that this is only an elongated variety of *nigripes*, and Mr. Clarke agrees.

(Mr. Clarke's *nigripes*, var. *dissecta*, appears to be typical *nigripes* rather more cut than usual.)

8. *ATHYRIUM SELENOPTERIS*. (*Kunze.*) Caudex stout, erect; stipes tufted, often 10 inches long, stramineous; fronds 10-20 inches long, lanceolate, somewhat attenuated towards the base, never deltoid; pinnæ 2-6 inches long, rather distant; texture rather flaccid, pinnatifid to a winged rachis in the smaller forms, or almost quite pinnate in the larger forms, the wing of the rachis being almost obsolete, the pinnules deeply pinnatifid, but with a somewhat decurrent base, ultimate segments more or less crenated, or even serrated; sori oblong or curved. *Kunze, Mett. Farngatt. Asp.* 196. *Asp. nigripes*, *selenopteris*, *Clarke, F. N. I. p.* 491. *Athyrium aspidioides*, *Hook. Syn. Fil. p.* 228. (in part) as far as the Nilgiri specimens are concerned.



ATHYRIUM SILENOPSIS (Kuntze.)

not the Madagascar plant of Schlecht. *Athyrium pectinatum*, *Bedd. F. S. I. t. 155* (not Wall.) *Athy. nigripes*, *Bedd. F. S. I. t. 157*.

Madras Presidency, Western Mountains, Nilgiris, common at 6,000–8,000 feet. Pulneys, Tinnevely Mountains; Ceylon at Newera Elya, and Maturatte, &c.

9. *ATHYRIUM GYMNOGRAMMOIDES*. (K7.) Fronds large, 5–6 feet long, bi-tripinnate, primary pinnæ 12–16 inches long, oblong-lanceolate, secondary pinnæ up to $4\frac{1}{2}$ inches long, by $1\frac{1}{2}$ inch broad, either deeply pinnatifid to a winged rachis, or pinnate, the wing being obsolete; ultimate segment oblong, lobed one-third of the way down; sori near the midrib. *Mett. Aspl. p. 193, t. 6, fig. 13–14*. *Bedd. F. S. I. t. 156*. *Thw. En. Pl. Zey. p. 335*.

Ceylon, central provinces 5,000–6,000 feet, nearly allied to *selenopteris*. Mr. Baker includes this under *nigripes*, but it is not very near the Himalayan plant.

VAR. β *ERYTHRORACHIS*. (*Bedd.*) The naked stipes and rachis deeply channelled, bright purple; fronds deltoid, much smaller than the type, 6–10 inches long, without the stipe, lowest pinnæ 3–4 inches long, the partial rachis channelled and winged on the upper side, the pinnules petiolate, $\frac{5}{8}$ inch long, auricled at the superior base and somewhat pinnatifid at the inferior base, crenate or subentire above, veins pinnate and forked terminating within the margin; sori asplenoid (never double), a few more or less curved. *Bedd. Sup. Ferns, p. 12*. *Athy. nigripes*, *Thw. En. Pl. Zey. p. 384*.

Ceylon, about Newera Elya.

This is probably a distinct species, it is included in the Kew bundle of *nigripes*, but is nearer to the Nilgiri *selenopteris*, some specimens of which are also in the Kew packet of *nigripes*, whilst others exactly similar are placed with the South African *aspidioides*. Mr. Thwaites has also referred it to *nigripes*, but he may not have seen the Himalayan plant.

10. *ATHYRIUM FILIX-FEMINA*. (*Bernh.*) Fronds 1–4 feet, lanceolate, narrowed at both ends, membranaceous, green, bi-pinnate;

rachis soft, appearing triangular or furrowed when dry; primary pinnae narrow linear-oblong, hardly narrowed at the base; secondary pinnae $\frac{1}{2}$ –1 inch, oblong, patent at right angles to the rachis of the primary pinnae, sessile or decurrent, serrate or pinnatifid; margin bluntly or acutely toothed; involucre in two rows on the secondary pinnae, short, oblong, subpersistent. *Bernh. in Schrad. Neu. Jour. i. Pl. 11. 26. Clarke. F. N. I. p. 491.*

I follow Mr. Clarke here, as he has an intimate knowledge of this fern and its varieties in a growing state, he says that there are two main types of the Himalayan *Filix-fœmina*, viz. :—1st, with a succulent rachis, appearing triangular or grooved when dry, green, involucre subpersistent; 2nd, with a firm rachis, appearing round in the dried specimens, often red; involucre less persistent, often fugacious.

VAR. 1. DENTIGERA. (*Wall.*) Cutting nearly as in the European type (not nearly so fine as in *pectinata*), green, rachis succulent, grooved when dry; involucre subquadrate, or horseshoe-shaped, smaller and less persistent than in the European type. (*Polypodium dentigerum, Wall. Cat. 334.*)

Himalayas 6,000–11,000 feet, from Kashmir to Bhotan, common from Nepal westwards.

VAR. 2. PECTINATA. (*Wall.*) Very finely cut, tripinnate, 4-pinnatifid, bright green; rachis slender, but scarcely succulent or grooved when dry; involucre subquadrate, or short oblong, little horseshoe-shaped, subpersistent. *Wall. Cat. 231. Clarke, F. N. I. t. 57. Bedd. F. S. I. t. 154, small form.*

Himalayas, Sikkim to Gurwhal 2,000–5,000 feet elevation; Parasnath 4,000–5,000 feet; Mountains of the Godavery and Central India; Mahableshwar; Scinde; Mount Aboo.

VAR. 3. ATTENUATA. (*Clarke.*) 1-pinnate; base of the stipes densely clothed with broad lanceolate scales; rachis succulent when dry, grooved or triangular; fronds small, very narrow, much tapering at both ends; pinnae patent, very close together, deeply regularly pinnatifid into oblong serrated segments, scarcely a quarter inch long;

involucre small, subquadrate, hardly ever horseshoe-shaped, not very fugacious. *Clarke, F. N. I. t. 59. fig. 1.*

Kashmir, 10,000–12,000 feet, north of the main valley ; appears to be only a stunted form of *dentigera*.

VAR. 4. RETUSA. (*Clarke.*) 1–2-pinnate, rachis firm, round when dry ; fronds red, never very large ; involucre small, fugacious ; sori, scattered, round. *Cystopteris retusa, Decne in Jacq. Voy. 176, t. 177. Clarke, F. N. I. t. 59–2.*

Himalayas 10,000–15,000 feet, from Kashmir to Bhotan.

VAR. 5. FLABELLULATA. (*Clarke.*) 2-pinnate ; rachis firm, round when dry, fronds red, 1–3 feet ; secondary pinnæ deeply pinnatifid ; segments lacinate ; involucre small, fugacious ; sori scattered, round. *Clarke, F. N. I. t. 60.*

Sikkim, 13,000 feet ; Yakla, Jongri, scarcely distinct from *dentigera*.

VAR. 6. POLYSPORA. (*Clarke.*) 2-pinnate, or sub-3-pinnate ; rachis firm, round when dry ; fronds red, 1–3 feet ; involucre large, approximate, very persistent, often reniform ; sori large, often ultimately thickly covering the whole of the pinnules. *Clarke, F. N. I. t. 61. fig. 1.*

N. W. Himalayas, 6,000–10,000 feet ; Kumaon to Chumba.

Very distinct from the other varieties, and approaching *oxyphyllum*.

VAR. 7. PARASNATHENSIS. (*Clarke.*) 1-pinnate, sub-2-pinnate ; rachis triangular when dry ; fronds reddish, 1½ foot, narrow-oblong, scarcely attenuated at the base ; primary pinnæ 1½ inch, broadest at the base ; secondary pinnæ ¼–⅓ inch, acutely toothed, scarcely pinnatifid ; involucre short, oblong, in two oblique rows, very persistent. *Clarke, F. N. I. t. 61. fig. 2.*


Parasnath 4,000 feet elevation ; probably only a small stunted form of *retusa*.

11. *ATHYRIUM OXYPHYLLUM.* (*Hook.*) Stipes with many

reddish linear-lanceolate scales at the base; fronds up to 3 feet, oblong-lanceolate, generally 2-pinnate, or in some forms only pinnate; slightly narrowed, never attenuated at the base, coriaceous, rigid, somewhat shining, striated when dry; primary pinnæ often falcate, sometimes only pinnatifid half-way to the rachis, $1\frac{1}{2}$ –2 inches long, generally much larger; 2–4 $\frac{1}{2}$ inches long, and pinnate, the pinnules more or less hastate from one or both of the basal lobes being enlarged or auricled, sub-entire to pinnatifid, and often sharply serrated, particularly at the apex; indusium small, fugacious or wanting. *Hook. Syn. Fil.* 228. *Bedd. F. B. I. t.* 67.

The exinvolucrate variety, called *Kulhaitense* by Clarke, is *eberneum*. (Wall.) It only differs in the absence of the involucre; the 1-pinnate form with short pinnæ looks different at first sight, but gradually runs into the larger and more compound forms.

Himalayas, from Gurwhal to Bhotan, 4,000–11,000 feet; Khasya 3,000–11,000 feet. I do not feel certain that the Khasya specimen called *stramineum* J. Sm. really belongs here, though it is so sorted in the Kew bundles; it is very like some of the Nilgiris forms of *Athyrium selenopteris*.



ATHYRIUM BREVISORUM. (Wall.)

12. *ATHYRIUM BREVISORUM*. (Wall.) Stipes $1\frac{1}{2}$ foot and more long stramineous-brown below, glossy; fronds $1\frac{1}{2}$ –2 feet long, ovate-lanceolate, membranaceous, bi-tripinnate; primary pinnæ long-petiole remote erect-patent, a foot and more long, broad-lanceolate, acuminate; secondary pinnæ 5–6 inches long, lanceolate, nearly sessile, pinnated; pinnules numerous, approximate, oblong-lanceolate, much acuminate, horizontally patent, 1 – $1\frac{1}{2}$ inch long, coarsely serrated, rarely subpinnatifid, the serratures very acute, almost

mucronate, lowest pinnules more distant on short petiolules, intermediate ones more approximate, the lower base decurrent, upper ones confluent; veins pinnated; veinlets forked once or more; sori generally in two rows, oblique near the costule in the pinnules that are pinnatifid, extending to the lobes, small, short-linear, straight or lunate; involucre slightly convex, jagged at the margin; main rachis stramineous, glossy, flexuose. - *Hook. Sp. Fil.* iii. 229. *Wall. n.* 220, (not *Mett.*) *Bedd. F. B. I. t.* 241.


Mishmee; Taongdong Mountains near Ava. The figure is taken from the Ava specimen collected by Wallich, which is bipinnate, the Mishmee specimen is tripinnate.

13. *ATHYRIUM FIMBRIATUM.* (*Wall. under Aspidium.*) Stipes stout, darkish brown, glossy, 1 foot and more long, copiously paleaceous, most so towards the base, with large glossy, tawny, lanceolate-subulate scales, fronds ample, 2-3 feet long, ovate or subdeltoid acuminate firm-coriaceous, 3-4-pinnate; primary pinnæ petiolate (as are the secondary and tertiary ones), 6-12 inches long, distant from a broad base, lanceolate, much acuminate; secondary ones also distant, of the same shape, but not acuminate, 1-1½ inches long, numerous; ultimate pinnules ovate-lanceolate, ¼ inch long, superior basal one the largest, acute, all pinnatifid with acute lobes, auricled at the superior base, terminal ones subconfluent; sori 2-5 on each pinnule, broad-oblong, or in age almost globose, very prominent and copious; involucre brown, membranaceous, broad, generally crescent-shaped or subreniform, fimbriated, extremely convex, so forced back by sori; rachises all pale brown, glossy, more or less flexuose. *Wall. Cat.* 339. *Hook. Sp. Fil.* iii. 234. *Syn. Fil.* p. 229. *Bedd. F. B. I. t.* 295.

Himalayas, from East Kashmir to Bhotan, 5,000-12,000 feet.

VAR. β *FOLIOSA.* (*Wall.*) Rachis often very red and wavy, glandular in the axis of the pinnæ, ultimate segments broader than in the type; sori small. *Wall. Cat.* 359. *Clarke, F. N. I. t.* 6 fig. 2.

Sikkim and Nepal at high levels.



C. 11. 11. 11.

ATHYRIUM FIMBRIATUM. (Hall.)

VAR. γ . SPHÆROPTEROIDES. (*Clarke*.) Involucre small, early disappearing; sori both lateral and terminal on the veins, becoming soon globose, often appearing elevated from the frond. *Clarke, F. N. I. t. 62, fig. 1.* *Asplenium Atkinsoni*, var. *Andersoni*, *Clarke, F. N. I. t. 57.*

Throughout the Himalayas, 9,000–13,000 feet.

GENUS XLII.—DIPLAZIUM. (*Swartz*.)

(*Diplazo* to be double; the double indusia or involucre.)

As in *Asplenium*, but some of the sori double, *i.e.*, on each side of the vein, each furnished with a linear indusium, the one opening interiorly, the other exteriorly.

1. DIPLAZIUM LANCEUM. (*Thunb.*) Caudex long-repent, rooting sparingly, scaly, stipes distant 4 inches to a span long slender, thickened at the base, and paleaceous with black scales, fronds chartaceous, firm, opaque, longer than the stipes, rarely an inch wide, lanceolate, attenuated, and acuminate at each extremity entire, sometimes a little repand; costa slender, prominent beneath; veins horizontal, fascicled, the superior, and sometimes the inferior branch only fertile; sori linear, distant, remote from the costa, often diplazioid. *Thunb. Fl. Jap. 333.* *Hook. Sp. Fil. iii. 235.* *Bedd. F. S. I. t. 227.*

Ceylon, Matale East and Saffragram 2,000–3,000 feet, dies in dry weather. Assam and N. E. Bengal 500–5,000 feet, extending to Chittagong and East Nepal.

(Also in China, Formosa, and Japan.)

2. DIPLAZIUM SUBSERRATUM. (*Bl.*) Caudex creeping, elongated; fronds membranaceous, $1\frac{1}{2}$ feet long, $1-1\frac{1}{2}$ inch wide, glabrous and satiny, elongate-oblong, lanceolate, sharply acuminate, moderately attenuated below, obtuse or acute at the base, towards the middle sinuate-crenate, coarsely dentate-serrated nearer the apex; costa slender, firm, prominent beneath; veins fascicled,


horizontal or nearly so, one to three of the exterior branches soriferous; sori distant, often curved; involucres often diplazioid, narrow-linear, occupying about two-thirds of the space between the costa and margin; stipes 1-2 inches long, slender, terete. *Bl. En. p.* 174. *Hook. Sp. Fil.* iii. 236. *Bedd. F. B. I. t.* 289.

Penang.

(Also in Java.)

3. *DIPLAZIUM ZEYLANICUM*. (*Hook.*) Caudex terete, repent, subterraneous, naked, black; stipes remote, solitary, 4 inches to a span long, paleaceous with lax dark subulate-lanceolate scales; fronds herbageous, a span to a foot long, 1-2 inches broad, lanceolate acuminate, deeply pinnatifid in the middle, pinnate at the base, serrated only towards the apex, lobes and pinnae horizontal, oblong, obtuse; veins pinnated, entire or forked; sori linear; involucres with the superior basal one principally diplazioid. *Hook. Sp. Fil.* iii. 237. *Bedd. F. S. I. t.* 228.

Ceylon, forests of Ambagamwa and Kotmallee.



DIPLAZIUM LANCEUM (*Thunb.*)

4. *DIPLAZIUM PALLIDUM*. (*Bl.*) Stipes a foot or more long, stramineous, deciduously scaly at the base; fronds 1½-2 feet long, ovate acuminate, subcoriaceous, often pale whitish-green when dry pinnated, pinnae numerous, approximate, mostly petiolate, horizontal, 5-6 inches long, ¼-½ of an inch broad, from a somewhat rounded

or obliquely cuneated base, linear-oblong, acuminate, falcate, sharp'y and subspinulose serrated, terminal one petiolate and similar to the rest, or larger and broad and pinnatifid at the base from being formed of dwarfed and confluent pinnae, very coarsely serrated towards the finely acuminate apex; veins very patent, immersed, generally twice dichotomous, the lowest superior branch (arising from near the costa) bearing the linear sorus along its whole length, rarely diplazioid, and only from 1-3 at the superior and mostly truncated

but not auricled base; involucre narrow, firm, membranaceous. *Bl. En. Fil. Jav.* 176. *Hook. Sp. Fil.* iii. 238. *Bedd. F. B. I. t.* 196.

Birma and the Malay Peninsula.

(Also in the Malay Islands and the Philippines.)



N°85

DIPLAZIUM PORRECTUM. (Wall.)

5. *DIPLAZIUM PORRECTUM*. (Wall.) Caudex erect or ascending; stipes tufted, 6-12 inches long; fronds 8-10 inches to 1½ foot long deltoid-oblong, pinnate; pinnae horizontal, numerous, 4-6 inches long, petiolate or sessile, and confluent at the acuminate pinnatifid apex, from a broad truncated and auricled or generally quite hastate base (having a distinct auricle above

and below), linear-lanceolate, obtuse or acuminate, lower ones more or less deeply pinnatifid, intermediate ones obtusely dentate, uppermost ones entire at the margins; veins patent, pinnated in the auricles, the rest once or twice forked; sori linear, elongated, confined to the superior or basal veinlet, and extending its whole length, or on the two outer ones, mostly diplazioid; involucre membranaceous when young. *Wall. Cat.* 204. *Hook. Sp. Fil.* iii. 250. *Bedd. F. B. I. t.* 245.

The Malay Peninsula, Penang, &c.

(Also in the Malay Islands.)

6. *DIPLAZIUM BANTAMENSE*. (Bl.) Rhizome creeping; stipes 6-12 inches long, with lanceolate caducous scales; fronds up to 1½ foot long, simply pinnate; pinnae generally 3-5 on each side, alternate or subopposite, with one terminal one, subcoriaceous, glabrous, generally sharply serrated towards the apex, otherwise nearly entire, 6-9 inches long, by 1-2 broad, somewhat narrowed at the base; apex acuminate or often caudate; veins pinnate; sori slender, irregular, starting from near the midrib and extending nearly to the margin, or only half-way towards it. *Bl. En. Fl. Jav. Fil.* 190-191. *Hook. Syn. Fil. p.* 231. *D. fraxinifolium* (Wall.), *Bedd. F. B. I. t.* 59.

Madras Presidency, on the Tinnevely and Travancore Hills. Khasya Hills, 4,000 feet elevation; Cachar 500 feet elevation; Malay Peninsula.

(Also in Malay Islands and Southern China.)



Nº 86.

DIPLAZIUM BANTAMENSE. (Bl.)

7. *DIPLAZIUM SYLVATICUM*.

(Presl.) Caudex decumbent;

stipes 1 foot long, firm, erect,

naked except at the base, where furnished with scales; fronds 1-3 feet long, simply pinnate, ovate-lanceolate in outline; pinnae numerous, about 6 inches long by 1 inch broad, the apex acuminate or caudate, the margin nearly entire or slightly waved or serrated, particularly towards the apex, but not to more than one line deep (at least in the type); base truncate or narrowed; texture thin, herbaceous; veins pinnate, fine; sori long, linear. *Presl. Rel. Haenk.* 1-42, under *Asplenium*. *Hook. Syn. Fil.* 232. *Bedd. F. S. I.* 161, a small specimen.

Madras Presidency, throughout the Western forests up 4,500 feet. Ceylon, central provinces; Malay Peninsula.
(Also in Mauritius, Fernando Po, Java, and Borneo.)

VAR. β PRESCOTTIANUM. (*Wall.*) Pinnæ narrower than type, more deeply serrated or cut, and more or less lobed toward the base. *Wall. Cat.* 235. *Hook. Sp. Fil.* iii. 251. *Bedd. F. B.* t. 243.

Singapore and Penang (perhaps a distinct species).

8. DIPLAZIUM SPECIOSUM. (*Mett.*) Stipes tufted, 1 foot more long, firm, erect, straw-coloured or brownish, nearly naked; fronds 1-2 feet long, 8-12 inches broad, with 10-20 pairs of pinnae below the pinnatifid apex, the lower ones often stalked, 4-6 inches long, about $\frac{1}{4}$ inch broad, the apex much acuminate, the edge lobed, the lobes reaching down half to two-thirds of the way to the rachis; generally falcate acute, obscurely crenate; texture herbaceous but firm; rachis erect, naked; veins pinnate in the lobes; sori slender, reaching nearly to the edge; rachis 4-sided with sharp angles. *acuminatum*, *Wall. Mett. (non H. and A.). Hook. Syn. Fil.* p. 231. *Bedd. F. B. I. t.* 290.

The Malay Peninsula; Tinnevely Mountains. Mr. Clarke considers it a form of *sylvaticum*.

9. DIPLAZIUM PINNATIFIDO-PINNATUM. (*Hook.*) Stipes 6-10 inches long, firm, erect, naked; fronds 9-12 inches long, 6-9 inches broad, pinnate, with a pinnatifid apex; pinnae 3-4 only on each side, 3-4 inches long, 1 inch broad, the apex acute, the margin sharply serrated; base cuneate; texture coriaceous, drying a dull blackish colour; veins obscure, pinnate; sori irregular, beginning at midrib, not reaching the margin. *Hook. Syn. Fil.* 231. *Bedd. F. B.* t. 244.

Mishmee; Mr. Clarke has united it with the Philippine *Lobelia*, of which there is only one authentic specimen at Kew, although this does not represent the whole frond, it has numerous pinnae of a lighter colour and more prominent venation; it appears to me to be quite a different plant.

10. *DIPLAZIUM LONGIFOLIUM*. (*Don.*) Caudex erect, small, densely clothed with the bases of old stipes, paleaceous, stipes tufted 5-6 inches long, fronds broad-lanceolate, acuminate, membranaceous, a span to 14 inches long, pinnated, pinnatifid at the apex, pinnæ, on slender petioles, $1\frac{1}{2}$ 3 inches long, very patent, broad-lanceolate, falcate, sharply acuminate, truncated and auricled at the superior base, inferior base excised, the margin more or less deeply lobed, lobes as well as the auricles spinulosely serrate; costa slender, flexuose, vein-like; veins in oblique fascicles corresponding to each lobe, dichotomous lowest superior branch only soriferous; sori mostly asplenoid, linear, neither attaining the costa nor the margin. *Don. Prod. Fl. Nep.* 7, under *Asplenium*. *Diplazium lobulosum*, *Bedd. F. B. I. t.* 247. *Hook. Sp. Fil. t.* 141. *Syn. Fil.* 234.

Mr. Clarke has placed this in *Asplenium*, but there are diplazioid indusia on my specimens, and on some of those at Kew. *Asplenium trapeziforme* (Roxb.), *Bedd. F. S. I. t.* 134, is not at all allied here, as suggested by Mr. Clarke.

Nepal and Kumaon, 6,000-8,000 feet.



DIPLAZIUM LONGIFOLIUM. (*Don.*)

11. *DIPLAZIUM TOMENTOSUM*. (*Hook.*) Caudex 1-2 inches, erect, sending down wiry black roots; stipes 6-9 inches long, firm, erect; fronds deltoid-lanceolate, 6-12 inches long, 3-4 inches broad at base and gradually narrowing upwards; pinnæ numerous, the lowest pair deflexed, the others horizontal, $\frac{3}{8}$ - $\frac{5}{8}$ inch broad, narrow-oblong, somewhat falcate, usually auricled at base on the upper side, pinnatifid half or two-thirds down to the midrib in the larger fronds, subentire in the smaller ones; segments falcate, $\frac{1}{2}$ inch broad;

(Also in the Malay Islands.)

12. DIPLAZIUM JAPONICUM. (*Thunb*
suberect ; stipes up to 18 inches long, pub
herbaceous, 8–18 inches long, 6–8 inches
late pinnate ; rachises and costa more or
hairs mixed with small scales, or almost qu
on each side, below the pinnatifid apex, ;
the lower ones quite opposite, some of the
ones sessile or decurrent, all pinnatifid eitl
or quite down to a winged rachis ; segme
toothed rounded or falcate apex, or rather
pinnate in the lobes ; veinlets simple or
hairy or with minute crisped scales ; sori
the midrib and not quite reaching the marg
Diplazium Thwaitesii (A. Br.), *Hook. Syn.*
291. *D. lasiopteris* (Mett.), *Hook. Syn. t.*
t. 160. *Diplazium decussatum* (Wall.),
D. polyrhizon, *Baker, Syn. Fil.* 490.

I have followed Mr. Clarke in redu
japonicum ; the typical *lasiopteris* of South
more hairy and generally lanceolate in shap

the rhizome creeping, though it is generally erect; in *lasiopteris* and *Thwaitesii* it is wide creeping. I have seen nothing like Mr. Clarke's variety *chattagramica*, with bipinnate fronds, the secondary pinnæ petioled and the rachis not winged, and can hardly fancy it belongs here, the specimen could not be found at Kew, but from the figure in Clarke's Review, it appears to be one of the numerous forms of *latifolium* and perhaps the same as the one he calls *succulentum*. Mr. Clarke is quite wrong in referring *Schkuhrii* Thwaites here, he did so on a specimen of my *decurrens*, wrongly named *Schkuhrii*, but *decurrens* (which is Thwaites's *polypodioides* var. β c. p. 3332, and Moore's *dilatatum*, var. β minor), can have no place here.

Madras Presidency, on the Western ghats, the supposed *lasiopteris* at Ootacamund, 7,000 feet, and Poombary on the Pulneys, 7,000 feet; *decussatum* on the Tinnevely Hills and Jeypore Hills, West of Vizagapatam, 3,000–4,000 feet. Ceylon, central provinces; N. W. Himalayas; Nepal; Nynee Tal; Kangra; Kumaon; Khasya, 3,000 feet; Malay Peninsula.

(Also in Japan, China, and Polynesia.)

13. *DIPLAZIUM SCHKUHRII*. (*Thwaites*.) Caudex erect; scales dense, small, lanceolate, dark-brown; stipes 9–18 inches long, naked; fronds deltoid-lanceolate, up to 2 feet long, bipinnate below, bipinnatifid above; pinnæ alternate, lower ones distant, 6–8 inches long, by $1\frac{1}{2}$ –2 inches broad; rachis winged; pinnules distant, sessile ligulate-oblong, all very regular in shape, $\frac{3}{4}$ – $\frac{7}{8}$ inch long, $\frac{3}{8}$ – $\frac{1}{4}$ inch broad, pinnatifid about $\frac{1}{3}$ down into rounded lobes; texture rather firm; surfaces glabrous, bright-green; veinlets simple or pinnated in the lobes; sori linear, generally reaching from the midrib to the sinus between the lobes, double or single. In younger plants or smaller specimens, the fronds are often pinnate only, *i.e.*, the primary pinnæ only pinnatifid half to two-thirds to the rachis. *Thw. En. Pl. Zey. p. 385.* (*Not Mett. or J. Sm. which is only a form of sylvaticum.*) *Hook. Syn. Fil. p. 491.* *Bedd. F. S. I. t. 230.* Ceylon; central provinces.

14. *DIPLAZIUM SORZOGONENSE*. (*Presl*.) Stipes tufted, scaly


below ; fronds up to 16 inches long, pinnate, with numerous pinnæ; pinnæ sessile, about 6 inches long, by $1\frac{1}{4}$ broad, cut down very regularly throughout two-thirds to the rachis into oblong blunt subentire lobes, which are of equal breadth, ($\frac{1}{4}$ inch) throughout to the apex; texture firm ; rachis slightly scaly ; veinlets simple, each occupied in its entire length from costule to apex by a sorus. *Hook. Syn. Fil. p. 236 (in part.) Bedd. F. B. I. t. 246.*

Malacca and Penang ; differs from the North India plant in its longer lobes with quite parallel edges, its simple veins, and its longer sori. The figure quoted above is from the Malay plant (though supposed at the time to be from North India), and there is another specimen of the same in the Kew Herbarium, labelled Khasya, Griffith, but it is probable that Griffith obtained it from Malacca.

15. DIPLAZIUM STOLICZKÆ. (*Bedd.*) Stipe and rachis glabrous or subglabrous ; fronds 1– $1\frac{1}{2}$ feet long, by 10 inches broad, pinnate ; pinnæ numerous, the lowest pair slightly the largest, and only a few of the upper ones gradually diminishing, the lower 2–3 pairs opposite or subopposite, the rest alternate ; texture subcoriaceous ; pinnæ very shortly petioled, 5–6 inches long, $1-1\frac{1}{2}$ inch broad, gradually tapering towards the apex, cut down $\frac{2}{3}$ to $\frac{5}{8}$ of the way to the rachis, into oblong, rounded pinnules, which are more than $\frac{1}{4}$ inch broad, and very regularly crenated ; main veins slightly wavy ; veinlets simple or forked towards their apex or below their centre, all very conspicuous ; sori 3–7 to each pinnule or segment (on the lower veins only), extending from the main vein $\frac{2}{3}$ of the way to the margin. *Bedd. F. B. I. p. 13 and t. 361.*

North India (Dr. Jerdon), probably Khasya. This is not in the Kew Herbarium, and had not been seen by Mr. Clarke when he wrote his Review.

VAR. β HIRSUTIPES. Differs from the type by the stipes and often the rachis being very fibrillose, in the pinnæ being narrowed, and the lower ones generally more or less reduced, and in the pinnules being much less incised. *Diplazium sorzogonense, Hook. Syn.*



A. J. A. S. P. C. O. D. R. C.

DIPLAZIUM ASPERUM. (Bl.)

Fil. 236 (*in part*). *Diplazium Stoliczkæ*, *Clarke, F. N. I. p. 5* 
Diplazium thelypteroides, *Bedd. F. B. I. t. 68*.

Himalayas ; Nepal to Bhotan, 7,000–10,000 feet ; very abundant ~~about~~
 about Darjeeling. This and the type may run into each other, but as
 yet only one specimen of the type is known ; they both look very
 different to the Malay *sorzogonense*, and differ by the lobes being
 more serrated, the veinlets being often forked, the sori never reach-
 ing the margin and being confined to the 3–4 lower veinlets.


16. *DIPLAZIUM ASPERUM*. (*Bl.*) Caudex erect, often quite a
 large trunk, (like *Alsophila*) stipes main and partial rachises prickly
 and scaly ; fronds large, rather coriaceous, bipinnate with the pin-
 nules pinnatifid nearly to the rachis, pinnules quite at right angles
 with the rachis, sessile or subsessile, 3–5 inches long, about 1 inch
 broad, segments very regular, forming an oblong parallelogram, or
 slightly falcate, obtuse, crenate, or serrate, often rather sharply ;
 veinlets numerous, 8–12 in each segment, generally simple, rarely
 forked, all or nearly all generally soriferous ; sori touching the costa
 but not the margin, generally only one of the lower ones double.
Bl. En. 195. *Diplaz. polypodioides*, *Hook. Syn. Fil.* 238 (*in part*).
Diplaz. polypodioides, *Bedd. F. S. I. t. 163*. *Diplaz. sikkimense*,
Clarke, F. N. I. p. 500, t. 65, fig. 1, scarcely differs, and is certainly
 nearer to this than *polypodioides*.

Madras Presidency ; throughout the Western forests, from no
 elevation up to 3,000 feet, often quite a tree fern ; Cuddapa forests ;
 Sikkim, banks of the Teesta, 500 feet.

(Also in Java.)

17. *DIPLAZIUM POLYPODIOIDES*. (*Mett.*) Caudex erect, stout,
 densely clothed at the crown with long brown fibrillose scales, 1 inch
 long ; stipes densely tufted, stout, green, 1 foot or more long ;
 fronds 3–4 feet long, and 2 feet wide, bipinnate, with 8–9 primary
 pinnæ on each side below the simple ones, the lower ones generally
 distant, secondary pinnæ 2–4 inches long, cut down nearly to the
 rachis in the typical form, but sometimes only $\frac{1}{3}$ down ; segments
 short oblong, crenate or serrate, or subentire ; veinlets about six
 generally forked, sometimes simple ; texture herbaceous ; surface

N°89



TRYPAS

DIPLAZIUM POLYPODIOIDES. (Mett.)

glabrous or nearly so ; sori in two oblique rows in the segments ; indusium linear, commencing at the midrib, but generally falling well short of the margin, lower ones generally double. *Mett. Hort. Fil. Lips.* 78. *Hook. Syn. Fil.* 238 (excluding *asperum* (Bl.), which differs in its more coriaceous texture, much more regular segments, numerous veinlets, and its prickles.) *Bedd. F. B. I. t.* 293. *Asplenium marginatum*, *Wall. Cat.* 391, *type sheet*.

Madras Presidency, in the Western forests up to 6,000 feet elevation. North India, throughout the Himalayas and Khasya hills ; Ceylon ; the Malay Peninsula. Clarke's variety "*vestita*" has the rachises more or less villous or subtomentose, but does not otherwise recede from the type, he says it is confined to the central Himalayas. His variety "*sublatifolia*" runs into "*latifolia*," and rather belongs to that species, if the two are really distinct, which is very doubtful, his variety "*effusior*" is *D. umbrosum*, var. *multicaudata*.

(Also in Australia.)

VAR. β DECURRENS. (*Bedd.*) Pinnæ more deltoid in form with the secondary pinnæ few and distant, and more or less decurrent on the rachis, only cut down a third or half-way to the rachis ; veinlets in the segments few, only 3-4 ; sori curved, generally occupying the whole length of the veinlets, and extending to the margin. *Diplazium decurrens*, *Bedd. F. S. I. t.* 229. *Diplaz. polypodioides*, var. β , *Thw. En. Pl. Zey. p.* 385. *c. p.* 3332. *Diplaz. dilatatum*, var. β minor, *Moore, Index Fil.* 327, *in part only*.

A very distinct looking fern, and I believe a good species, but I now prefer to follow Mr. Thwaites (who alone can have seen it growing), and consider it a variety of *polypodioides*. Sir W. Hooker, *Sp. Fil. t.* 258, has mentioned it under *polypodioides* as perhaps distinct species. Mr. Baker has included it under *maximum* (Don) which is otherwise, as far as the Kew bundle is concerned, one the large forms of *latifolium*, and Mr. Clarke in his Review named it *Dipl. Schkuhrii*, but this was owing to his having received a specimen of it so labelled (evidently by mistake), from Mr. Thwaites.

Ceylon ; Ambagamwa.

18. *DIPLAZIUM LATIFOLIUM*. (*Don.*) Like *polypodioides*, only the secondary pinnæ are generally much less cut down, and often much broader, they are sometimes almost entire, or with only shallow serratures, when the fern has quite the appearance of “*sylvaticum*,” only bipinnate instead of simply pinnate; other forms have the secondary pinnæ 2 inches (or even more) broad at the base, cut down a third or half-way to the rachis; segments always more or less crenate or serrate; veinlets simple or forked, rather distant, their number depending on the size of the segments (never so numerous as in *asperum*, except when the segments are double the size of those of that plant); sori narrow, often occupying the whole length of the veinlet, and reaching the margin; indusium sometimes obsolete or early caducous. *Don. Prod. Fl. Nep.* 8. *Hook. Syn. Fil.* 239. *Dip. dilatatum*, *Hook. Sp. Fil.* iii. 258. *Bedd. F. S. I.* 162, a form running nearer to *polypodioides*. *Dip. maximum*, *Hook. Syn. Fil.* 232, in part.

Madras Presidency, in all the Western forests. North India, throughout the Himalayas and Khasya Hills. Ceylon; Malay Peninsula; from very low altitudes up to 9,000 feet elevation.

(Also in Australia, China and the Philippines.)

If we only included here species with very broad secondary pinnæ, not cut more than half-way down to the rachis, this plant would be very distinct from “*polypodioides*,” but I fear this is not possible, and in all large herbaria, I find specimens that it is very difficult to say which species they should be referred to. *Diplazium torrentium* and *succulentum* of Clarke cannot be made into distinct species, unless we also make many more, but they belong to types which, though generally referred here, must, from the deeper cutting of their secondary pinnæ, go into “*polypodioides*,” if the two are to be kept distinct; no figures and no description could enable any one to distinguish some of these forms as species, the only difference often being the extent of the cutting of the secondary pinnæ. *D. decurrens* unless a distinct species (which view its venation I think supports), should rather be referred here than

19. *DIPLAZIUM TRAVANCORICUM*. (J. Sm.)
with trunk-like caudex, secondary pinnæ lanceolate, somewhat attenuated at the base, caudate, quite pinnate towards the base, the petioled, the upper ones gradually becoming so, the pinnæ terminating with a long broad pinna, the upper 8 inches or so is only very shallow pinnules smaller than the adjoining ones, next long, by about 1 inch broad, lanceolate in shape towards the apex ; veins numerous, prominent central costa ; texture subcoriaceous and striated ; sori commencing a little distance not nearly reaching the margin.

Travancore Hills ; Athraymally forests, a

20. *DIPLAZIUM UMBROSUM*. (J. Sm.)
Stipes 1 foot or more long, strong, erect, clothed with scales, and sometimes muricate ; fronds 3-5 feet long, 1-2 feet broad ; primary pinnæ ovate-lanceolate to deltoid, 1-2 feet long ; secondary pinnæ very various, some pinnatifid (in the simpler forms), to quite pin-

into the other even in their restricted areas, and that others are geographical varieties which would not be constant in cultivation, and secondly, because I think any attempt to separate them into species would only add to the present confusion, it being a task that should only be undertaken after a cultivation of all the different forms.

Mr. Clarke has proposed the subgenus *Pseud.-Allantodia* for the species with allantodioid sori and free veins, but I am afraid these allantodioid sori are not constant or to be depended on, *Australe*, from the Nilgiris, has the sori often quite allantodioid as figured by me, *tab.* 158, *F. S. I.*; but I have specimens where all the sori are perfectly diplazioid; again, I have specimens of *Asplenium resectum*, from two different localities, with nearly all the sori allantodioid.



DIPLAZIUM UMBROSUM, VAR. AUSTRALE. (*R. Br.*)

VAR. AUSTRALE. (*R. Br.*) Stipes often muricate; fronds 3-pinnatifid, often rather flaccid in texture, ultimate segments broad and blunt, sori sometimes quite allantodioid, at other times quite asplenioid or diplazioid; rhizome creeping or erect. *Athyrium australe*, *Bedd. F. S. I. t.* 158. *Diplaz. bellum*, *Clarke, F. N. I. p.* 496. *Diplaz. multicaudatum*, var. *tristis*, *Clarke, l. c. p.* 512.

Sikkim, Bhotan; Khasya; Nilgiris and other mountain forests, 5,000-7,000 feet, in South India; Birma.

(Also in Australia and Tropical Africa.)

VAR. PROCERUM. (*Wall.*) Rhizome creeping; stipes muricate; fronds very large, moderately firm in texture, primary pinnae up to nearly 2 feet long, secondary pinnae 5-5½ inches, again pinnated with the pinnules about 1 inch long, and pinnatifid about half-way to the rachis; segments oblong, crenate; veinlets in the segments once forked; sori mostly in two rows, near and parallel to the midrib, but other very short sori are scattered on the segments; indusium allantodioid. *Wall. Cat.* 2203. *Hook. Syn. Fil.* 489. *Dipl. procerum*, *Clarke, 495.*

perpetuated 3 inches long and 1 in. in breadth
 nearly to the rachis, the segments oblique
 long, rather narrow ligulate, more or less s



No 91

ADIANTUM UMBROSUM,
VAR. MULTICAUDATUM. (Wall.)
 segments rounded entire, to slightly crenate
 to simple or forked ..

forked

or st

chis;

Wall

Bedd.

Griffith

Bedd.

F.

to Bh

comme

5,000 f

2,000 f

V.

Fronds

ceous,)

the ter

very su

slightly crenate

GENUS XLIII.—ANISOGONIUM. (*Presl.*)

(From *anisos*, unequal; *gonia*, an angle; the angles of the venation.)

As in *Diplazium*, but the veins anastomosing.

1. ANISOGONIUM CORDIFOLIUM. (*Mett.*) Stipes 6–12 inches long, firm, erect, scaly below; fronds 8–12 inches long, 3–4 inches broad, entire, caudate at the base, the point acuminate or subdeltoid, with a large terminal and 1 or 2 pairs of smaller but similar spreading lateral pinnae; texture coriaceous; veins in close groups of about 4, anastomosing copiously in the outer half of the frond; sori reaching from the midrib to the edge.

Mett. Fil. Hort. Lips. p. 74, t. 12. f. 6.

Hook. Syn. Fil. 243. Bedd. F. B. I.

t. 331 (the simple form.)

Malacca and Malay Peninsula.

(Also in Philippines and Malay Islands.)


2. ANISOGONIUM LINEOLATUM. (*Mett.*) Stipe 6–9 inches long, firm, erect, grey, scaly below; fronds occasionally simple, usually with a terminal pinna and 3–6 pairs of lateral ones, which are 6–12 inches long, 2 inches or more broad, entire, often suddenly acuminate; texture coriaceous; rachis naked; veins 4–8 to a cluster, uniting slightly towards the edge.

Mett. l. c. p. 74. tab. 11. f. 5. Hook. Syn. Fil. 243. Bedd. F. B. I. t. 330.

Malacca and Malay Peninsula.

(Also in Philippines and Malay Islands.)


3. ANISOGONIUM HETEROPHLEBIUM. (*Mett. MSS.*) Stipes 1 foot long, grey, scaly throughout; fronds 12–18 inches long, 8–9 inches broad, with 6–8 opposite pairs of pinnae below the pinnatifid apex; the lowest 2 inches or more apart, 3–4 inches long, 1–1½ inches broad, the edge undulated, the point acute, the base cordate



ANISOGONIUM CORDIFOLIUM. (*Mett.*)


East Nepal to Malacca, 1000-6000

4. *ANISOGONIUM ESCULENTUM*. (Presl)



ANISOGONIUM HETEROPHLEBIUM. (Mett.)

midrib, with lines of sori often on all the lateral veins. *Hook. Syn. Fil.* 241. *Bedd. F.*




ANISOGONIUM ESCULENTUM. (Presl.)

both surfaces and rachis naked, the lower part, veins in groups of 2-3 on a side half-way across to the edge; sori often $\frac{1}{2}$ p. 245. *Bedd. F. B. I.* 332 (*Callipteris*)
Ceylon, Matale East and Oodawella.

GENUS XLIV.—HEMIDIC

(*Hemi*, half; *diktyon*, net; outer portion)

As in *Asplenium*, but veins anastomosing to the margin of the frond.



(*Linn.*
1-3 in
chaffy,
one in
pinnate
lobes
inch long
a rounded
texture
surface

(Also in Afghanistan, Western Asia, Europe, and North-west Africa.)

2. *HEMIDICTYUM FINLAYSONIANUM*. (*Hook.*) Stipes 6–9 inches long, green, subcompressed, nearly naked; fronds usually 1 foot long (often much more), 6–8 inches broad, occasionally simple and entire, generally pinnate with 2–6 opposite pairs of lateral pinnæ, the lower ones 4–8 inches long, $1\frac{1}{2}$ –2 inches broad, the apex very acuminate, the base narrowed into a distinct petiole, the margin entire or irregularly lobed; texture leathery; veins subflabellate, very oblique, anastomosing slightly towards the margin, sometimes bounded by an irregular intramarginal line; sori often $1\frac{1}{2}$ –2 inches long, rooting buds are often produced from the midribs of the pinnæ near the apex. *Hook. Syn. Fil.* 245. *Bedd. F. B. I. t.* 72.

North and East Bengal, with Assam, up to 3,000 feet, very common, especially in Khasya; Malay Peninsula.



N° 96

HEMIDICTYUM FINLAYSONIANUM.
(*Hook.*)

GENUS XLV.—*ALLANTODIA*. (*Wall.*)

(*Allantos*, a sausage; form of indusium.)

Sori dorsal, linear-oblong, attached to the primary veins; indusium the same shape as the sorus and quite enclosing it, bursting in an irregular line down to the centre, or rarely asplenoid; veins anastomosing and forming several elongated areoles, smaller towards the margin, with free marginal clavate veinlets; fronds pinnate, very membranaceous, (the character of the indusium is not to be relied on, similar sausage-shaped indusia being found in species of true *Asplenium* and *Diplazium*.)

1. *ALLANTODIA JAVANICA*. (*Bl. under Asplenium.*) Fronds



often 2 feet long, 1 foot broad, pinnæ 4–8 inches long, 1–1½ inch broad, oblong, entire or slightly crenulate, particularly at the caudate apex, veins forked near the midrib, 2–3 of hexagonal areoles occupying the outer half of the space between midrib and margin; sori confined to the anterior vein of the first fork or areole. *Bl. En. Pl. Jav. Fil.* 175. Allant. Brunoniana (Wall.), *Hook. Syn. Fil. p.* 246. *Bedd. F. S. I.* 159.

Nepal and Bhotan, 4,000–7,000 feet elevation; Khasya, Mikir Hills; Ceylon.

(Also in Java and Samoa.)

GENUS XLVI.—ACTINIOPTERIS. (*Link.*)

(*Actin*, rays; *pteris*, a fern.)

Sori linear, elongated, submarginal; indusium the same shape as the sorus, folded over it, placed one on each side of the narrow segments of the frond opening towards the midrib: a single species like a miniature palm.

1. ACTINIOPTERIS DICHOTOMA. (*Forsk*, under *Acrostichum*.) Stipes densely tufted, 2–6 inches long; fronds like fans, 1–1½ inch deep, composed of numerous dichotomous segments which are rush-like in texture, not more than ½ line broad, the veins few and subparallel with the indistinct midrib, the segments of the fertile frond longer than those of the barren one. *Acrostichum dichotomum*, *Forsk. Fl. Ægypt. Arab.* 184. *Actiniopteris radiata*, *Hook. Syn. Fil. p.* 246. *Bedd. F. S. I. t.* 124.

Throughout India, especially the Peninsula in dry rocky places below 3,000 feet elevation; Ceylon.

(Also in North Africa, Mascareen Islands, Persia, Cabul.)



TRIBE IX.—ASPIDIEÆ.

Sori dorsal, subglobose, rarely elliptical; indusium superior, similar in shape to the sorus, fixed either by the centre or a sinus.

GENUS XLVII.—DIDYMOCHLÆNA. (*Desv.*)

(*Didymos*, double; *chlæna*, a cloak;—the double indusium.)

Sori elliptical, terminal on a veinlet, but distinctly intramarginal; indusium elliptical, marginate at the base, attached longitudinally to the linear receptacle, free all round the edge; veins subflabellate, free, their apices clavate; fronds bipinnate, pinnules articulate with the rachis.

1. DIDYMOCHLÆNA LUNULATA. (*Desv.*) Caudex erect, subarborescent; fronds densely tufted, 4–6 feet long, bipinnate; pinnules $\frac{3}{4}$ –1 inch broad, dimidiate, subquadrangular, entire or slightly sinuated; texture subcoriaceous; veinlets forked or pinnate, their apices marked with white dots; sori 2–6 to a pinnule. *Desv. in Mem. Soc. Linn.* vi. p. 282. *Hook. Syn. Fil.* p. 248. *Bedd. F. B. I.* t. 15.

Birma and the Malay Peninsula.


(Also in Tropical America; East African Islands; Malay Islands; Fiji.)

GENUS XLVIII.—MESOCHLÆNA. (*R. Br.*)

(*Mesos*, middle; *chlæna*, cloak, attachment of the indusium.)

Sori and indusium of Didymochlæna, but habit and venation of Nephrodium, i.e., the lower pair of veinlets anastomose, the others free, parallel; fronds bipinnatifid.

1. MESOCHLÆNA POLYCARPA. (*Bl. under Aspidium.*) Caudex erect; stipes tufted, short; fronds 2–3 feet long, 12–18 inches broad, pinnate; pinnæ close, very numerous, spreading, dwindling down below to mere auricles, the largest 6–9 inches long, $\frac{1}{2}$ inch broad, cut down half-way or more to the rachis into linear-oblong lobes; rachis and lower surface villose; veins very close, pinnated,



twelve or more on each side in a lobe ; sori small, close, ultimately confluent. *Aspidium polycarpum*, *Bl. En. Fil. Jav.* 156. *Didymochlæna* *Hook. Syn. Fil. p.* 248. *Mesochlæna polycarpa*, *Bedd. F. B. I. t.* 344. *Nephrodium javanicum*, *Hook. Sp. Fil. iv. p.* 67.

In the *Syn. Filicum* this genus has been joined with *Didymochlæna* ; but, as Mr. Smith has pointed out, it is really a *Nephrodium* with the indusium elongated.

Malay Peninsula.

(Also in the Malay Islands.)

GENUS XLIX.—POLYSTICHUM. (*Roth.*)

(*Polys*, many ; *stichos*, order.)

Sori subglobose, dorsal or terminal on the veinlets ; indusium orbicular, fixed by the centre ; veins all free ; texture generally coriaceous and teeth generally awned.

1. *POLYSTICHUM SEMICORDATUM*. (*Sw.*) Stipes scattered, 6—12 inches long, fibrillose at the base ; fronds 2—3 feet long simply pinnate ; pinnae 4—6 inches long, $\frac{1}{4}$ inch broad, nearly entire, acuminate, cordate or truncate at the base ; texture herbaceous or somewhat firm ; surfaces naked and rachis nearly so ; veins pinnate, the lower ones ending short of the margin ; sori in 1—3 rows on each side, the inner one the most constant and regular, close to the midrib.

Sw. Syn. Fil. p. 45. *Hook. Syn.*


Fil. 249. *Bedd. F. B. I. t.* 35.

Birma and the Malay Peninsula.

(Also in the Malay Islands, Philippines, and Tropical America.)



POLYSTICHUM SEMICORDATUM. (*Sw.*)



2. *POLYSTICHUM LONCHITIS*. (*Linn. under Polypodium.*) Caudex short, stout oblique, densely paleaceous; stipes short paleaceous, lower portion of the rachis with ferruginous large scales; fronds 6–18 inches long, densely tufted, erect rigid, linear or lanceolate, tapering at both ends, pinnated; pinnæ numerous, approximate from a broad nearly sessile obliquely truncated base, ovate or lanceolate, falcate acute spinulose-serrate, the superior base truncated and auriculate; sori confined to upper portion of the frond in two or more series upon the pinnæ; indusium denticulate on the margin. *Hook. Sp. Fil.* iv. 8. *Sw. Syn. Fil.* p. 43. *Hook. Syn. Fil.* p. 250. *Bedd. F. B. I. t.* 128. *Polypodium Lonchitis*, *Linn. Sp. pl.* 1548.

North West Himalaya, Kashmir, Sonamurg 11,000 feet elevation. (Also in Arctic and Alpine Europe, Asia, and North America.)

3. *POLYSTICHUM LACHENENSE*. (*Hook.*) Rhizome very scaly; stipes very numerous, thick, densely tufted, scaly, 2–4 inches long; fronds narrow-linear, 4–8 inches long by $\frac{1}{2}$ – $\frac{3}{4}$ inch broad, pinnate, the lower pinnæ remote but hardly reduced in size; pinnæ 2–3 inches broad, deltoid-ovate, subcoriaceous, lobed more or less or subentire, the apex blunt, the margins spinosely serrated or crenate-serrate; sori 6–12 to a pinna, often covering the whole under side; indusium incised. *Hook. Syn. Fil.* p. 250. *Clarke, F. N. I.* p. 506. *Bedd. F. B. I. t.* 32.

Sikkim, 13,000–16,000 feet, Lachen, Tungu, Samding, 7,000–15,000 feet; Kashmir, Palgram, 13,000 feet.

4. *POLYSTICHUM ATKINSONI*. (*Bedd.*) Rhizome small, scaly; stipes 2–3 inches long, thin, stramineous, and with the rachis furnished with a few chaffy scales; fronds 1–3 inches long, $\frac{1}{2}$ inch broad, pinnate, the lower pinnæ not reduced; pinnæ ovate, very coriaceous, spinosely-serrated, the superior base often with an auricle; sori 1–3 to a pinna, generally on the upper half only. *Bedd. F. B. I. Suppl. t.* 362.

Sikkim, Yakla Pass 10,000 feet; Lachen 10,000 feet; Bhotan.

5. *POLYSTICHUM AURICULATUM*. (*Linn. under Polypodium.*) Stipes tufted, 4–6 inches long, scaly below, or throughout; fronds

indusium often quite
Hook. Syn. Fil. 251.

Throughout India and Ceylon.

(Also in Formosa.)

TYPE. Pinnæ submembranaceous
inches long, serrate; upper base square,
auricled, lower base much cut away; in-
vally altogether absent. *Hook. Sp. Fil.* i
t. 120. *Polypodium harpophyllum* (Ze.


Nilgiris, Pulneys and other Western
at the higher elevations.

VAR. β MARGINATUM. (*Wall.*)
shining, scaly below, ovate-oblong, closely
lobed; indusium conspicuous, sometimes
366. *Don. Prod. Fl. Nep.* 3. *Bedd. Su*
F. N. I. 507.

North India; Kunawur to Bhotan, 7

VAR. γ CÆSPITOSUM. (*Wall.*) F
margin scarcely spinulose or serrate, s
scarcely coriaceous; indusium conspicu
367. *P. cœspitosum*, *Bedd. F. B. I. t.* 3
Prod. Fl. Nep. 3. *Clarke, l. c.* 507.

Khasya, 3,000–4,500 feet elevation
v....



POLYSTICHUM AURICULATUM. (Linn.)

6. *POLYSTICHUM ILICIFOLIUM*. (*Don.*) Stipes densely tufted, 2-4 inches long, clothed with large scales throughout; fronds linear or lanceolate, not attenuated at the base, very coriaceous, pinnate; pinnae either small simple triangular or subrhomboidal, or much larger and again pinnate, naked on both sides and shining, angles spinulose-mucronate; rachis fibrillose; sori mostly in two rows, usually large; indusium peltate; veins immersed. *Don. Prod. Fl. Nep.* 3. *Hook. Syn. Fil.* 251. *Polystichum stimulans* (*Kze.*), *Bedd. F.B. I. t.* 31.

Himalayas, from Chumbra to Sikkim, 7,000-11,000 feet elevation.

The smaller forms with simple pinnae are very distinct, but the larger ones with the pinnae again pinnated, quite graduate into *P. aculeatum*, var. *rufobarbatum* till it becomes quite indistinguishable from forms of that fern, and is probably only a variety of it.



No. 103.

POLYSTICHUM ILICIFOLIUM. the upper with the basal lobe much enlarged; veins pinnate or forked in the segments; sori terminal on a veinlet; indusium peltate, bluntly crenated or subentire. *Hook. Syn. Fil. p.* 251, and *2nd Cent. Ferns, t.* 25, in part only. *Bedd. F. B. I. t.* 126.

Himalayas; from Balti to Sikkim, 7,000-13,000 feet elevation.

Very near the smaller forms of *Prescottianum*, but the stipe is fibrillose, not scaly. Hooker's plate is taken from specimens of both species, and it is very probable that they are only varieties of the same plant.

VAR. β . GRACILIS. Fronds smaller, (3–5 inches long with the stipe by $\frac{1}{4}$ – $\frac{1}{2}$ inch broad), margin of pinnæ crenate, with a sudden acumination (not bristly serrate), involucre reniform. *Clarke, F. N. I. t. 508. Lastrea gracilis* (Moore), *Bedd. F. B. I. t. 198. Lastrea sparsa*, var. *gracilis*, *Hook. Syn. Fil. 498.*

Sikkim; 11,500 feet elevation. A very doubtful species, which Mr. Clarke considers only a variety of Thomsoni, so I place it here pending further material. Mr. Clarke says that the involucre is polystichoid.

8. POLYSTICHUM ACULEATUM. (*Sw.*) Stipes tufted, 6–12 inches long, scaly and fibrillose below or throughout; fronds large, ovate-lanceolate, bipinnate, coriaceous, not attenuated at the base (or only slightly so in some forms); pinnæ numerous close lanceolate, pinnules oblong, unequal at the base, lower margin excised, upper generally more or less auricled, margins bristly or spinulose; texture more or less coriaceous, under surface generally fibrillose; veins pinnate or forked in the segments of the pinnules; sori towards the apex or at the apex of the lower veinlets. *Sw. Schrad. Journ. ii. 37. Hook. Syn. Fil. p. 252.*

Throughout the Indian region on the mountains.

(Also throughout the whole world.)

VAR. β LOBATUM. (*Eng. bot. t. 1563.*) Fronds narrowly lanceolate; pinnæ hardly pinnate, the lower secondary pinna sessile, or decurrent. *Clarke, F. N. I. p. 509.*

Throughout the Himalayas.

VAR. γ . RUFO-BARBATUM. (*Wall. Cat. 369.*) Fronds bipinnate, very coriaceous, usually reddish; stipes densely clothed with large red scales; rachis with red fibrillæ; pinnules naked and fibrillose beneath, in cutting like ilicifolium. *P. aculeatum*, *Bedd. F. S. I. t. 121. Polyst. brachypterum*, *Kunze, No. 906.*

Nilgiris and Western mountains of South India; Himalayas, from Kashmir to Bhotan.

VAR. δ . ANGULARE. (*Presl. Newm. 173.*) Lax and of thinner



POLYSTICHUM ACULEATUM, VAR. RUFO-BARBATUM. (Wall.)

ture, pinnules smaller, more numerous, orbicular, rhomboid, mostly auriculate, the serratures setiferous rather than spinulose. *Ed. F. S. I. t. 121.*

Nilgiris and Western mountains of South India.

VAR. ζ SEMIFERTILE. Base of the frond fertile, upper one-third barren. (*Clarke, l. c.*)
Sikkim.

VAR. η BIARISTATUM. (*Bl. En. Pl. Jav. Fil. 164.*) Pinnules large, oblong-falcate, sparingly serrate or spinulose; sori generally round the margin.

Khasya; Ceylon; Malay Peninsula.

VAR. θ SETOSUM. (*Wall. Cat. 371.*) Lower surface of frond with very long fibrillæ; rachis with very large scales as well as fibrillæ, pinnules small, quite entire, except the spinulose apex or with inconspicuous crenatures to represent the usual lobes; sori small on the lower veinlet of the forked or pinnate vein of the segment (or what would correspond to the segment where the pinnule is entire).

Himalayas, from Kumaon to Sikkim, 5,000–8,000 feet elevation. Well marked form, considered a distinct species by some botanists.

VAR. ι ANOMALUM. (*Hook. Sp. Fil. iv. 27.*) Segments blunt slightly mucronate; sori often, but not always, on the upper face of the fronds. *Polystichum anomalum, Bedd. F. S. I. 19.*


Ceylon; Horton plains and Happootalee, 5,000–6,000 feet elevation.

VAR. κ TRAVANCORICUM. (*Bedd.*) Pinnæ rather distant, each pair generally deflexed; pinnules prominently petioled, either entire, large broad deltoid, with the lobing blunt not mucronate, elongated up to 3 inches long, and pinnatifid or completely pinnate except an indistinct wing to the rachis), the pinnules distant and each bluntly lobed, 3–4 pair below the pinnatifid apex; rachis and petiole scaly and fibrillose, under surfaces fibrillose.

Travancore and Tinnevely mountains, 4,000–5,000 feet elevation. A distinct variety, but evidently a form of *aculeatum*, the more entire pinnules are nearest in shape and cutting to those of *anomalum* and *biaristatum*; the same frond will often have some pinnæ with the pinnules short and only slightly lobed, and others with them all elongate and pinnate or deeply pinnatifid.

VAR. λ MUCRONIFOLIUM. (*Bl. En. Pl. Fav. Fil.* 164.) Fronds 3-pinnate or sub-3-pinnate. (*Clarke, l. c.* 509.)

Khasya and Assam; this is very distinct looking, it is tripinnate like the last, but very different, being finely cut with small pinnules.



NO 105.
POLYSTICHUM PRESCOTTIANUM.
(*Wall.*)

9. POLYSTICHUM PRESCOTTIANUM. (*Wall.*) Stipes short, thick, flaccid, with many straw-coloured scales and fibrillæ; fronds from narrow to broad lanceolate, tapering at the base; pinnæ either short oblong, $\frac{1}{4}$ –1 inch long, more or less divided, but not to the rachis, or elongate, 2–3 inches long and deeply pinnatifid to the rachis or even pinnate; segments with the margin serrate and often hair-pointed, more or less fibrillose beneath. *Wall. Cat.* 363. *Bedd. F. B. I. t.* 34. *Hort. K. Syn. Fil.* 253.

Himalayas; from Kashmir to Bhotan, 10,000–13,000 feet elevation –

VAR. β BAKERIANUM. (*Clarke.*) Fronds large, very flaccid, broad-lanceolate, (9–10 inches broad), truncate at the base. *Clarke, F. N. I. t.* 66.

Sikkim; Yakla, 12,000 feet; very distinct-looking at first sight, but evidently only a flaccid variety of the above.

VAR. γ CASTANEUM. (*Clarke.*) Stipe round, naked, scales on the rachis blackish, pinnæ little pinnatifid.

Sikkim, 15,000 feet elevation.

GENUS L.—CYRTOMIUM. (*Presl.*)

(Derivation not known.)

Indusium orbicular, peltate ; veins pinnate from a central costa, lower anterior veinlets free, the rest angularly and irregularly anastomosing, forming unequal and subhexagonal areoles within which are 1–3 excurrent venules, or the upper veinlets only angularly anastomosing ; fronds pinnate, coriaceous.

1. CYRTOMIUM FALCATUM. (*Sw.*) Stipes tufted, 6–12 inches long, densely clothed below with large dark scales ; fronds 1–2 feet long, by 6–9 inches broad, pinnate ; pinnæ numerous, the lower ones stalked, ovate-acuminate, falcate, 4–6 inches long, by 1–2 inches broad, the edge entire or slightly undulated, the upper side narrowed gradually, sometimes auricled, the lower rounded or obliquely truncate at the base ; texture coriaceous, both surfaces naked, the upper one glossy, primary veins from the midrib of the pinnæ parallel to the other nearly to the margin, throwing off pinnately 1–2–3 times branching veinlets ; sori small, copious, scattered. *Swartz, Syn. Fil.* t. 3. *Hook. Syn. Fil.* 257.

VAR. β CARYOTIDEUM. (*Wall.*) Pinnules larger, sharply pointed, sometimes deeply lobed towards the base, and generally with a long sharp auricle on one or both sides at the base. *Wall.* t. 376. *Bedd. F. S. I. t.* 119.

Nilgiris at the higher elevations. Himalayas from Gurwhal to Shotan, 3,000–8,000 feet elevation. Khasya, 3,000–4,000 feet ; common. There is also a variety common on the Nilgiris known as *Proptera*, which has more numerous and much smaller pinnæ, scarcely at all auricled or lobed, but it graduates into caryotideum, though extreme forms look very distinct. The type of falcatum, which only differs slightly, is from China, Japan, South Africa, Sandwich Islands, and Madagascar.

2. CYRTOMIUM CADUCUM. (*Wall.*) Stipes tufted, 1 foot long, firm erect, stramineous, scaly below ; main rachises sometimes

with a terminal rooting bud ; fronds 1-2 feet long, 6-9 inches broad, pinnate ; pinnæ petioled, numerous, narrow lanceolate acuminate, from subentire to deeply lobed ; texture coriaceous ; primary veins from the midrib of the pinnæ repeatedly dichotomous at an acute angle into nearly equally strong veins, not anastomosing usually till near the margin, many of them free to the margin ; sori rather large and in two rows half-way between the rachis and the margin, or much smaller and scattered. *Wall. Cat.* 381. *Hook. Syn. Fil.* 257. *Bedd. F. B. I. t.* 45. *Clarke, F. N. I. p.* 512.

Himalayas ; from Nepal to Bhotan, 4,000-7,000 feet ; Khasya, 3,000-5,000 feet elevation.

GENUS LI.---ASPIDIUM. (*Swartz.*)

(*Aspidos*, the indusium being like a shield.)


Indusium orbicular or reniform, or sometimes irregular and abnormal, being linear and curved, or sometimes absent ; veins compoundly anastomosing with generally free veinlets in the areoles, receptacles compital or often at the apex of the free veinlets ; fronds very various, from simple to tripinnatifid, often membranaceous and flaccid.

1. ASPIDIUM SINGAPORIANUM. (*Wallich.*) Rhizome creeping ; stipes subtufted, those of the fertile fronds 6-12 inches long, fronds 12 inches or more, 2-4 inches broad, oblong entire, acuminate at the apex, narrowed rather suddenly and then decurrent gradually at the base ; texture herbaceous ; primary veins distinct nearly to the edge, united by transverse arched ones, the areoles of which enclose netted veinlets, the ultimate ones free ; sori numerous, close, 4-6 in a line between the main veins ; indusium peltate orbicular. *Nephrodium* (Baker), *Hook. Syn. Fil.* 296. *Bedd. F. B. I. t.* 168.

Singapore and the Malay Peninsula.

(Also in the Malay Islands.)


2. ASPIDIUM VASTUM. (*Blume.*) Rhizome creeping, stipes scattered, narrowly winged often nearly or quite to the base, furnished with numerous linear subulate brown persistent scales ; fronds



GENERA


CYRTOMIUM FALCATUM, VAR. β CARYOTIDEUM. (Wall.)

2-4 feet long, cut down to a broadly-winged rachis, into entire linear-oblong pinnæ 6-12 inches long, by 1-2 inches broad; texture subcoriaceous; main veins distinct nearly to the margin, connected by numerous prominent cross veins forming 7-8 areoles between the costa and margin, with netted and free veinlets in the areoles; sori small, scattered on the netted veins; indusium reniform. *Bl. En. Pl. Far. Fil.* 142. *Hook. Syn. Fil.* 296. *Aspid. alatum*, *Wall. Cat.* 378. *Sagenia alata*, *Bedd. F. B. I. t.* 169.



N°107
ASPIDIUM SINGAPORIANUM.
(*Wall.*)

East Bengal, up to 1,000 feet elevation; Mishmee; Khasy—
Chittagong; Malay Peninsula.
(Also in the Malay Islands.)



N°108
ASPIDIUM VASTUM. (*Blume.*)

3. *ASPIDIUM SUBCONFLUENS.* (*Bedd.*) Stipes tufted, 1-2 feet long, furnished with dark coloured linear-subulate scales; fronds about 1 foot each way, triangular in outline, tripartite, lateral pinnæ opposite or subopposite, shortly petioled, unequal sided with the

er pinnules small falcate subentire, the lower ones much produced lanceolate acuminate, the 2-3 basal ones cut about one-third into falcate lobes, terminal pinnæ deeply pinnatifid, lobes 3-6 lobes, pinnatifid a quarter of the way down to the midrib; texture moderately firm, surface glabrous above, pubescent on the veins beneath; veins anastomosing with a few free included veinlets; sori clustered, small, generally on the netted veins; indusium variable, uniform, curved, horseshoe-shaped or nearly peltate, persistent. *edd. Sup. Ferns, t. 364.*

Khasya, 3,000-5,000 feet elevation, below Umwai.

4. *ASPIDIUM SEMIBIPINNATUM.* (*Wall. under Polypodium.*) Stipes 1 foot or more long, pale brown, naked; fronds 12-18 inches long, 6-9 inches broad, terminal pinnæ ear-oblong, 6-9 inches long, each broad, entire, narrowed towards both ends, lateral pinnæ 4-6 on each side smaller, the lower one or two bipartite; texture hercynaceous; veins inconspicuous, the primary ones lost before they reach the margin, the secondary variously anastomosing, the areoles with free included veinlets; sori minute, scattered,



Nº 109

ASPIDIUM SUBCONFLUENS. (*Bedd.*)


confined to the connected veinlets; indusium reniform and irregular in the last. *Hook. Sp. Fil. iv. 59, t. 231. Syn. Fil. 297. Bedd. B. I. t. 137. Polypodium semibipinnatum, Wall. Cat. 388.*

Penang; nearly allied to some forms of polymorphum, but I think sufficiently distinct.

(Also in Borneo.)

5. *ASPIDIUM SUBTRIPHYLLUM.* (*Hook.*) Rhizome creeping;

Hook. Sp. Fil. iv. 52. *Syn. Fil.* 296. *Bedd. F. S. I. t.* 242, and
F. B. I. t. 48.



ASPIDIUM SUBTRIPIHYLLU M.


(Hook.)

Ceylon and the Malay Peninsula. My Ceylon specimens are much smaller than those from Birma, but I believe they are the same species; in habit it is quite like *variolosum*, but the sori are always on the netted veinlets, instead of at the apex of free veinlets.

6. ASPIDIUM VARIOLOSUM. (*Wall.*)

Rhizome suberect, stipes tufted, scaly near the base only, 12-18 inches long in the fertile, less in the sterile fronds, more or less dimorphous, deltoid in outline, herbaceous in texture, and somewhat puberulous, especially in the sterile ones; pinnae about three pair below the pinnatifid apex, the lowest pair again pinnated with the pinnules often pinnatifid, the next pair often pinnatifid c pinnated in a less degree, the pinnae c

the fertile fronds generally narrower and often more divided, *ma*
veins not very distinct but more so in the fertile than in the sterile
areoles copious, with many free included veinlets; sori rather larg
almost always at the apex of the free veinlets, very rarely on the nette
veins; indusium peltate and reniform on the same frond. *Wal-*
Cat. 379. *Redd. Sup. Ferns*, t. 365 (not *F. B. I.* 170, which i



ASPIDIUM VARIOLOSUM. (Wulf.)

cicutarium.) *Aspidium Zollingerianum*, *Bedd. F. B. I. t.* 251, *fertile frond only*. *Nephrodium* (*Sagenia*) *variolosum*, *Hook. Sp. Fil.* 29 and *Zollingerianum* (*l.c.*) *as far at least as the Birmese plant*.

East Bengal Plains, abundant from Assam to Chittagong ; Tenasserim near Moulmein ; Penang.

7. *ASPIDIUM POLYMORPHUM*. (*Wall.*) Rhizome suberect, stipes tufted, yellowish or brown, paleaceous only at the base ; fronds large, 1-4 feet long, by 1 foot or more broad, pinnate ; pinnae 3-6 on each side, oblong or elliptic, acuminate, unequal at the base, sometimes contracted when fertile, quite entire to crenate or coarsely toothed, stalked or subsessile, the terminal one often more or less lobed or subpinnatifid, the lowest pair generally (not always) bifurcate ; texture herbaceous to subcoriaceous ; main veins prominent and distinct to the margin, with many free included simple or forked veinlets ; sori on the netted veins, small and scattered in the uncontracted fronds, large and more or less in two rows between the main veins in the contracted ones ; indusium reniform or often quite absent. *Wall. Cat.* 382. *Hook. Syn. Fil. p.* 297. *Bedd. F. S. I. tabs.* 116, 117. *Sagenia siifolia*, *Baker, Syn. Fil. p.* 29, *as far as the Courtallum plant*. *Neph. Wightii*, *Clarke, F. N. I.* 538, *the contracted form*.

Western forests of Madras Presidency, very common up to 4000 feet. (I do not consider the contracted form even a constant variety.) Northern India from Gurwhal to Mishmee and Chittagong ; Ceylon ; Birma.

(Also in the Malay Islands, the Philippines and Fernando Po.)

8. *ASPIDIUM SIMONSHI*. (*Baker.*) Very like *polymorphum*, but with the stipes and rachis polished blackish-purple, lowest pair of pinnae pinnate below, deeply lobed upwards, others more or less bifurcate below and rather deeply lobed throughout, venation as in *polymorphum* ; sori rather large (in my specimens), in two series between the main veins, or irregularly scattered sub-4-serial between the main veins. *Hook. Syn. Fil.* 504. *Bedd. Sup. Ferns. t.* 367.

Seetakand Hill, Chittagong ; Sikkim, and Bhotan. Mr. Clarke makes it a variety of *polymorphum*, and he may be right, but until

re specimens are gathered, I consider it safer not to lump it with that. Mr. Clarke says he does not see how it differs from *polymorphum*, var. *macrocarpum*, but he has made a new species of that. Both Baker and Clarke make two sections of *Sagenia*, one with "sori in more than two rows between the principal veins, often irregularly scattered," the other with "sori in two rows between the principal veins," both forms occur in this species and in *polymorphum*, and more or less in *semibipinnatum* and other species, the definition is only misleading.

9. *ASPIDIUM HETEROCARPUM*. (*Bedd.*) Rhizome creeping densely; stipes solitary, erect, with lanceolate-linear brown persistent scales at the base; pinnate, very much as in *polymorphum*, but lower pinnae not bifurcate; pinnae 8 inches long, narrow-lanceolate, entire, caudate at the apex, sessile or very shortly stalked, uniparous in the axils; main vein distant to nearly the margin, venules copious with free included veins; sori very small on reticulated veins, much scattered; sori reniform or horseshoe-shaped, or sometimes curved or as in *Athyrium* and *Asplenium*.



Nº 112

ASPIDIUM HETEROCARPUM. (*Bedd.*)

Sagenia heterocarpa, *B. I. t.* 47. *Sagenia*

ra (Baker), *Hook. Syn. Fil.* 504. *Clarke, l. c.* 537.

in and Chittagong, in wet flats near rivers, forming large clumps about 6 feet high.

ASPIDIUM DECURRENS. (*Presl.*) Rhizome creeping; stipes solitary or quite to the base, furnished with numerous linear-lanceolate brown persistent scales; fronds often 3 feet long, dimor-

phous, the fertile much contracted, pinnatifid down to a broadly winged rachis into 3-4 pair of pinnæ or lobes on each side, 6-12 inches long, 1-2½ inches broad, lowest pair often bifurcate, margins quite entire in the sterile, more or less crenated in the fertile, subcoriaceous, glabrous on both sides; main veins distinct nearly to the margin, with cross veins forming series of large areoles in which are included netted veinlets and simple or forked free veinlets; sori large, generally in two pretty regular series between the main veinlets, nearly always at the apex of the free veinlets, often continued all down the wing of the stipes; indusium generally more or less reniform, sometimes elongated and irregular. *Presl. Rel. Hanck*, 28. *Hook. Syn. Fil.* 299. *Sagenia pteropus*, *Bedd. F. S. I. t.* 82.

Eastern Bengal at the foot of the hills, Assam, Cachar, Chittagong; South India in the Bolampatty Hills between Coimbatore and Palghat, 2,000 feet elevation; Tinnevely and Travancore Hills, 2,000-2,500 feet; Ceylon; Malay Peninsula. In habit very like *vastum*, but fertile fronds contracted, and sori differently situated.

(Also in the Malay Islands, South China, Polynesia.)

VAR β MINOR. (*Bedd.*) Fronds very small (1 foot long only), sterile 3-5 partite rather thin pinnatifid, fertile deeply pinnatifid, very much contracted, the lobes or pinnæ being very narrow. *Sagenia pteropus-minor*, *Bedd. F. S. I. t.* 245.

Ceylon only, in the Doombera district at no great elevation.

11. *ASPIDIUM CICUTARIUM*. (*Sw.*) Stipes tufted, 1 foot or more long, with many scales close to the base; fronds very variable in size, from 8 inches up to several feet, deltoid in outline, apex pinnatifid, below which are 1-4 pair of pinnæ, which are either only lobed or completely pinnated with the secondary pinnæ deeply pinnatifid, or even again pinnated (*i.e.* tripinnate), with the tertiary pinnæ deeply pinnatifid (quadripinnatifid), the lowest pair of pinnæ are always deeply pinnatifid or pinnate on the lower margin, but the other pairs are generally much less so and often only lobed $\frac{1}{3}$ - $\frac{1}{2}$ down, texture thin, surfaces glabrous in some varieties, but in others hairy on the upper surface and puberulous on the costas and costules



L. A. C. FAG. 745

ASPIDIUM DECURRENS, VAR. β MINOR. (Bedd.)

beneath ; principal veins tolerably conspicuous to the margin, others copiously netted with often free included veinlets (there are no prominent cross veins forming large well-defined areoles as in some of the species, and the free veinlets are often few only) ; sori rather large, in two rows between the main veins, on the netted veins (compital) or at the apex of the free veinlets ; indusium reniform or peltate. *Swartz. Mett. Farngatt. Pheg. and Asp.* 117. *Hook. Syn. Fil.* 299. *Sagenia coadunata* (Wall.), *Bedd. F. S. I. t.* 81. and *t.* 170 (*variolosa*).

Throughout the Indian region, from the plains up to 5,000 feet. In South India (both Eastern and Western sides), the more hairy variety known as *coadunata* only is found, but in Northern India the glabrous variety is also common.

(Also in the tropics throughout the globe.)

12. *ASPIDIUM MULTICAUDATUM*. (*Wall.*) Stipes 1-3 feet, often densely clothed nearly its whole length with linear-subulate brown persistent scales ; fronds 3-4 feet long, with a large terminal pinna deeply pinnatifid, lobes lanceolate acuminate, and several pinnæ on each side, the lowest deltoid 1 foot or more long and nearly as broad, deeply pinnatifid above and pinnate below ; texture firm herbaceous, rachises beneath rusty ; main veins distinct to the margin, areoles with free included veinlets ; sori rather large in two rows between the principal veins, or more numerous and scattered, generally at the apex of free veinlets, rarely on the netted veins (compital). *Wall. in Herb.* *Nephrodium Griffithii* (Baker), *Hook. Syn. Fil.* 300. *Sagenia Griffithii*, *Bedd. F. B. I. t.* 337.

Khasya Hills, South side up to 1,000 feet elevation. Birma *Aspidium*? *t.* 118, *F. S. I.*, is probably this species, or at least closely allied, it has twice been gathered in the Anamallay forests, but young fronds without fructification.

In the *Synopsis Filicum* *Sagenia* is kept up as a section of *Nephrodium* for all species supposed to have a reniform indusium, and *Euaspidium* as a section of *Aspidium* for those with an orbicular indusium, the result of this has been that allied species like *repandum* and *polymorphum* (probably one and the same plant) and *membranaceum* and (*Trimeni*) *giganteum* (closely allied if not

varieties one of the other), are placed far apart ; it is a fact, however, that most of the species have both orbicular and reniform involucres, sometimes on the same individual, the involucres often varying much more than this, being athyrioid and asplenoid. Mr. Clarke has placed both *Aspidium* and *Sagenia* in *Nephrodium*, a genus widely different in habit.

GENUS LII.—PLEOCNEMIA. (*Presl.*)


(*Pleos*, full ; *knemia*, rays ; in allusion to the venation.)

Only differs from *Aspidium* in its less compound venation, only the lower veins anastomose arcuately forming one series of areoles near the costa (as in *Campteria*), without any free veinlets within them, the other veins generally all free, or some angularly anastomosing ; sori generally at the apex of the free veinlets outside the areoles, but sometimes some are on the netted veins, (*i.e.*, compital).

1. PLEOCNEMIA THWAITESII. (*Bedd.*) Stipes tufted 1–1½ feet long, shining blackish-purple as is the rachis and main costa, basal scales linear dark-brown ; fronds deltoid, 1½–2 feet each way, tripinnatifid ; lowest pinnæ largest deltoid stalked, with several large lanceolate deeply pinnatifid pinnules ; texture rather thick and opaque, the veins not conspicuous, glabrous beneath, and also above except the costa and veins which are scurfy or puberulous, lower veinlets forming elongate costal arches, no free included veinlets, the rest free with clavate apices, none reaching the margin ; sori rather large in a single series on each side of the midrib of the segments, all at the apex of free veinlets ; indusium reniform, with the lobes of the sinus broad and overlapping, so that it looks orbicular. *Sagenia Thwaitesii*, *Bedd F. S. I. t. 244*. *Nephrodium* (*Sagenia*) *simulans* (*Baker*), *Hook. Syn. Fil.* 300. Ceylon, southern provinces near Galle, abundant on the Haycock Mountain. (*C. P.* 3331. *Thw. En. p.* 390.)

2. PLEOCNEMIA TRIMENI. (*Bedd.*) Stipes tufted, 1–1½ feet, grey, basal scales linear ; fronds 1½–3 feet long, deltoid or deltoid lanceolate, pinnatifid at the apex, below pinnate with about 5–6 pair of lateral pinnæ, which are more or less stalked, and all pinnatifid

more than half-way to the rachis, about 8-9 inches long by 2 b and a much larger basal pair which are deltoid, again pinnated the lower basal pinnules deeply pinnatifid, ultimate lobes blunt slightly crenated or nearly entire; texture rather thin; surfaces glabrous on both sides; costas and veins rusty above, less so or nearly glabrous beneath; lower veins forming elongated costal arches near the rachis of the pinnae, with generally a second series of areoles formed by the veins again anastomosing angularly, veins of the segments only anastomosing near the midrib and forming one series of arches, the rest free, no free veinlets in the areoles; sori in a single row on each side of the midrib, those towards the margin and apex of the segments apical on the free veinlets, those lower down and near costa of the pinnules generally on connected veins (*i.e.*, *compitodiusium reniform*. *Sagenia gigantea*, *Bedd. F. S. I. t. 80* (*description*.) *Pleocnemia gigantea* (Baker), *Hook. Syn. Fil. Edit. p. 503*, *not Blume*.



PLEOCNEMIA TRIMENI. (*Bala*

The Kew packet of *giganteum* from which Sir W. Hooker his description (*Sp. Fil.* iv. p. 50,) contains several different specimens the Java specimen being a single pinna of *Aspidium giganteum* species distinct from this; the Birma and Assam specimens *multicaudatum* (Wallich); and the Ceylon specimens being *plant, Pleocnemia membranacea*, and *P. Thwaitesii*. This not Blume's *giganteum* I have had to rename it.

Ceylon, central provinces, (*C.P.* 1357); South India, Anam Palghat side, in the forests near the Nelliampady coffee estates


3. PLEOCNEMIA MEMBRANACEA. (*Hook, under Aspidium.*)

Stipes tufted up to 9 inches long, straw-coloured, basal scales linear; fronds up to about 18 inches long, deltoid, bi-tripinnate, pinnatifid at the apex, the lowest pair of pinnæ much the largest, long stalked, with the lowest secondary pinnæ stalked and again quite pinnate at the base, the pinnules being stalked and deeply pinnatifid, ultimate segments crenulate, upper pinnæ gradually less compound; texture thin, membranaceous, upper surface somewhat hairy towards the margins; costas and veins rusty or puberulous, under surface slightly puberulous; lower veins forming long costal arches, the rest free; sori at the apex of free veins; indusium both orbicular and reniform. *Hook. Syn. Fil. p. 259.* *Sagenia gigantea*, var. *minor*, *Bedd. F. S. I. t. 243*, a small frond less compound than described above. *Thw. En. Pl. Zey. p. 390.* (*C. P. 1358.*)

Ceylon; in the Doombura district at no great elevation. Mr. Thwaites had doubts whether this was a variety of the last, or a distinct species; the venation is quite that of the last species, but rather less compound, owing to the fronds being more cut and narrower; it is quite that of *Pleocnemia*, and I find some of the involucre reniform; it is of thinner texture, much more cut and less glabrous than the last, and, I think, may safely be considered a distinct species.

(Also in Java, Philippines, China, and Formosa.)

4. PLEOCNEMIA MEMBRANIFOLIA. (*Presl.*) Stipes tufted, furnished with black linear subulate scales more copious towards the base, up to about 1 foot long; fronds seldom more than 1 foot long, from deltoid to ovate, pinnate with a pinnatifid apex, lateral pinnæ few pairs opposite or subopposite, 2-4 inches long, $\frac{3}{4}$ -1 $\frac{1}{4}$ broad, more or less deeply pinnatifid, the segments rounded nearly entire, lowest pair of pinnæ much larger, deltoid, pinnatifid or almost pinnate, the lower basal segments being elongated and pinnatifid; main rachis and both surfaces more or less softly hairy, with multicellular hairs; texture softly herbaceous; fertile fronds uniform with the sterile, or more or less contracted, often very much so, lower veins in the sterile and broad fertile fronds anastomosing and



PLEOCNEMIA MEMBRANIFOLIA. (*Prest.*)

forming loops near the costa, sometimes very regularly, sometimes only occasionally, in the contracted fertile fronds all the veins free; sori generally apical on the free veinlets; involucre reniform. *Presl. Rel. Haenck.* 36, t. 5, fig. 3. *Aspidium fuscipes*, *Bedd. Sup. Ferns*, t. 366. *Aspidium fuscipes*, *Wallich*, partly, but not the type sheet, which is *Lastrea sagenioides*.

East Bengal Plains, extending into Assam, Cachar, and Chittagong. Khasya and Sikkim hills up to 3,000 feet elevation. Birma and the Malay peninsula. In habit much like small *Aspidium cicutarium*, but venation different, and easily known by the persistent black scales towards the base of the stipe; in the *Synopsis Filicum* it has been erroneously lumped with *Lastrea dissecta* by Hooker and Baker.



NO 116
PLEOCNEMIA CLARKEI. (Bedd.)

5. **PLEOCNEMIA CLARKEI.** (Bedd.) Stipes and rachis slightly pubescent; fronds 2-3 feet long, lanceolate, narrowed at the base into distant auricles, softly shortly villous, herbaceous, pinnate; pinnæ numerous, alternate, $1\frac{1}{4}$ -2 inches broad, pinnatifid nearly to the rachis, the pinnules lanceolate from a broad base, irregular as to length, and from subentire to deeply pinnatifid (on the same pinna) acute at the apex, lowest veins forming arcs along the costa of the pinnules from which proceed free veinlets; veins of the pinnules pinnate, or once or twice forked, the lowest veinlet of a group often looped with the next group; involucre kidney-shaped, hairy and ciliate. *Bedd. Sup. Ferns*, t. 368. *Nephrodium artinexum*, *Clarke, F. N. I.* p. 536.


Sikkim; near Dikeeling, 4,500 feet elevation.

6. *PLEOCNEMIA LEUZEANA* (*Hook.*) Caudex subarborescent, densely scaly at the crown, stipes 2-3 feet long, stout, striated, shortly hirsute without scales; fronds up to 6 feet long, subdeltoïd, bipinnate; pinnae 1-1½ foot long, 6-8 inches broad, simple, or the lowest with 2-3 large (12-14 inches long) secondary pinnae from the lower side, which are again pinnate with the pinnules stalked, 2-4 inches long, and deeply pinnatifid, ultimate segments oblong-rounded, entire or denticulate, usually with a tooth in the sinus at their base; lower veins anastomosing in costal arches near the costae of the pinnules and along the midrib of the segments, other veins

free and excurrent, or all the veins of the segment are free; sori copious, usually in single rows on each side of the midrib, mixed with yellow glandular hairs; indusium reniform, early fugacious. *Hook. Syn. Fil.* 295. *Bedd. F. B. I. t.* 134-

North and East Bengal, base of the hills up to 2,000 feet elevation, Sikkim, Assam, Cachar, Khasya, Chittagong; Birma, and Malay Peninsula.

(Also in Malay Islands, China, Polynesia, and North Australia.)



N°117

PLEOCNEMIA LEUZEANA.
(*Hook.*)

GENUS LIII.—LASTREA. (*Presl.*)

(After Chev. de Lastre, a French nobleman.)

Sori subglobose, dorsal or terminal on the veins; indusium reniform, attached by the sinus, or in the first four species sometimes orbicular; veins all free; fronds pinnate or compoundly 2-4 pinnate.

* *Indusium reniform or orbicular, texture more or less coriaceous.*

1. *LASTREA AMABILIS.* (*Bl. under Aspidium.*) Rhizome

creeping; stipes scattered, 6–12 inches long, slender, slightly scaly below; fronds about 1 foot long, by 6–9 inches broad, with a pinnate apex and 2–6 lateral pinnæ on each side, which are 1–1½ inch broad, all simply pinnated or the lowest sometimes slightly compound at the base, pinnules subrhomboidal with the lower side much cut away ¼–1 inch long, cut down about a third or half-way to the costa into rounded lobes which terminate with a long bristle, sometimes being more or less serrated; texture subcoriaceous; rachis and both surfaces naked; veins pinnate in the lobes; sori submarginal; indusium reniform or orbicular. *Bedd. F. S. I. t. 109.* *Aspidium amabile*, *Bl. En. Pl. Jav. Fil. p. 165.* *Hook. Syn. Fil. 254.* *Aspid. rhomboideum*, *Wall. Cat. 364.*

I keep this species in *Lastrea* as I have always found the involucre reniform; they appear, however, to be sometimes orbicular, so that the species has equal right to be a *Polystichum*.

South India, on the Anamallays, Lower Pulneys, Travancore, and Tinnevely Hills, 3,000–4,000 feet elevation; Ceylon, central provinces; North India, Nepal, Jaintea, about 4,000 feet, rare.

(Also in the Philippines and South China.)

2. *LASTREA ARISTATA*. (*Sw. under Aspidium.*) Rhizome long, creeping, very scaly; stipes scattered, about 1 foot long, more or less fibrillose, as are the main and partial rachises; fronds about 12–15 inches long, deltoid in outline, pinnate at the apex, below which are 5–6 lateral pinnæ on each side, which, except the lower pair, are generally simply pinnate or with a tendency to be bipinnate in their upper basal pinnules, lower pair bipinnate towards the base, the lower basal pinnules being much the largest and quite pinnated, pinnules subrhomboidal with the lower base cut away, more or less lobed, the upper basal lobe being the deepest, lobes copiously toothed, teeth aristate; texture more or less coriaceous; veins pinnate in the lobes, the veinlets being simple or forked; sori apical on the veinlets and well within the margin; indusium reniform or orbicular. *Bedd. F. S. I. t. 101.* *Aspidium aristatum*, *Sw. Syn. Fil. 53.*

South India, very general in the Western forests of the Madras Presidency; Ceylon; Birma.

I have always found the involucre strictly reniform, and Hooker has described it both under *Polystichum* and *Lastrea* (in the last as *Lastrea platypus*); it is a much less divided fern than *coniifolia* besides having an utterly different caudex; Mr. Clarke, who unites it with *coniifolia*, states, that he has never been able to see the creeping caudex accompanies a less divided form, but Mr. Clarke's specimens are all typical *coniifolia*, and there is no *aristata* from the Himalayas in the Kew Herbarium, and only one poor specimen from Khasya, labelled *Lastrea platypus*.

(Also in Japan, Formosa, South China, Java, Luzon, Norfolk Island.)


3. *LASTREA AFFINIS*. (*Wall.*) Very similar to *aristata* or *amabilis*, but the lower pinnæ less compound than in *aristata* often quite similar to the upper ones; texture much more coriaceous and sori very large. *Wall. Cat.* 370. *Lastrea aristata*, var. *Hamiltonii*, *Bedd. Fern. Sup. pl.* 369. *Hook. Syn. Fil.* 256. *Aspidium speciosum* (Don.), *Aspidium aristatum*, var. *affinis*, and *assamica*, *Clarke, l. c.* 511.

I formerly considered this species a variety of *aristata*, I have never seen it growing, but Mr. Clarke tells that the rhizome is erect and not creeping, so I cannot consider it a variety of either *aristata* or *amabilis*.

Sikkim, Nepal, Assam, Jaintia.

4. *LASTREA CONIIFOLIA*. (*Wall.*) Rhizome erect; stem tufted; fronds very large, ovate, 2-3 feet long, 4-5 pinnate, ultimate pinnules and segments small, coriaceous shining, aristate; sori large or small; indusium reniform or orbicular. *Bedd. F. B. I.* 261. *Aspidium*, *Wall. Cat.* 341.

I have always found the involucres reniform in their young stage, I have had both this and *aristata* in cultivation on the Nilgiris for many years, and they show no tendency to run one into the other.



LASTREA CONIIFOLIA. (Wall.)

A—Root of *L. aristata*.


Himalayas, Kumaon to Bhotan, 4,000–10,000 feet, Khasya, 3,000–6,000 feet; South India on the Western mountains; Ceylon; Malay Peninsula.

(Also in Natal, Samoa, and Australia.)

* * *Indusium* always reniform.

† *Pinnæ* subentire or only slightly lobed (except in forms *v* and *δ* of *Walkeræ*.)

5. *LASTREA CUSPIDATA*. (*Mett. under Aspidium.*) Stipes about 1 foot long, reddish as is the rachis, scaly towards the base; fronds 1–3 feet long, pinnæ numerous, linear-lanceolate from a broad base, 6 inches long, $\frac{1}{4}$ – $\frac{3}{4}$ inch broad, cuspidate at the base, very shallowly lobed (not $\frac{1}{2}$ to the rachis), the lobes sharply serrated; texture scarcely coriaceous; rachis and both surfaces naked; veins pinnate with 3–4 veinlets on each side; sori towards the base of the veinlets; indusium reniform fugacious. *Mett. Farngatt. Phlegm. and Asp.* 92. *Polypodium elongatum*, *Wall. Cat.* 309. *Bedd. F. B. I. t.* 118. *Nephrodium cuspidatum*, *Hook. Syn. Fil.* 260.



Nº 119.

LASTREA CUSPIDATA.
(*Mett.*)

Khasya Hills, 3,000–4,000 feet elevation, near Shillong.

6. *LASTREA HIRTIPES* (*Bl. under Aspidium.*) Caudex erect; stipes tufted up to about 1 foot long, densely clothed, as is the rachis, with long blackish hair-like scales; fronds up to 3 feet long, pinnæ numerous, linear-lanceolate from a broad base, 4–8 inches long, by nearly 1 inch broad, with broad shallow lobes reaching $\frac{1}{4}$ to $\frac{1}{2}$ towards the rachis, lower ones not reduced; texture herbaceous,

surfaces naked; veins pinnate; veinlets 4-5 on a side; sori low down on the veinlets, indusium reniform. *Bl. En. Pl. Jav. Fil* 148. *Bedd. F. S. I. t.* 96. *Asp. atratum*, *Wall. Cat.* 380.

Himalayas, 5,000-9,000 feet, from Nepal to Bhotan, Khasya 4,000-6,000 feet; South India, Nilgiris, &c.; Ceylon; Birma.

(Also in the Malay Islands and Polynesia.)

7. *LASTREA WALKERÆ*. (*Hook under Polypodium*.) Stipes up to 1½ foot long, clothed with linear scales; fronds up to 3 feet long, oblong-lanceolate, pinnate; pinnæ linear-lanceolate, short-stalked, acuminate, slightly crenate along the margins, lower ones about 6 inches long, by about 1 inch broad, lower base cut away, upper base square; texture subcoriaceous, glabrous; veins in pinnate groups; sori 2-3 serial, between the midrib and the margin, dorsal on the veinlets; indusium reniform, very fugacious. *Polypodium Walkeræ*, *Hook. Syn. Fil.* 1st ed. p. 305. *Lastrea*, 2nd ed. 493. *Bedd. F. S. I. t.* 234.

Ceylon.

VAR. β *MACROCARPA*. Fronds very large, the pinnæ slightly lobed, often furnished with a large auricle at the superior base. *Bedd. F. S. I. t.* 235.

Ceylon.

VAR. γ *PINNATIFIDA*. Pinnæ pinnatifid nearly to the rachis, leaving only a broad wing along the rachis, lower ones sometimes pinnate. *Bedd. l. c. t.* 235.



LASTREA HIRTIPES. (Bl.)

VAR. ♂ BIPINNATA. Pinnæ sub-bipinnate. *Bedd. l. c. t. 235.*
Ceylon.

† † *Pinnæ cut more than half-way to the rachis, compound in some forms of Filix Mas.*

A. Veins generally simple.

8. *LASTREA GRACILESCENS.* (*Bl.*) Stipes tufted, long, slender; fronds oblong-lanceolate, pinnate, lower pinnæ scarcely reduced, often deflexed, rachises pilose above, pinnæ cut down nearly to the rachis, segments narrow-oblong, obtuse, not much narrowed upwards, veins about six on each side of the costule, the two lowest terminating above the sinus, all generally simple, rarely forked, sori medial on the veins, indusium glabrous or minutely pilose. *Aspidium gracilescens*, *Bl. En. Pl. Jav. Fil.* 155. *Hook. Syn. Fil.* 262. *Bedd. F. B. I. t.* 253. *Clarke, F. N. I.* 513 in part.

Mr. Clarke includes here *L. flaccida* (Hooker), which is certainly a mistake; he also includes the Assam forms of *L. immersa* (*Bedd. F. B. I.* 252) united with the Malay Peninsula plant by Hooker; in this he may be correct, as this plant appears distinct from the Malay one, it is however, much larger than any of Mr. Clarke's specimens of *gracilescens*, and has more veins to the segments.

Nepal, Sikkim, 6,000–8,000 feet elevation, Khasya, 4,000 feet; South India, Mudumally forests, a small form.

(Also in China, Japan and Java.)

Mr. Clarke's variety *decipiens* (Darjeeling, 7,000 feet, and Khasya, Dingling, 5,000 feet) has the fronds shorter and more deltoid, and the veins often branched with the sorus subterminal on the upper branch; it is probably a distinct species. His variety *hirsutipes* only differs in having the stipes hirsute and in being of rather thicker texture (Khasya and Jaintea, 4,000–5,000 feet); his variety *didymochloenoides* has longer narrower pinnæ with very large sori, but runs into the type.

9. *LASTREA IMMERSA.* (*Bl.*) Caudex creeping, stipes elongated 3–4 feet long, fronds rather coriaceous in texture, pinnate, ovate

long, or oblong in outline, pinnæ 6–10 inches long, sessile, about 1 inch broad, cut down close to the rachis, glabrous above on the segments, or with occasionally a few distant inconspicuous hairs on the veins, the rachis strigose, frosted beneath on the veins and the rachis with minute yellow glandular dots; segments linear, rather distant; veins simple (or the lowest one very rarely forked), 12–14 even more on each side of the costa; sori medial, one to each segment, much immersed, the depression plainest on the upper side, annulus large, persistent, reniform, but appearing orbicular from the overlapping at the sinus. *Aspidium immersum*, *Bl. En. Fil.* 156. *Phrodium immersum*, *Hook. Sp. Fil.* 112. *Syn. Fil.* 263.

Malacca and the Malay Peninsula.

(Also in the Philippines and New Caledonia.)

Mr. Clarke considers that the Assam plant (*Bedd. F. B. I. t.*) should be referred to *gracilescens*, it is of a different texture, less downy, and with much fewer veins.

10. *LASTREA CALCARATA*. (*Bl.*) Caudex erect, stout, stipited, fronds lanceolate pinnate, sometimes with auricles below, up to about 2 feet long, pinnæ spreading 1–4 inches long by $\frac{1}{2}$ inch to nearly 1 inch broad, cut down two-thirds or more towards the rachis into oblique falcate linear oblong segments, the upper basal one sometimes considerably longer than the others, rachis villous, under surface more or less villous, upper surface villous, or glabrous except the secondary rachis, veinlets 3–8 on a side, simple, sori medial, involucre glabrous or villous. *Asp. calcaratum*, *Bl. En. Fil. Jav. p.* 159. *Lastrea calcarata*, *Hook. Sp. Fil. iv.* 93, and *falciloba*, *p.* 108.

VAR. *CILIATA*, type (*Wall. Cat.* 351), texture herbaceous, lowest pinnales deflexed, no auricles on the stem; stipes rounded, pinnæ about 4 inches long, caudate at the apex, nearly 1 inch broad; indusium hairy or glabrous. *Lastrea falciloba*, *Bedd. F. S. I.* 105. *Asp. ciliatum*, *Wall. Cat.* 351, and *Asp. canum*, *Wall. Cat.* 387. *Asp. ciliatum*, *Clarke, F. N. I.* 515, except syn. *L. Bergiana* (*Bedd.*), which is *L. cana* (*Baker*).



LASTREA CALCIVATA, VAR. CILIATA. (H. S. Gentry)

South India, very common in all the Western forests; and in Ceylon (exactly corresponding with Wallich's two type sheets). Specimens sent from both these localities were partly referred by Hooker to his *calcarata* and partly to *falciloba*, but he doubted whether the two species were really distinct. Also in Khasya, the Himalayas; Birma.

ASP. CALCARATUM, VAR. β . *Thw. En.* 391, *L. calcarata, Bedd. F. S. I. t.* 246, is an abnormal form, with small narrow pinnæ tapering at both ends and less pinnatifid, except sometimes quite at the base, where the segments are almost free and distant; but, as Thwaites says, it passes into the type. A form from Birma also has very narrow pinnæ and is densely hairy.

VAR. β SERICEA. (*J. Scott, M.S.*) Pinnæ short oblong, 1–2 inches long by $\frac{1}{2}$ – $\frac{3}{4}$ inch broad, quite obtuse at the apex, or ending in a short sudden point (never caudate), involucre glabrous, otherwise as in the type. *L. sericea, Bedd. F. B. I. t.* 308. This is, perhaps, scarcely distinct as a variety from the type, but its geographical limits are curious, in North India it has only been found in Chittagong, elevation 200 feet, and in South India only on the Jeypore Hills west of Vizagapatam, elevation 2,000 feet. I have had it for several years in cultivation and it quite kept its character.

VAR. γ FALCILOBA. (*Clarke.*) Stipes angled and furnished with auricles below the frond; pinnæ more numerous and narrower, 4 inches long by $\frac{1}{2}$ inch broad; texture subcoriaceous (more like *ochthodes*); indusium glabrous. *Lastrea falciloba, Hook. Sp. Fil.* iv. p. 108, in part only. *Aspidium hirsutulum, Wall. Cat.* 7083, type sheet example b, has no auricles on the stipe, and appears to me to belong to the type (*ciliata*), it only consists of a single small frond, and might belong to either, but I fear *falciloba* and *ciliata* are not well defined as varieties, but run one into the other. *Clarke, F. N. I.* 515, *excl. t.* 105, *Bedd. F. S. I.*

Khasya and Sikkim mountains up to 3,000 feet and river banks in the plains.

(Clarke's variety *pubera* does not belong here, *Wall. Cat.* 338, being *Nephrodium arbuscula* (typical), and from the Sirumalays near Dindigal, not from Nepal.

11. *LASTREA VISCOSA*. (*J. Sm.*) Stipes 6 inches or more long, firm, erect, reddish-brown, slightly scaly, finely villous; fronds 9-12 inches long, 4-5 inches broad, oblong-lanceolate, pinnæ close 2-3 inches long, the lower ones somewhat reduced and deflexed, cut nearly to the rachis into close spreading linear-oblong lobes under 1 line broad, textures herbaceous, colour dark green; rachis villous, under surface densely glandular, veinlets 5-6 on a side, sori medial. *Hook. Syn. Fil. p. 264. Bedd. F. B. I. t. 334.*

Malay Peninsula, at Malacca.

(Also in the Philippines and Borneo.)

12. *LASTREA CRASSIFOLIA*. (*Bl.*) Stipes tufted 1½ foot long, channelled above slightly hirsute; fronds deltoid ovate, firm herbaceous, pinnate, pinnæ pinnatifid, rachis, costa and costules beneath hirsute, the rest glabrous, pinnæ petiolate (petioles ¼ inch long) subsessile, lower ones not reduced in size, pinnatifid nearly two-thirds down to the rachis, segments obtuse subfalcate subentire, basal one reduced in size, costa deeply channelled above, veinlets simple, the lower pair free (as in *Lastrea*) or joining just before entering the sinus of the segments (as in *Nephrodium*) remaining ones simple (or rarely forked) sori one on, or a little below the centre of each vein involucre reniform glabrous. *Aspidium, Bl. En. Fil. Jav. 158. Hook. Syn. Fil. 266. Lastrea nephrodioides, Bedd. F. B. I. t. 199.*

Birma and the Malay Peninsula. I also have a specimen collected in Sikkim, 3,000 feet elevation, by the late Mr. Atkinson which I refer here, as it exactly corresponds in venation; the pinnæ however, are subsessile, and the upper basal segment (or pinnule somewhat enlarged.


(Also in the Philippines and Java.)

13. *LASTREA CANA*. (*Baker.*) Stipes densely tufted, 6-10 inches long, finely villous and often furnished with pellucid scales, chiefly towards the base; fronds 8-12 inches long, elongate lanceolate, not narrowed at the base, or suddenly or gradually narrowed into auricles on the stipe, pinnate with numerous pinnæ, central ones two inches long by ½ inch broad, cut down nearly to the rachis into close

renated lobes, texture herbaceous, rachis slender, villous, and both sides slightly so, veinlets 6–8 on a side, simple (very rarely forked); sori small medial on the veins, indusium reniform, glabrous or pilose. *Baker, Hook. Syn. Fil.* 267, not *Wallich, Bedd. F. B. I. t.* 307. *L. Bergiana, Bedd. F. B. I. Suppl. t.* 370. *Aspid. appendiculatum, Wall. Cat.* 349, type sheet only. *Aspid. eberneum, Wall. Cat.* 389, type sheet only.

Himalayas near Simla, Sikkim, Yakla 8,000 feet elevation. Mr. Clarke thinks this may be only a variety of *ochthodes*, but it has no glands at the base of the pinnæ, the texture also is different, and it is more hairy.

14. *LASTREA BEDDOMEI. (Baker).* Rhizome slender, wide creeping; stipes 6–9 inches long, slender, glossy; fronds pinnate up to about 1 foot long, lanceolate elliptic, central pinnæ the largest, about $1\frac{1}{2}$ inches long, $\frac{3}{8}$ inch broad, cut down almost to the rachis into close rather acute entire segments under 1 line broad, lower pinnæ distant and dwindling down very gradually, texture rather firm, veins beneath furnished sparingly with longish white needle-like hairs, veinlets 4–5 on each side, simple sori near the margin. *Baker, Hook. Syn. Fil. p.* 267. *L. gracilescens, Bedd. F. S. I. t.* 110.



Nº 122.
LASTREA BEDDOMEI.
(*Baker.*)

South India on the Travancore and Tinnevely Mountains in swampy ground. Ceylon, common in swampy places about Newera Elya. A pretty delicate species much cultivated at Ootacamund.

15. *LASTREA ELWESII. (Baker.)* Stipe not seen, main rachis minutely pubescent; fronds oblong-lanceolate, narrowed from the midrib at both ends, 16 inches long, by $4\frac{1}{2}$ inches broad, glabrous, pinnate; pinnæ patent, subobtuse, cut down only one-half towards

the midrib ; segments broad, short, obtuse, lower pinnæ much deflexed and gradually shorter ; texture moderately firm, surfaces naked, bright green ; veinlets distant, simple, about 4 on each side ; sori small marginal ; indusium small fugacious. *Hook. Syn. Fil.* 491. *Bedd. Fern. Sup. t.* 376.

Sikkim (Elwes, only once collected, and a single frond without stipe.)

16. *LASTREA OCHTHODES*. (*Kze.*) Caudex erect, stipes tufted, 1-2 feet long, firm, naked or nearly so ; fronds 2-4 feet long, 8-12 inches broad ; pinnæ $\frac{3}{4}$ -1 inch broad, cut down nearly to the rachis into entire or bluntish, often falcate segments 1-1 $\frac{1}{2}$ lines broad, with a prominent gland at the base beneath, the lower ones gradually reduced ; texture subcoriaceous ; rachis and underside more or less villous ; veinlets prominent, simple 10-15 on a side ; sori submarginal. *Aspidium ochthodes*, *Kze. in Linnaea*, xxiv. 282. *Bedd. F. S. I. t.* 106. *Nephrodium prolixum*, *Baker, Syn. Fil.* 268.

Northern India, throughout the Himalayas and Khasya Mountains ; South India and Ceylon, throughout the mountain forests ; Malay Peninsula.

VAR. β *TYLODES*. (*Kze.*) More coriaceous, rachis and underside glabrous, lower pinnæ not gradually reduced, but suddenly abortive and reduced to tuberculated glands, sori at the base of the veins, close to the costules. *Bedd. F. S. I. t.* 107.

South India mountains, rare, and probably the same localities as *ochthodes* ; it is, I believe, quite entitled to rank as a species. I have had both in cultivation for many years, and Mr. Thwaites, who cultivated them in Ceylon, considered them distinct species.

17. *LASTREA FAIRBANKII*. (*Bedd.*) Caudex long creeping, 2 thick as a crow quill, very black, glabrous, but furnished with a few scales, copiously rooting, stipes 8-20 inches long stramineous, black at the base, rachis channelled above, glabrous or slightly pubescent with weak glandular hairs ; fronds coriaceous 6-10 inches or more long, narrow oblong pinnated, pinnæ opposite or alternate, sessile or nearly so, 1-1 $\frac{1}{2}$ inches long, less than $\frac{1}{2}$ inch broad, linear obtuse or acuminate at the apex, pinnatifid nearly to the rachis, the lower ones

reduced in size and often more distant, the partial rachis channelled and glabrous on the upper side, slightly pubescent or glabrous beneath, and there furnished with numerous broad-ovate or lunate very transparent scales, segments oblong to ovate entire much recurved at the margins; veins more or less undulate; veinlets simple excurrent at the margin; sori one on the centre of each veinlet; involucre reniform, fimbriate at the margin. *Bedd. F. B. I. t. 254.*

Pulney Mountains, in swamps about the south base of Peermall Malley, 4,000 feet elevation. It was also formerly found about the lake at Kodiekarnal on the same mountains, but has disappeared from that locality. In my Supplement I have reduced it to *thelypteris*, but now believe it to be distinct. In its reduced lower pinnæ it approaches *Beddomsi*, but it is much more rigid in texture, and its scaly secondary rachis is quite unlike that fern. I hope it may be further collected. There is a specimen in the British Museum.


B. Veins forked.

18. *LASTREA THELYPTERIS*. (*Desv.*) Rhizome slender, wide-creeping, stipes about 1 foot long, slender stramineous; fronds 1-2 feet long, 4-6 inches broad, lanceolate, truncate at the base; pinnæ spreading, $\frac{1}{4}$ inch broad, cut down very nearly to the rachis into entire spreading linear oblong lobes, those of the barren frond the broadest, lower pinnæ equalling the others, veins forked (upper ones simple) texture herbaceous, rachis and both sides naked, sori small, not confluent, in rows near the recurved edge; indusium small inconspicuous. *Desv. in Mém. Soc. Linn. 6. 257. Bedd. F. B. I. t. 44. Hook. Syn. Fil. 271.*

South India, on the Nilgiris, swamps near Ootacamund; North India, Kashmir, Bandipoor, City Lake, 5,600 feet elevation, Kunawur, 6,000 feet.

(Also in Europe, North Asia, North America, Cape Colony and New Zealand.)


19. *LASTREA GRISEA*. (*Baker.*) Stipes firm, erect, grey, villous upwards; fronds $1\frac{1}{2}$ -2 feet long, 8-9 inches broad, oblong-lanceolate;



mostly forked; sori small, submarginal; involucre ciliated.
Syn. Fil. p. 271. Bedd. F. B. I. t. 335.

South India, near Cochin, only once collected.

20. *LASTREA FUSCIPES*. (*Wall.*) Stipes tufted, slender 6-12
 inches long, dark purplish-brown, scaly below, fronds 12-24 inches
 6-12 inches broad, oblong-lanceolate, pinnate; pinnæ 3-6
 inches long, 1 inch broad, the lowest pair rather shorter than the
 others, and deflexed, and with some of its lower segments often
 notched; segments cut down to a broadly-winged rachis, linear-
 oblong, blunt, entire or slightly crenate; texture herbaceous,
 shiny polished like the stipe; veinlets 6-8 on each side, simple



LASTREA SYRMATICA.

forked, generally excurrent when sterile, but terminating well
 in the margin when fertile; sori generally apical on a veinlet,
 sometimes well below the apex; indusium reniform fugacious.
V. Cat. 361. Lastrea melanopus, Hook. Sp. Fil. iv. 110.
L. F. B. I. t. 38. L. sagenioides, Hook. Syn. Fil. 271. Poly-
um obscurum, Hook. Sp. Fil. iv. 237. Syn. Fil. 308.
L. F. B. I. t. 230.

Birma and the Malay Peninsula.

(Also in the Philippine Islands.)

The type sheet of Aspidium fuscipes in Wallich's Herbarium, No. 1, is this plant, but the other sheets are Pleocnemia membranifolia.

21. *LASTREA SYRMATICA*. (*Willd. under Aspidium*). Stipes

...
 ; abrous, scarcely coriaceous in tex
 sides; veins all forked 10-15 on eac
 on each side of the costule, nearer
 always medial on the superior vein




No 125
 LASTREA FLACCIDA (Hb. K.)

reniform
 237. *L.*
 bilis, *H.*
S. I. t.

Mr.
 veins of
 seen in
 rarely th
 each oth
 margin; t
 tinued be
 found a t
 often app
 held up t

South
 ghat, 2,00
 Travancor



Nº126.

LASTREA BRUNONIANA. (Wall.)

... rachis costules and
brown hairs above, and beneath w
veins pinnate in the ultimate seg
forked or even simple in smaller e
or near the apex of a veinlet; indus
Fil. p. 274. Bedd. F. S. I. t. 250.

*Mr. Clarke has joined this with
in my opinion, its real affinity is with
resembles, and of which it may be a m*

South India, Travancore Hills,
Elya and the highest part of centra
6,000 feet.

(Also in Java.)

23. *LASTREA BRUNONIANA.* (*W*
long, black, densely clothed with large
fronds 12-18 inches long, 2-4 inches
subequal oblong-lanceolate blunt pinn
largest $1\frac{1}{2}$ -2 inches long, $\frac{3}{4}$ inch b
rachis into sharply toothed rounded lo
herbaceous, rachises ebeneous and m
fibrillose scales, under surfaces naked
veinlets. *Asp. Brunonian*



Nº 127
LASTREA BARBIGERA. (*Hook.*)



Nº 128.
LASTREA ODONTOLOMA. (*Moore.*)

the pinnules.
F. B. I. t. 227. L. Falconeri, *Hack*
t. 41.

Himalayas; Kashmir to Sikkim
Nearly allied to *Brunoniana* if it be re

25. *LASTREA ODONTOLOMA*. (L
thick chestnut-coloured, with scattered
fronds up to 10 inches long, by 6–8 in
truncate at the base; pinnæ often w
secondary pinnæ elliptic-oblong, pinn
ments rounded, sharply serrate; te
towards the margin; venation subfl
Clarke, F. N. I. p. 521. L. *Filix-mas*
Fil. p. 498. *Bedd. F. Sup. t.* 373, *not*
Filix-mas.)

Himalayas, Chumba to Bhotan, 1

I have followed Mr. Clarke in s
but I fear I may be wrong in so doing,
Filix-mas, var. *panda* seem to connect
but if it goes into *Filix-mas*, *Brunonian*
referred there. I find the greatest diffi
this species and *Brunoniana*, the fimbri

pinnæ lanceolate, cut down nearly (scarcely ever quite) to the rachis into close, blunt, regular segments which are subentire, crenate, or serrulate, lower pinnæ sometimes gradually reduced, sometimes not so; texture generally more or less coriaceous, rachis scaly or scabrous, under surface generally naked, sometimes fibrillose, veins pinnate, or the lower ones pinnate, sori medial, involucre large, reniform or orbiculate. Polyp. Filix-mas, *L. Sp. Pl.* 1551. *Hook. Syn. Fil.* 272; *Sp. Fil.* iv. 116. *Lastrea* *patentissima*. *Bedd. F. S. I. t.* 11 (typical Filix-mas). *Clarke, F. N. I.* p. 519.


Throughout the Indian region, but generally confined to considerable elevations on the mountains.

(Also found throughout the world in temperate places) except in Australia and America south of Peru).

VAR. β PARALLELOGRAMMA. (*Hook. Fil.* iv. 116.) Pinnate, rarely bipinnate, pinnæ often reduced towards the base, segments oblong-parallelogram, generally close and compressed, entire, crenated or serrated.

I include under this as sub-varieties, *patentissima*, *fibrillosa*, *apiciflora*, *nidus*, *Clarkei*.

Taking *patentissima* (common throughout the Indian region) as the type, they all show very slight differences, but these are not constant; *apiciflora*, which has generally been made a distinct species, is remarkable on account of the sori being confined to the apex of its segments; Mr. Clarke has, however, found it frequently with the sori scattered, though he says it can be distinguished by being remarkably free from fibrillæ and hairs, but I have specimens which I gathered, near Darjeeling, with fibrillæ on all the veins beneath; the fronds are generally more truncate at the base than *patentissima*,



Nº 129.

LASTREA FILIX-MAS. (*L.*)

VAR. NIDUS. (*Clarke.*)

... pinnæ a little reduced and scattered, but with a tendency to be feet. *Bedd. F. B. I. Sup. t. 372.*—tapering much at the base, almost smaller than the type, but quite to 11,000 feet. *Bedd. Fern Sup. t.* from Dalhousie with the under surface (British Museum).—*Fibrillosa* differ surfaces of the pinnæ copiously clothed with hairs, 9,000–12,000 feet from Kumaon

VAR. γ ELONGATA. (*Hook and Greville*) pinnate at the base, bi-tripinnate, or sometimes from high elevations, ultimate segment from a broad base, rachises glabrous or tomentose. *Bedd. F. S. I. 112.* Var. *marginata* from Canariense, *A. Brown.*


South India on the Western mountains; Ceylon; Himalayas and Khasi. *Aspidium rigidum* (*Desv.*) seems quite distinct from *Schimperianum* (*Hochst.*) [intermedia, a form at a higher elevation. 2-pinnate, large sori.]

rachises glabrous, or sometimes scaly. *Don. Prod. Fl. Nep.* 6
Bedd. F. S. I. t. 115. *Arthobotrys macrocarpa*, *Wall. Cat.* 395.

North India up to 4,000 feet elevation; South India, Western mountains 2,000–4,000 feet; Malay Peninsula. A very distinct looking plant in its typical form at low elevations, but running into *elongata* at higher elevations (*vide* forms of my collecting on the Nilgiris and Brumagherries in the British Museum).

VAR. ϵ PANDA. (*Clarke, F. N. I. p.* 519, *t.* 68, *fig.* 1.)
 Fronds simply pinnate, truncate at the base, narrow oblong, pinnae pinnatifid $\frac{1}{2}$ – $\frac{2}{3}$ towards the midrib, segments sub-spinulose.

Himalayas, Dhurmsala, 10,000–11,000 feet elevation. This has much the aspect of *odontoloma*, and it will probably prove to be a luxuriant form of that plant; if a variety of *Filix-mas*, it certainly connects *odontoloma* with it.



Nº130.

LASTREA FILIX-MAS. (L.)

VAR. δ COCHLEATA. (Don.)

† † † *Fronds compound, 2–4-pinnate.*


27. LASTREA RIGIDA. (*Desv.*) Stipes tufted, 6 inches long, densely clothed below with large lanceolate or ovate scales; fronds up to 2 feet long, oblong-lanceolate, largest pinnae 3–4½ inches long, 1–1½ inch broad, the pinnules of the lower half free ovate rhomboidal, cut down nearly to the rachis below, teeth mucronate; texture herbaceous; rachises usually scaly, under side naked or slightly glandular; veinlets subpinnate in the lower lobes of the pinnules; sori near the midrib; indusium firm prominent. *Hook. Syn. Fil. p.* 275. *Clarke, F. N. I. p.* 523.

Very near *L. Filix-mas elongata*, and I cannot say I know any character to distinguish it.

deltoid, 3-4 inches long, 1½-2 inches
equal to the next; pinnules ovate-lance
long, ½ inch broad, cut down to the ra
lobes with copious aristate teeth; text
mineous, scarcely scaly; under surface
green; involucre not gland-ciliated. *H*

VAR. γ REMOTA. (*A. Br.*) Scal
extending to the rachis; fronds oblong-la
6 inches broad; pinnæ lanceolate, clo
only the lowest free, the largest about
cut half-way down to the rachis or more,
side and involucre not glandular. *Hoo.*
F. B. I. t. 336. *Asp. eberneum*, *Wall*
Kashmir to Nepal, 6,000-9,000 feet
(Also in Northern and Alpine Europ

29. *LASTREA SPARSA.* (*Don.*)
densely covered with lanceolate-linear
stipes long, 1 foot or more, with scatt
generally glabrous or subglabrous; fronds
pinnæ the largest, deltoid with . . .



SCIENTIA

LASTREA SPARSA. (Don.) VAR. γ *OBTUSISSIMA.* (Mett.)

...pinnæ less divided, pinnul
Bedd. F. Sup. t. 374.

Nepal to Bhotan, 9,000-12,000

VAR. γ OBTUSISSIMA. (*Met*
light brown broad chaffy scales, low
frond less cut, the ultimate segme
oblong to obovate. *Bedd. F. Sup.*
Ceylon, Nepal.

VAR. δ DELTOIDEA. (*Bedd.*)
and lower basal pinnules being
/ *F. S. I. t. 248.*
Ceylon.

VAR. ϵ MINOR. (*Thw.*) Frond
simply pinnate, the pinnæ only $\frac{1}{2}$ i
larger specimens, however, running
deltoidea.

Ceylon ; Simla, North Cachar.

VAR. ζ ZEYLANICA. (*Bedd.*) I
sissima, but much more compound
than the others. main second


on (near Newera Elya), 5,000 feet elevation. *Lastrea undulata* (Walters), *Bedd. F. S. I. t.* 271.

Mr. Clarke's variety *latesquama*, does not differ from what I consider the type of *sparsa*, which agrees exactly with *Wall. Cat.* 390, *densum*, and *Wall. Cat.* 7080, *Asp. oppositum*.

30. *LASTREA PULVINULIFERA*. (*Bedd.*) Stipe up to 30 inches furnished with a dense cushion-like mass of golden scales at base (as in *L. odorata*), and clothed throughout (especially in the younger specimens), as are the rachis and partial rachises, with numerous brown black scales, which proceed from a broad base, and which are usually more copious at the axils of the pinnæ; fronds ample, up to 24 inches long (without the stipe) by 18 inches broad near the base, rigid, tripinnate, coriaceous-herbaceous; pinnæ gradually decreasing in size from the base to the apex, lower pair opposite, large deltoid, the pinnules of the lower margin much the largest, remaining alternate or rarely subopposite or opposite; pinnules up to 4 inches long in the basal pinnæ, with their secondary pinnules equal to the primary pinnules on the upper parts of the frond; segments usually rounded, entire or lobed, and occasionally dentate or crenate, serrate on both sides, but sometimes furnished with a few weak teeth on the costules and veins on the upper side; veinlets inconspicuous, terminating within the margin; sori terminal on the lower margin; involucre reniform glabrous. *Bedd. F. B. I. t.* 333. *Lastrea pulvinulifera* (Baker), *Hook. Syn. Fil. p.* 500, *in part only*. *Clarke, F. N. I. p.* 525, *in part only*. *Lastrea sparsa*, var. *squamulosa*, *Clarke, l. c. p.* 524, *in part*.

Sikkim, very common about Darjeeling. Perhaps only a very robust form of *sparsa*.

31. *LASTREA BUCHANANI*. (*Baker.*) Stipes tufted up to 1 foot long, dark coloured at the base, straw-coloured upwards, densely clothed as is the rachis, with long spreading rather rigid black hairs; blade ovate to deltoid, tripinnate, with the tertiary pinnules from entire to pinnatifid, secondary rachis hairy, lowest pinnæ largest and unequal sided from being more compound at the base,



tertiary rachis a little scaly, ultimate segments oblong blunt; texture subcoriaceous; veins pinnate or forked in the ultimate segments; sori below the middle of the veins and near the costules (never apical on a veinlet as in *pulvinulifera*); involucre reniform, glabrous, persistent. *Hook. Syn. Fil. p.* 498. *Lastrea pulvinulifera*, *Hook. l. c.* 500, *in part only*. *Nephrodium pulvinulifera*, *Clarke, F. N. I. p.* 525, *in part only*.

Himalayas, near Darjeeling, 7,500 feet elevation.

(Also in Natal and Bourbon.)


32. *LASTREA SPECTABILIS*. (*Wall. under Aspidium.*) Stipes 1-1½ foot long and with the rachises purple-coloured and furnished with patent linear subulate reddish-brown scales; fronds 2-4 feet long, more than 1 foot broad, subdeltoid, tripinnate with the tertiary pinnules deeply pinnatifid, and their segments again crenated or serrated, lowest pinnæ largest, deltoid, unequal-sided, the two lower basal secondary pinnæ being much produced; texture moderately firm, the costa of the pinnules and veins of the segments sparingly furnished with long weak setæ most copious on the upper side, otherwise glabrous, sori medial on the upper basal veinlet of the segment; involucre reniform. *Wall. Cat.* 372. *Clarke, F. N. I. p.* 526. *Lastrea Hendersoni*, *Bedd. Fern Sup. t.* 377.

Khasya on Shillong Hill 5,000-6,500 feet elevation; Nepal.

33. *LASTREA DEPARIOIDES*. (*Moore under Diclisodon.*) Stipes about 1 foot, long, firm, slightly scaly below; fronds up to 2 feet long, oblong-lanceolate, bipinnate with the pinnules deeply pinnatifid, pinnæ 4-6 inches long, 2 inches broad, lanceolate, the secondary rachis with a very narrow inconspicuous wing, pinnules stalked, deltoid, unequal sided, with broad blunt subquadrangular lower segments and prominent teeth, texture rather firm, rachis and both surfaces naked; sori terminal in the teeth which the convex involucre quite conceal. *Diclisodon*, *Moore, Index Fil.* 316. *Lastrea deparioides*, *Bedd. F. S. I. t.* 104. *Hook. Syn. Fil.* 277.

South India, on the Anamallay Hills, 2,000 feet elevation; Ceylon, southern provinces.

normal form, of it, as t
 genus. *H. k. Syn. L. l. p. 277.*
E. S. I. t. 246.



Nº133
 LASTREA DEPARIoides.
 (Moore.)

colour pale green - red

(Also in Tropical Africa ; Mauritius ; South China.)

36. *LASTREA PARISHII*. (*Hook.*) Caudex creeping, stipe 6-9 inches long, slender, naked, ebeneous, fronds small, 6-8 inches each way, deltoid, the lower pinnæ much the largest, lower pinnules larger than the others, which are oblong-lanceolate acuminate $1\frac{1}{2}$ -2 inches long by $\frac{1}{4}$ - $\frac{1}{2}$ inch broad, cut down nearly to the rachis into close oblong-lanceolate crenated lobes, texture membranaceous, rachises and under surfaces pubescent ; veinlets pinnate or forked in the lobes ; sori in two rows, not far from the midrib, apical on a veinlet or situated well below the apex, indusium reniform. *Hook. Syn. Fil.* p. 281. *Bedd. F. B. I.* t. 43.

Birma, Moulmein, moist shady places on limestone rocks.

37. *LASTREA SIKKIMENSIS*. (*Bedd. under Polystichum.*) Stipes and rachises rather copiously furnished with persistent ovate acute blackish scales, fronds ovate acuminate up to 3 feet long, subcoriaceous-membranaceous, bipinnate, primary pinnæ 4-6 inches long, oblong acuminate, secondary pinnules oblong, deeply pinnatifid, $\frac{1}{2}$ inch to 1 inch long, tertiary pinnules sharply and deeply cut, with the segments acuminate and falcate, costa of the pinnules waved, veins simple or forked, terminating within the margin, sori generally one to each segment, terminal on a veinlet, involucre reniform. *Polystichum sikkimense*, *Bedd. F. B. I.* t. 127. *Nephrodium*, *Clarke, F. N. I.* p. 525. *Hook. Syn. Fil.* 256.

Sikkim, Mon Lepcha, 10,000-12,000 feet elevation.

38. *LASTREA BLUMEI*. (*Hook.*) Stipes up to about 2 feet long, stout, very shaggy, with long linear rufous scales, rachises



NO 134.

LASTREA PARISHII. (*Hook.*)

rufous-pubescent, and copiously furnished with rufous scales, fronds very large ovate in outline, tripinnate, primary pinnæ 10–15 inches long, tertiary pinnæ 1–1½ inch long, pinnatifid nearly to the rachis, the segments oblong entire or rarely rather deeply incised, sparingly hairy above, pubescent on the costa on both sides; veinlets forked or simple; sori small medial on the veinlets; involucre reniform, glabrous. *Lastrea Blumei*, *Hook. Sp. Fil.* iv., 135, *in part at least*. *Bedd. F. S. I. t.* 249. *L. intermedia* (*Baker*), *Hook. Syn. Fil.* p. 283. *L. rhodolepis*, *Clarke, F. N. I.* 526.

North India, Sikkim, Assam and Khasya 5,000–7,000 feet; Ceylon, common in the central provinces, 2,000–3,000 feet elevation. (Also in Japan; Malay Islands; Polynesia.)

Mr. Clarke states that it is not the *intermedia* of Blume. I do not know whether he is correct in this, but in any case a new name is unnecessary, as it has been published and figured as *Blumei*, a name otherwise unoccupied.

39. *LASTREA RECEDENS.* (*J. Smith.*) Rhizome creeping, stipes erect, villous, clothed at the base with linear scales; fronds very large deltoid, 3-pinnate above, 4-pinnate below, with the quaternary pinnules deeply pinnatifid, lower pinnæ up to nearly 2 feet long, (in large specimens) secondary pinnæ on the lower side 9–10 inches long, quaternary pinnæ ¾ inch long, the ultimate segments entire or slightly crenated, texture herbaceous, rachises villous, under surface softly hairy on the veins, upper surface nearly glabrous except on the costules, veins forked, sori terminal on, or a little below, the apex of a veinlet. *Phegopteris recedens* (*J. Smith*) *Hook. Jour. of Bot.* iii. 394. *Lastrea recedens*, *Bedd. F. S. I. t.* 98 *two-thirds of a small frond.* *Nephrodium*, *Hook. Syn. Fil.* 281.

South India, on the lower Pulney Mountains, and Travancor and Tinnevely Hills, 3,000–5,000 feet elevation; Ceylon, central provinces, 2,500–5,000 feet elevation.

(Also in the Philippines.)

40. *LASTREA DISSECTA.* (*Forst.*) Rhizome erect, stipes 1 foot or more long, a little scaly at the base, fronds deltoid to ovate, up to

nearly 3 feet long ; texture herbaceous, pinnate above, with the pinnæ 6-9 inches long by $1\frac{1}{4}$ inch broad, pinnatifid two-thirds towards the rachis, bipinnate below, from some of the lower basal pinnules of the lowest pinnæ being stalked and much elongated and deeply pinnatifid, and equal to some of the upper primary pinnæ ; ultimate segments crenulate, a little hairy above (on the lamina, not the veins), with scale-like hairs, quite glabrous below in South Indian examples, but with scale-like hairs in specimens from Ceylon ; veinlets forked, always free, a forked veinlet often proceeds from the costa to the margin of the sinus, between the segments ; sori apical on the superior veinlet, forming a single row on each side of the segments nearer the margin than the midrib, involucre reniform, glabrous ; fertile portions of the frond often much contracted and frond often only partially seeding. *Lastrea membranifolia*, *Bedd. F. S. I. t.* 102, *a small frond*.

The descriptions of *Lastrea dissecta* in *Hooker's Syn. Fil.* and of *L. membranifolia* in *Hook. Sp. Fil.* includes this species, the variety *ingens* and *Pleocnemia membranifolia*.

Madras Presidency, Western mountains, common up to about 5,000 feet elevation ; also in the N. Arcot and Vizagapatam Hills ; Ceylon, up to 5,000 feet ; Birma.

(Also in Malay Islands and Polynesia.)

VAR. INGENS. (*Clarke.*) A very large fern, stipe and main rachis pubescent, and with lax narrow-lanceolate sparse brown deciduous scales ; fronds up to 9 feet long, the lower pinnæ 2 feet and more long, bipinnate above, the pinnæ about 1 foot long by 6 inches broad, the secondary pinnæ quite distinct but sessile from a very broad base and deeply pinnatifid, tripinnate below, the lower basal pinnules of the lowest pinnæ being stalked, elongated and again pinnated, their pinnules (tertiary pinnæ) being deeply pinnatifid, and equalling the secondary pinnæ above ; ultimate segments entire, nearly glabrous above except on the costa, and costules, but there are a few scale-like hairs on the lamina, costa, costules and often the veins beneath furnished with white glistening multicellular patent hairs ; texture, venation and sori exactly as in *dissecta* ; involucre reniform,

but the lobes of the sinus often overlapping, so that it appears to be circular; fertile portions of the frond often much contracted, and fronds often only partially seeding. *Clarke, F. N. I. p. 526.*

Himalayas, Sikkim, and Bhotan, 4,000–7,000 feet, very abundant between Darjeeling and Surail; Khasya, 3,000–5,000 feet.

Mr. Clarke says that it seeds but rarely, and then usually only very partially, this is an error, as I found many clumps near Darjeeling (end of November) with most of the fronds entirely covered with seed, and not in the least contracted. I had written out a full description, believing this to be a distinct species from *dissecta*, differing in having copious multicellular hairs beneath, the South Indian plant being glabrous; I have, however, since seen Ceylon specimens in the British Museum which have similar hairs beneath, so I can now only regard it as a variety; it only differs in its larger size, and in being more compound, and in the ultimate segments being more entire.

41. *LASTREA SPLENDENS.* (*Hook.*) Stipes 2–3 feet long, stout, polished bright chestnut coloured and scaly, as is the main rachis; fronds long-lanceolate up to 4 feet, by about $1\frac{1}{2}$ broad, bipinnate, primary pinnæ narrow lanceolate, about 2 inches broad, and nearly the same breadth throughout; secondary pinnæ 20–30 pairs, about $\frac{1}{4}$ inch broad, lanceolate-oblong, rounded at the apex, cut about one-third towards the rachis into small lobes, the upper basal one being generally rather larger than the others, all more or less serrated; texture subcoriaceous, glabrous on both surfaces, or slightly scaly beneath; rachis of primary pinnæ deeply channelled above, and winged upwards glabrous above, often scaly beneath; veinlets pinnate or forked, not reaching the margins; sori well below the middle of upper veinlet, forming a row on each side of the pinnule near the costa. *Hook. Syn. Fil. 282. Bedd. F. B. I. t. 42.*

Sikkim and Bhotan, 6,000–7,000 feet elevation; Malay Peninsula.

42. *LASTREA ANGUSTIFRONS.* (*Moore.*) Rhizome wide-creeping; stipes 1–2 feet long, ebeneous, deciduously scaly; fronds 2 feet

6-8 inches, glabrous, elongate, 3-pinnate, pinnæ distant, erect-tent, flexuose, lanceolate, two inches broad; pinnules distant lanceolate, cut down to the rachis below into subquadrangular toothed segments (tertiary pinnæ) $\frac{1}{2}$ inch broad, texture subcoriaceous; rachis and both sides naked; veinlets forked; sori small, low down on the veins near the midrib of the pinnules. *Moore in Hook. Bot. Fil. iv. p. 126. Hook. Syn. Fil. p. 283. Bcdd. F. B. I. t. 226. splendens, var. β , Hook. Sp. Fil. 126.*

Nepal, Sikkim (a little-known plant, perhaps abnormal *Filix-mas* or *splendens*).

43. *LASTREA SCABROSA* (Kunze.) Caudex stout, suboblique, densely leucous, stipes 1-3 feet long, densely palcaceous towards the base, scabrous above; fronds membranaceous, semitransparent, deltoid-ovate acuminate up about 2 feet long, bipinnate, or subtripinnate above, tripinnate below, pinnæ alternate, distant 6-inches long, the lowest pinnule deltoid with some of the inferior basal secondary pinnules enlarged to double the size of the others, which are generally $1\frac{1}{2}$ -2 inches



LASTREA SCABROSA. (Kunze.)

long and pinnatifid down to a winged rachis (or almost pinnate); the pinnules $\frac{1}{4}$ - $\frac{1}{2}$ inch long, cut down one-third to half-way to the costa into more or less obtuse segments, veinlets prominent, one on each segment, simple or forked, terminating within the margin and clavate at the apex; costa and veins furnished with weak whitish

hairs on both sides ; sori medial on the veinlets, one to each segment, becoming jet black when ripe, indusium (?) *Aspidium scabrosum* (*Kunze*), *Hook. Syn. Fil.* p. 283. *Polypodium nigrocarpum*, *Bedd. F. S. I. t.* 169.

Nilgiris, common in the sholas at the higher elevations, 6000-7,500 feet ; in texture and habit the pinnæ much resemble those of *Cystopteris setosa*. I have never been able to detect any indusium even in the youngest stage, though I have long had it in cultivation ; I suspect it is really a *Phegopteris*.

44. *LASTREA FERRUGINEA*. (*Bedd.*) Caudex short, stout, erect, stipes 1-1½ foot long, densely clothed throughout with broad-ovate very transparent scales and scabrous with tubercles ; fronds 1-2 feet long, deltoid-ovate, tripinnate below, bipinnate tripinnatifid above, texture herbaceous, turning blackish when dried, pinnæ 8-10 inches long, spreading at a right angle, the lowest pair deltoid and larger than the others and more compound, its lower basal secondary pinnæ being elongated and pinnate, bipinnatifid ; secondary pinnæ 1-2 inches long, pinnate in the lower and larger pinnæ, pinnatifid above, pinnules about ½ inch long, cut about half down into oblong rounded crenate or subentire segments ; rachises and both sides ferruginous, with minute glandular scale-like pubescence, veinlets simple or forked, terminating within the margin and clavate at the apex, sori medial or subterminal on the veins, submarginal, one to each lobe of the tertiary pinnæ, in the South Indian plant confined to the upper lobes, in the Ceylon plant more lobes are seeded and the sori are not quite so marginal, indusium reniform, glabrous, persistent. *Bedd. F. S. I. t.* 100. *Hook. Syn. Fil.* 283. *Lastrea obtusiloba* (*Baker*), *Hook. Syn. Fil.* 284. *Bedd. F. S. I. t.* 296. *Asp. Blumei*, var. ? *Thwaites*, p. 392.

South India, Nilgiris, rare in sholas between Avalanche and Sispara, 7,500 feet elevation, not observed elsewhere ; Ceylon (*C. P.* 3142), about Newera Elya, 6,000 feet elevation. The Ceylon and South Indian plants are certainly the same ; I had both in cultivation for a long time and could not have distinguished them except for the labels ; it is nearly allied to *scabrosa*, but quite a different colour both living and dried, more ferruginous, and not scabrous.



LASTREA FERRUGINEA. (Bedd.)

45. *LASTREA BORYANA*. (*Willd. under Aspidium.*) Stipes : feet long, sparingly scaly near the base ; fronds very large, up to 8 feet long by 4 feet broad, herbaceous or almost membranaceous, bipinnate with the secondary pinnæ ($3\frac{1}{2}$ –6 inches long) pinnatifid down to a very regularly but narrowly winged midrib, pinnules blunt $\frac{1}{2}$ – $\frac{3}{4}$ inch long, more or less wide apart at the base, pinnatifid about one-third or half down into small entire segments ; rachises and both surfaces naked or with a little mealy pubescence and a few hairs ; veins forked simple ; sori low down on the veins, in a single row on each side of the pinnules near the midrib, generally very small, large in some Himalayan examples, involucre often much lacerated on the margin, very fugacious. *Willd. Sp. Pl.* v. p. 285. *Baker, Syn. Fil.* 2. *Lastrea divisa*, *Hook. Sp. Fil.* iv. 133. *Wall. Cat.* 393. *Bedd. S. I. t.* 97. *Polypodium subtripinnatum*, *Clarke, F. N. I.* p. 54. 80, fig. 1.

South India, on the Western mountains, 3,000–7,500 feet elevation ; Ceylon, central provinces ; Himalayas, Gurwhal, Bhotan, 4,000–8,000 feet elevation ; Khasya ; Malay Peninsula. Clarke's variety *microstegioides* does not recede from the type.

(Also in Bourbon, Java, and China.)

46. *LASTREA TENERICAULIS*. (*Wallich under Polypodium*) Caudex erect, furnished with numerous fibrous roots, stipes tufted below a little scaly, above very glaucous, with a blueish bloom, which easily rubs off ; rachis with a line of hairs on the sulcated upper side, glabrous beneath ; fronds 1–3 feet and more long, broad ovate-lanceolate acuminate, membranaceous, bipinnate with the primary pinnatifid nearly to the costæ, pubescent with long white hairs on the costa and costules on both sides, rachis of the primary pinnæ furnished with a very regular line of dense hairs on the upper side, glabrous and convex below, pinnæ rather remote, opposite or alternate lanceolate acuminate, inferior ones 10–12 inches long by 3 inches broad ; pinnules up to 2 inches long, oblong lanceolate from a broad adnate base (which is decurrent so as to form a winged rachis, particularly in the upper pinnæ), pinnatifid (in the lower pinnæ almost to the costa) ; secondary pinnules pinnatifid in the lower pinnæ, entire

with the apex crenated in the upper portion of the frond; veins in the larger ultimate segments pinnate with veinlets simple, in the smaller segments veinlets once-forked only, all terminating a little within the margin; sori medial or terminal on the veinlets, 1-6 to each ultimate segment, involucre very deciduous. *Polypodium tenericaule*, *Wall. Cat.* 335. *Lastrea flaccida*, *Bedd. F. S. I. t.* 90. *Polyp. Russellianum*, *Wall. Cat.* 7077. *Lastrea setigera*, *Baker, Syn. Fil.* 284, in part.

Hooker's and Baker's descriptions of *tenericaulis* and *setigera* are made to include *Phegopteris ornata*, a very different fern. I have never been able to detect an indusium even in the youngest stage of our South Indian plant, and I should follow Wallich and include it in *Phegopteris*, only Mr. Clarke says it is present in the Bengal examples.

South India, on the Western mountains, 2,000-3,000 feet elevation, very common; Ceylon, 1,500-3,000 feet; North India, Himalays from no great elevation up to 4,000 feet; Malay Peninsula.

(Also in China, Australia, and Polynesia.)


GENUS LIV.—NEPHRODIUM. (*Schott.*)

(From *nephros*, a kidney; shape of indusium.)

Sori subglobose, dorsal on the veins; (often athyrioid in *otaria*) indusium reniform, sometimes wanting; veins pinnate, one or more pair anastomosing angularly with an excurrent veinlet from their junction which is either free or joined in the angle of the next superior pair; fronds always simply pinnate with pinnatifid pinnæ in the Indian species (rarely simple in species not Indian); caudex erect or creeping. Most of the supposed species of *Goniopteris* fall in here, as an indusium is often present in the very young stage though sometimes entirely absent.

1. *NEPHRODIUM OTARIA*. (*Kze.*) Rhizome creeping, stipes 6-12 inches long, pale-coloured; fronds 12-15 inches long, pinnate, lateral

pinnæ distant, 4-6 inches long, $1\frac{1}{2}$ inch broad, linear-oblong, truncate at the base, caudate at the apex, coarsely but shallowly lobed, the lobes serrated; terminal pinna sometimes with larger lobes or auricles towards its base; texture thin herbaceous; rachis and surfaces smooth; veins pinnate in each lobe, 6-8 on each side, the lowest pair anastomosing angularly, with an excurrent vein which either runs out to the margin or occasionally joins one or even two of the superior veins (but the anastomosing is not so systematic as in the other species of the genus); sori medial on the veinlets, most



Nº137

NEPHRODIUM OTARIA. (A. n.)

general; Ceylon, 2,000-3,000 feet elevation.

(Also in Luzon.)

of which are soriferous; fertile frond often contracted but not always; indusium reniform or often asplenoid and athyroid. *Aspidium otaria*, Kunze, *Herb. Mett. Aspid. p.* 34. *Hook. Syn. Fil.* 288. *Pleocnemia aristata*, *Hook. Sp. Fil.* iv. 62. *Bold. F. S. I. t.* 83. The sori are quite abnormal, the venation and position of the sori are not at all that of *Pleocnemia*.

South India, Anamallay Hills, the dry teak forests, 1,000-3,000 feet elevation, appearing in the rainy season, Tinnevely Hills and elsewhere on the Western ghats, but by no means

2. *NEPHRODIUM UNITUM*. (*Linn. under Polypodium.*) Rhizome creeping, stipes 1-1 $\frac{1}{2}$ foot long, brownish naked; fronds about 2 feet long, pinnate, pinnæ numerous, 3-5 inches long, $\frac{1}{2}$ inch broad, cut down about one-third, rarely more, into triangular acute entire or subentire lobes, lower pinnæ not reduced; texture coriaceous; veins pinnate in the lobes, 4-8 on each side, the lowest pair anastomose

gularly, with an excurrent veinlet which runs to the margin or joins a rarely two pair of the superior veins; upper surface glabrous, under surface generally very hairy; sori medial on the veins; indurum reniform, hairy. *R. Br. Prod. Fl. Nov. Holl.* 148. *Hook. Sp. Fil.* 289. *Polypodium unitum*, *Linn. Sp. Pl.* 1548, *in part.* *propinquum*, *Hook. Sp. Fil.* iv. 79. *Bedd. F. S. I. t.* 89.

Throughout the Indian region in swampy places and tanks.
(Also in Tropical Asia, Africa, Australia and America.)

3. *NEPHRODIUM PTEROIDES*. (*Retz. under Polypodium.*) Rhizome wide-creeping, stipes up to 2 feet long, slender, slightly scaly below; fronds up to 4 feet or rather more long and 2 feet broad, bipinnate, pinnæ spreading $\frac{3}{4}$ to nearly 1 inch broad, apex acuminate, lower ones not reduced), cut down only one-third, the lobes triangular, entire; texture herbaceous; rachis and upper surface nearly glabrous, under surface with minute glandular pubescence on all the veins, veins pinnate on the lobes, the lowest pair always anastomosing angularly with a free excurrent veinlet, the other veins, (7-9 pair,) rays free; sori towards the apex of the upper veinlets always connected to the lobes or the apex of the lobes. *Polypodium pteroides*, *Retz.*, *Hook. Syn. Fil.* 289. *Nephr. terminans*, *Wall. Cat.* 386. *Hook. Sp. Fil.* iv. 73. *Bedd. F. S. I. t.* 90.

Madras Presidency, Western mountains, 2,000-4,000 feet elevation; Ceylon, up to 3,000 feet; Birma; (not in N. India).

(Also in the Philippines.)

Very near *extensum*, but less cut, the anastomosis of the veins angular and the sori terminal.

4. *NEPHRODIUM EXTENSUM*. (*Bl.*) Rhizome creeping, stipes up to 2 feet, glabrous or slightly pubescent; fronds up to 4 feet long, and 1½ foot broad, bipinnate; pinnæ about $\frac{3}{4}$ inch broad (lower ones not reduced), cut down two-thirds into linear oblong-falcate slightly enulate lobes; texture herbaceous; rachis subglabrous, upper surface naked, or with a few distant hairs, under surface with minute glandular pubescence on the veins; veins pinnate in the lobes, 10-13 pair, the lowest pair anastomosing at an angle with a free excurrent

veinlet, or meeting just at the sinus without the excurrent veinlet, or the lowest pair is quite free (as in *Lastrea*), the other pairs always all free; sori towards the apex of the veins and present on all or all but the 1-2 lower ones, often much immersed and visible as punctiform dots on the upper surface. *Asp. extensum*, *Bl. En. Pl. Fav. Fil.* 156. *Hook. Syn. Fil.* 289, (not *Bedd. F. S. I. t.* 85, which is large molle). *Asp. multijugum*, *Wall. Cat.* 348, (not 355, which is *truncatum*). *Nep. punctatum*, *Bedd. F. B. I. t.* 131, (not *punctatum*, *Baker, Syn. Fil. p.* 513, which is *Lastrea ochthodes*, vide specimen in Kew Herbarium.)

South India, Tinnevely Hills, rare; Ceylon, central provinces, 3,000-4,000 feet elevation; Birma, Penang, and Malay Peninsula (Khasya, 2 sheets of Griffiths in the Kew Herbarium, but not found by any one else in North India).

(Also in the Malay Islands.)

5. *NEPHRODIUM MICROSORUM*. (*Clarke*.) Rhizome wide creeping, stipes angled, furnished with linear-subulate long soft brown persistent scales towards the base, more or less softly pubescent as is the rachis; fronds about 2 feet long; pinnæ rather distant, sessile, about $4\frac{1}{2}$ inches long, by $\frac{7}{8}$ inches broad, the lower ones somewhat reduced or the stipe with auricles, deeply cut almost to the rachis into linear-oblong tongue-like entire or subentire segments, the upper and lower basal ones often considerably longer than the others; texture thinly herbaceous, both surfaces, especially the under one, furnished with scattered long white needle-like hairs; veins about 13 pairs, the lowest pair always regularly anastomosing at an acute angle very near the costa of the pinnæ with an excurrent vein, the other pairs always all free; sori minute, sometimes only near the costa of the pinnæ and scarcely extending to the lobes; involucre glabrous caducous. *N. extensum*, var. *microsorium*, *Clarke, F. N. I. t.* 530, and var. *late-repens*.

Sikkim up to 4,000 feet elevation, very abundant about Siligori, near the foot of the Darjeeling Pass.

6. *NEPHRODIUM CUCULLATUM*. (*Bl.*) Stipes about 1 foot



2. 11. 1919

NEPHRODIUM CUCULLATUM. (Bl.)

long, hairy, strong erect ; frond with the main rachis hairy beneath, up to 2 feet long, pinnate ; pinnæ very close, 3–6 inches long, by $\frac{3}{8}$ – $\frac{1}{2}$ inch broad, cut down scarcely one-third into triangular acute subfalcate lobes, lower pinnæ dwindling suddenly into mere auricles ; texture very coriaceous, lower surface very pubescent ; veins (raised when dry), 8–10 pair, 3–5 lower ones uniting ; sori towards the apex of the veins, which are generally all soriferous ; indusium prominent, glabrous, persistent. *Aspidium cucullatum*, *Bl. En. Pl. Jav. Fil.* 151. *Hook. Syn. Fil.* 290. *N. unitum* (*Sieber*), *Wall. Cat.* 358. *Bedd. F. S. I. t.* 88.

Madras Presidency, Western ghats, not common ; Ceylon, Western and Southern provinces, marshy places, common ; Malay Peninsula ; (Assam doubtful.)

(Also in Mauritius, Malay Islands, and Polynesia.)

7. *NEPHRODIUM ELATUM*. (*Baker*) Rhizome wide-creeping, stipes 1 foot long, downy ; rachis downy ; fronds oblong lanceolate, 3–4 feet long, pinnate, pinnæ sessile, 9–15 inches long by 1–1 $\frac{1}{2}$ inch broad, cut down nearly two-thirds into close rather falcate blunt lobes ; lowest pinnæ distant and much reduced or not reduced at all, texture firm, herbaceous, above rachis of pinnæ strigose, and a few hairs on the veins, beneath costules and veins hairy, with longish white hairs, veins 13–16 pair, the lower 3–4 pair anastomosing with excurrent veinlets, sori large medial, involucre minute fugacious. *Baker, Syn. Fil.* 502. *Bedd. Fern. Sup. t.* 378.

Himalayas (Dr. Jerdon.) Mr. Clarke has stated in his Review that Dr. Jerdon's specimen is a *Goniopteris*, but on examining it with me, he acknowledged that it is unlike any *Goniopteris*, and that it is closely allied to, if not identical with, the Mauritius fern ; the lower pinnæ are not reduced in Jerdon's specimen at Kew, but much so in the Mauritius plant, which is the only difference between them, and this variation occurs as much in others of the genus.

(Also in the Mauritius.)


8. *NEPHRODIUM ARIDUM*. (*Don.*) Stipes 1 foot or more long, erect, slightly pubescent ; fronds 3–4 feet long 12–18 inches broad ;

æ rather distant, 6-9 inches long, $\frac{1}{4}$ -1 inch broad, cut about one-third of the way down into subtriangular sharp pointed lobes; textureaceous; rachis and lower surface hairy; veinlets 8-10 on a side, pair anastomose; sori in rows about midway between the midrib edge; indusium naked, lower pinnae reduced. *Aspid. Don. Prod. Nep. p. 4.* *A. venulosum, Wall. Cat. 352, type sheet.* *A. obscurum, Blume.* *Neph. aridum, Hook. Syn. Fil. p. 291.* *Bedd. F. B. I. t.*

Throughout East Bengal abundant, from the Soonderbun to Assam the Dehra Doon, up to 3,000 feet elevation.

Malay Peninsula (not in South India or Ceylon.)

9. *NEPHRODIUM GLANDULOSUM.* (*Hook.*) Stipes proximate, main rachis densely villous, frond 1 foot long; adpressedly strigose on upper surface; pinnae truncate at the base, subentire, entire or pinnatifid scarcely one-sixth the way to the midrib; veinlets beneath minutely punctate, several pairs uniting, indusium reniform, elongate, prominent, firm; sori ultimately often confluent, *Hook. Fil. iv. 76, partly.* *Clarke, N. I. p. 531, t. 74, and 1.* *Not Bedd. F. B. I. t.*, which is *urophyllum*.



Nº 139.


NEPHRODIUM ARIDUM. (*Don.*)

The above is Mr. Clarke's diagnosis from Blume's specimens collected in Java, but there is an exactly similar specimen in the Kew herbarium from Griffith, supposed to have been collected in Assam, Mr. Clarke thinks it is probably from Malacca, but in either case it must be recorded in this work; it differs from the next in being strigose above, but I strongly suspect it is only a slight variety of

it; it has glandular dots on the under surface, 6-7 pairs of veins, of which the four lower anastomose.

10. NEPHRODIUM UROPHYLLUM. (*Wall. under Polypodium*—)

Rhizome short creeping; stipes approximate, 2 feet or more long, stout, erect, furnished with lanceolate-linear brown scales below; fronds up to about 4 feet long, pinnate; pinnæ narrow-oblong, subsessile or sessile, acuminate or caudate at the apex, 6-12 inches long by $1\frac{1}{2}$ -2 inches broad, the margin subentire, slightly crenated with very short serratures; upper surface quite glabrous, or the



N^o 140.

NEPHRODIUM UROPHYLLUM.

(*Wall*)

rachis of the pinnæ a little strigose, under surface quite glabrous or with some minute pubescence on the veins, copious glandular punctiform dots are present on both sides; texture subcoriaceous or herbaceous; veins 8-10 pair, rarely 14, all or nearly all anastomosing, the current veinlets either free or joining the pair above; small, one to each vein, about its centre; indusium reniform, generally absent. *Wall. Cat.* 199 *Hook. Syn. Fil.* p. 314. *Goni*

opteris, *Bedd. F. S. I. t.* 239. *Nephrodium lineatum*, *Pruet. Epim. Bot.* p. 48. *Hook. Sp. Fil.* iv. 72. *Bedd. F. B. I. t.* 133. *Asp. lineatum*, *Bl. En. Fil. Fav.* 144, (not *Wall.*) *Nephro. glandulosum*, *Bedd. F. B. I. t.* 132, (not *Blume.*)

Himalayas, from Gurwhal to Bhotan and Chittagong 1,000-5,000 feet elevation; Ceylon, Oodawella Forest; not in South India. There can be no doubt that this is a *Nephrodium*, and that involucres are often present, though Mr. Clarke has still kept it in *Goniopteris*. Sir W. Hooker described it both as a *Nephrodium* and as a *Goniopteris*.

(Also in Java and Philippines.)

11. *NEPHRODIUM MOULMEINENSE*. (*Bedd.*) Stipes firm, erect, 2 feet or more long; fronds up to 4 feet long, pinnate, pinnæ often with a large gland at base, numerous coriaceous, about 1 foot long and $1\frac{1}{4}$ – $1\frac{1}{2}$ inch broad, narrow oblong, caudate at the apex, the margin with sharp serratures, which are falcate and only about one line deep, sometimes nearly obsolete or obscure, the frond being nearly entire, upper surface generally quite glabrous and shining, under surface with obscure hairs on the costa and veins; veins very prominent 16–21 pair, all but the upper 2–3 pair anastomosing with a zigzag excurrent veinlet united throughout; sori medial on the veins or often near their apex in a single row on each side of the costule; indusium reniform. *Bedd. F. Sup.*

P. 18. *Goniopteris lineata*, *Bedd. F. B. I.*

t. 3. *G. multilineata*, *Bedd. F. B. I. t.* 231,

(not 232, *Penangiana*, which is the next species). *Polypodium multilineatum* (*Wall.*),

Hook. Syn. Fil. 316. I have been obliged

to rename this as it is a *Nephrodium*, and

there is another *Neph. multilineatum*, *i. e.*

Wallich's (our *truncatum*).

South India, very common on the Golconda Hills (west of Vizagapatam) 2,000–

3,000 feet elevation, involucre distinct; (not


on the Western mountains). Throughout

North India, from the plains up to 5,000

feet elevation, very common; Birma, very common near Moulmein

(involucre distinct); this differs from *urophyllum* in being more

rigid in texture and in having many more veins.



Nº 144.

NEPHRODIUM

MOULMEINENSE. (*Bedd.*)

12. *NEPHRODIUM COSTATUM*. (*Wall. under Polypodium.*) Stipes

$1\frac{1}{2}$ foot and more long, rather stout, scaleless, glossy, and as well as the

rachis strongly tinged with red (rarely stramineous); fronds 1–3 feet

long, broad-oblong or lanceolate coriaceous-submembranaceous

pinnated glabrous, pinnæ numerous, rather distant, patent, sessile 5–8

inches long, $\frac{1}{4}$ an inch or a little more broad (on sterile fronds some-

times exceeding an inch) from an obliquely cuneate truncate sessile

base (lower ones rather more attenuated and subpetiolate) lanceolate or elongate-oblong, finely acuminate at the apex, the margin coarse and sharply submucronate-serrated, serratures pointing a lit forward, uniform; costæ prominent beneath, of the same colour as rachis and stipes, generally reddish, glabrous on both sides obscurely hairy on the costa and veins beneath; veinlets about 6 pairs, of which all are connivent except 2-3 short pairs in the tee of the serratures, sori in two series generally at the base of the ve:



NEPHRODIUM ARBUSCULA. (Desv.)

close to the costules, but sometimes near the centre of the veins, involucre reniform, very fugacious and only to be detected on very young fronds. *Bedd. F. B. I. t. 220.* *Polypodium costatum*, (Wall. *Hes not Brack.* P. (*Goniopteris lineatum*, *Coleb. in Herb. W. and Wall. Cat. 300.* *Hook. Fil. v. 12.* *Polypodium pergamum*, *Hook. Sp. Fil. v. Bedd. F. B. I. t. 232* (slightly broader in the pinnae).

North India, Nepal Kumaon at high elevations specimens from Dr. Stuebel (Gurwhal) have the involucre quite distinct; Penang.

13. NEPHRODIUM ARBUSCULA. (Desv.) Caudex small erect at trunk-like, stipes tufted, short, slightly pubescent; fronds 12-15 inches long, pinnate, lanceolate in outline, pinnae numerous close, 2-3 inches long by about $\frac{3}{8}$ inch broad, very shortly petioled, rather blunt at the apex, generally auricled at the superior base, the margin with shallow serratures, lower ones gradually reduced to mere deltoid auricles; texture firm, herbaceous; veins 6-8 pair, the 3-4 lower pairs

anastomosing, the upper ones free, rachis pubescent, upper surface glabrous in age, under surface more or less hairy on all the veins; sori solitary near the middle of the 3-4 lower veins, indusium small reniform. *Desv. Mém. Soc. Linn. vi. p. 253. Hook. Syn. Fil. 292. Bedd. F. S. I. t. 87. Neph. falcilobum, var. pubera, Clarke, F. N. I. p. 515. Wall. Cat. 338, Asp. puberulum.*

South India, Western forests, 2,000-4,000 feet, in beds of rivers; Ceylon, central provinces.

(Also in the Mascarene Islands.)

14. *NEPHRODIUM PENNIGERUM. (Bl.)* Caudex erect; stipes villous or subglabrous up to about 1 foot long; fronds large up to 4 feet long, pinnate, generally with the lower pinnae more or less reduced, sometimes not so; pinnae numerous, spreading 8-10 inches long, 1-1½ inch broad (the fertile sometimes much contracted), apex acuminate, margin cut down only about ¼ towards the rachis into oblong falcate lobes; texture herbaceous, rachis villous, upper surface strigose on the main costa, otherwise glabrous or often more or less strigose particularly near the margins, under surface villous on the veins; veins 8-12 pair, the 5-6 lower ones generally, rarely only 3, anastomosing; sori medial on the veins; involucre reniform, generally persistent. *Aspid. pennigerum, Bl. En. Pl. Far. Fil. 153. Hook. Syn. Fil. 292. Neph. abruptum, Bedd. F. S. I. t. 86. Asp. multilineatum, Wall. Cat. 353. Neph. glandulosum, var. læte-strigosa, Clarke, F. N. I. p. 532. Neph. pennigerum var. multilineata, Clarke. l. c. p. 532. Neph. abortivum (F. Sm.), Hook. Syn. Fil. p. 292.*

South India, common in all the Western mountain forests; Ceylon, central provinces, 2,000-4,000 feet; East Bengal, from Mishmee to Chittagong, at no great elevation. (Clarke's specimens of *glandulosum* var. *læte-strigosa* and *pennigerum* var. *multilineata* are quite the same as the South Indian plant.) Malay Peninsula.

(Also in the Malay Islands and Tropical Africa.)

15. *NEPHRODIUM MOLLE. (Desv.)* Rhizome tufted, or more rarely creeping; stipe and main rachis beneath pubescent, shortly

hairy or glabrous, with few or no scales ; fronds oblong-lanceolate, 1-3 feet long, pinnate, pinnæ spreading generally about 4-6 inches long by $\frac{3}{4}$ inch broad, the lower ones generally more or less reduced, often into mere auricles, sometimes not reduced at all, generally cut down about half-way to the midrib into blunt subfalcate lobes ; texture herbaceous, generally more or less pilose on both sides, sometimes glabrous or subglabrous, except in the costas and costules ; veins 6-8 pair, pinnate in the lobes, the lowest pair (rarely two) anastomosing with an excurrent veinlet, involucres reniform glabrous or hairy. *Desv. Mém. Soc. Linn.* vi. 258. *Hook. Syn. Fil.* 293. *Bedd. F. S. I. t.* 84. *and* *extensum*, *t.* 85, *not* *Bl.* *Polypodium parasiticum* (*Linn.*) *Asp. solutum*, *Wall. Cat.* 350 *and* *tectum*, *Cat.* 394. *Neph. didymosorum*, *Bedd. F. B. I. t.* 200.

Throughout the Indian region from the plains up to 6,000 feet.

(Also throughout the world in tropical and subtropical regions.)

VAR. β AMBOINENSE. (*Presl.*) Fronds smaller, papyraceous, more glabrous, pinnæ smaller, less cut, being generally only slightly crenated or cut down about $\frac{1}{8}$ to $\frac{1}{4}$, very gradually reduced towards the base, the lower ones often being hastate and very acuminate, veins 4-5 pairs, 2 rarely 3 lower pairs anastomosing ; sori generally on 2-3 lower veins, sometimes confined to the lowest pair (didymosorous). *Neph. amboinense*, *Hook. Sp. Fil.* 292. *Nephr. extensum*, var. β minor, *Bedd. F. B. I. t.* 201. Evidently only a form of *molle*, sometimes confounded in herbaria with *Arbuscula*.

North India, Bengal, and Assam, near the foot of the hills ; Ceylon, Ambagama. (*C. P.* 3390.)

NEPHRODIUM PROCURRENS is a name given by Baker (*Syn. Fil. p.* 290.) to *molle* with a creeping root, but there are no other characters to distinguish this form, as I have the very large form of *molle* with the lower pinnæ reduced to deltoid auricles (sent from Ceylon) with the root wide-creeping, as well as forms which have the fronds truncate at the base and no reduced lower pinnæ. (*Wall.* 349, 3rd sheet in this form.)

MR. γ AUREUM. (*Clarke, F. N. I. p. 533.*) Differs by having mute pubescence bright and golden, the fertile fronds longer e barren, and the pinnæ sometimes contracted near the base ; and truncate at the base, *i.e.*, the lower pinna not reduced. It is feared that none of these characters are permanent. It is from Sikkim and Bhotan, 1,000–4,000 feet elevation, and ssam.

PIDIUM MULTIJUGUM. (*Wall. Cat. 348.*) Taken by Mr. in his type of *var. multijuga*, is typical *extensum* (*Bl.*)

MR. DIDYMOSORUM. (*Parish, Bedd. F. B. I. t. 200.*) From rim ; is a very hairy form of molle, lower pinnæ not at all, sori twin, confined to the two lower veinlets (the latter or probably not constant.) *Wall. Cat. 354*, *Asp. canescens*, ngapore, second sheet, is this plant.

p. tectum (*Wallich, Cat. 394*, in Linnæan Herb.) is a deltoid molle with no reduction of the lower pinnæ, and a creeping

. NEPHRODIUM CRINIPES. (*Hook.*) Stipe up to $1\frac{1}{2}$ foot long, rect, clothed, as is the main rachis, with lanceolate-linear scales ; fronds about 2–3 feet long, pinnate, dwarfed below ricles ; pinnæ horizontal, $4-6\frac{1}{2}$ inches long, $\frac{1}{2}-\frac{3}{4}$ inch broad, vn $\frac{1}{2}-\frac{3}{4}$ towards the midrib into oblong scarcely acute lobes, er basal one often somewhat enlarged ; texture herbaceous ; beneath minutely puberulous ; costa, costules and veins sparsnished with longish hairs above ; veins 6–8 on a side, the rest pair anastomosing ; sori medial, but a little nearer the apex ie costule ; indusium reniform glabrous persistent. *Hook. l. 294. Bedd. F. B. I. t. 263.*

orth-east Bengal, up to 1,500 feet elevation, from Nepal to and Chittagong ; Malacca (Mr. Clarke's Nilgiri example is on of a contracted fertile frond of *pennigerum*.)

. NEPHRODIUM FEROX. (*Moore.*) Stipe 1–2 feet long

strong, densely fibrillose, the hairs long black and leaving the stem rough when they fall; fronds 3-4 feet long, pinnate; pinnæ rather close spreading, up to 15 inches long, by $\frac{3}{4}$ -1 $\frac{3}{8}$ inches broad; apex very acuminate, cut down about one-third into lanceolate-falcate lobes; texture rigid coriaceous, both surfaces glossy; veins very prominent, 12-15 on each side, the lower (about 5) pairs anastomosing, the excurrent veinlet joined throughout and very prominent; sori basal on the veins close to the costule. *Hook. Syn. Fil.* 294. *Bedd. F. N. I.* t. 129.

Kumaon. (Mr. Clarke doubts the Himalayan locality, but there are specimens in Kew Herbarium supposed to have been collected there, and I received it from the Calcutta Herbarium as Himalayan.)


(Also in Java and the Philippines.)

18. *NEPHRODIUM TRUNCATUM*. (*Presl.*) Rhizome short, erect, stipes tufted stout erect up to 2 feet long, greyish naked or slightly pubescent; fronds pinnate up to 4 feet long, the lower pinnæ reduced often to mere auricles; rachis slightly pubescent or glabrous; pinnæ numerous up to 11 inches long by 1 $\frac{5}{8}$ inch broad, acuminate at the apex, sessile or subsessile and often somewhat contracted towards the base, cut down about half-way to the partial rachis into oblong lobes which are more or less square at the apex and crenated; texture herbaceous, upper surface generally glabrous, lower glabrous or slightly pubescent on the costa and veins, and sometimes with glandular dots; veins 6-9 pairs, generally the two, rarely 3-4 lower anastomose; sori small medial or nearer the costule than the margin involucre reniform. *Presl. Tent. Pter.* 81. *Hook. Syn. Fil.* 29. *Neph. eusorum* (*Thw.*), *Bedd. F. S. I.* 130. *Asp. prionophyllum* *Wall. Cat.* 355. *Asp. multilineatum*, *Wall. Cat.* 353, from Singapore. *Wall.* 352, 3rd sheet.

South India, Tinnevely and Travancore Hills; Ceylon, Central Provinces, 2,000-5,000 feet elevation; Cachar and Chittagong Hills at no elevation; Malay Peninsula.

(Also in Malay Islands, North Australia, and Polynesia.)

19. *NEPHRODIUM BRACHYODON*. (*Hook*) Stipes tufted, 1-2 feet long, greyish naked : fronds up to 3 feet long, pinnate ; pinnae petiolate, often contracted at base, 6-9 inches long, $1\frac{1}{4}$ - $1\frac{1}{2}$ inches broad cut about one-third or half-way down into blunt slightly falcate lobes, lower pinnae not much smaller than the others ; texture sub-coriaceous ; rachis naked or slightly pubescent, upper surface glabrous, except the partial rachis which is slightly hairy, under surface glabrous except the rachis and costules which are slightly hairy ; veins 8-12, all free or the lower one anastomosing and with sometimes an excurrent veinlet, a single vein generally rises from the rachis of the pinnae, between the pinnate groups of veins and runs to the margin ; sori medial on the veins ; involucre reniform. *Hook. Syn. Fil.*



Nº 143.

NEPHRODIUM TRUNCATUM. (*Presl.*)

295. *Bedd. F. B. I.* 379. The lower pair of veins only occasionally anastomose in the Malay Peninsula plant, but in the West Indian plant several pairs generally join ; the two plants, however, seem quite to run one into the other, and I am afraid they also connect *Lastrea crassifolia*.

Malacca and Malay Peninsula.

(Also in West Indies and Peru.)

Nephrodium multijugum (*Baker*), given as a Malay Peninsula

fern (*Syn. Fil.* 291) is, as far as the Kew specimen of Wall. Cat. 348 is concerned, *Lastrea ochthodes*. *Aspidium multijugum*, Wall. Cat. 348, in the Linnæan Herbarium is, however, *Nephrodium ex-tensum*, typical, though referred by Clarke to *molle* as a variety. *Nephrod. Hoenkianum* (*Presl.*), given in *Syn. Fil.* p. 291, as a Ceylon fern, does not occur in that island or in our limits.

GENUS LV.—NEPHROLEPIS. (*Schott.*)

(*Nephros*, kidney; *lepis*, scale—the indusium kidney-shaped and scale-like.)

Sori round, arising from the apex of the upper branch of a vein generally near the edge; indusium reniform or roundish; veins forked, free clavate; fronds simply pinnate with the pinnæ articulated at the base, furnished with white cretaceous dots on the upper surface.


1. NEPHROLEPIS CORDIFOLIA. (*Linn. under Polypodium.*) *Caudex* suberect, the wiry fibres often bearing tubers; stipes tufted, with 1–4 inches long, slightly scaly; fronds up to 2 feet, pinnate; pinnæ numerous, crowded, often imbricated, 1–1½ inch long, by ¼–⅝ inch broad, usually blunt; margin entire or slightly crenate, the underside rounded or cordate, the upper distinctly auricled at the base; texture rather coriaceous; rachis scaly, both sides nearly glabrous; sori about half-way between the midrib and margin in a single row; indusium firm, persistent, lunate or reniform. *Polypodium cordifolium*, *L. Sp. Pl.* 1549. *Nephrolepis tuberosa* (*Presl.*), *Hort. Sp. Fil.* iv. 151. *Bedd. F. S. I. t.* 92.

Throughout the Indian region up to 5,000 feet elevation.

(Also in the tropics of the whole world, and in Japan and New Zealand.)

2. NEPHROLEPIS EXALTATA. (*Linn. under Polypodium.*) *Rhizome* suberect; stipes tufted, 4–6 inches long, firm, slightly scaly; fronds up to nearly 3 feet long, pinnate; pinnæ rather close,

Nº144



145

NEPHROLEPIS CORDIFOLIA. (Linn.)

2-3 inches long, $\frac{1}{4}$ - $\frac{1}{2}$ inch broad, usually acute at the apex, the margin entire or slightly crenate, the superior base with a small narrow auricle, the lower base with a smaller rounded one; texture rather coriaceous, both surfaces nearly glabrous, or very scaly particularly when young; sori submarginal; indusium firm, reniform. *Polypodium exaltatum*, *L. Sp. Pl.* 1548. *Nephrolepis exaltata* (*Schott*), *Hook. Syn. Fil.* 301. *Bedd. F. S. I. t.* 93.

South India, the mountain forests of the Eastern and Western sides; East Bengal from Assam to Chittagong, up to 1,000 feet elevation; Ceylon; Malay Peninsula.

(Also in the tropics of nearly all the world.)

3. *NEPHROLEPIS VOLUBILIS*. (*J. Smith, MS.*) Rhizome climbing up to 50 feet over trees, with adpressed chesnut scales on the short lateral distant spurs, whence spring clusters of stipes; pinnae obtuse or not very acute; venation, sori, &c., as in *exaltata*. *Aspid. exaltata* *Wall. Cat.* 1031, partly. *Clarke, F. N. I. p.* 541, *t.* 78.

Sylhet and Chittagong; Malacca.

(Also in Borneo.)

4. *NEPHROLEPIS ACUTA*. (*Presl.*) Rhizome short; stipes tufted up to 8 inches or more long, firm, naked or slightly scaly; fronds up to 8 feet long, pinnate; pinnae up to 8 inches long and $\frac{1}{8}$ inch broad, oblong acuminate, sometimes shortly petioled, the margin slightly crenate; texture thin, both surfaces hairy when young, as is the rachis, but quite glabrous in age; sori near the margin; involucre reniform to suborbicular. *Presl. Hook. Sp. Fil.* iv. 153. *Bedd. F. S. I. t.* 94. *Neph. biserrata* (*Schott*).

South India, common on the North Arcot Hills, at no great elevation; Ceylon; North India, Chittagong Hills, up to 1,000 feet.


(Also in Tropical Africa.)

5. *NEPHROLEPIS RAMOSA*. (*Beauv. under Aspidium.*) Rhizome slender, wide-creeping; stipes very short, scattered; fronds up to 12 inches long, 1-3 inches broad, pinnate; pinnae $\frac{1}{4}$ - $1\frac{1}{2}$ inch long.

$\frac{1}{2}$ – $\frac{3}{4}$ inch broad, slightly crenate, the upper edge auricled and truncate, parallel with the stem, the lower oblique; texture papyraceous, rachis villous, both surfaces slightly villous or glabrous in age; sori near the margin; involucre cordiform, roundish fugacious. *Beauv. in the Flore d'Oware. Hook. Syn. Fil.* 301. *Neph. oblitterata (Hook), Bedd. F. S. I.* 251.

Ceylon, central provinces, 2,000–5,000 feet, climbing on trees; Malay Peninsula.

(Also in Tropical Africa; Philippines; Australia; Fiji.)



NEPHROLEPIS RAMOSA. (*Beauv.*)

GENUS LVI.—OLEANDRA. (*Cav.*)


(Resemblance to the Oleander.)

Sori round, inserted in a row near the base or below the centre of the compact free veinlets; involucre reniform; fronds entire lanceolate-elliptical, stems jointed, rhizome wide-creeping.

1. *OLEANDRA NERIIFORMIS. (Cav.)* Shoots woody, wide-creeping but often suberect, clothed with short adpressed scales which are often deciduous; stipes short, seldom 1 inch long, with the joint below the middle; fronds 4–8 inches long, $\frac{1}{2}$ – $1\frac{1}{2}$ inch broad, in opposite pairs or often in terminal whorls, or more rarely scattered, from narrow-linear to oblong-acuminate; texture subcoriaceous, both sides glabrous or hairy underneath; sori in two rather irregular rows near the midrib. *Hook. Fil. Exot. t.* 58. *Syn. Fil.* 302. *Bedd. F. B. I. t.* 264.

Himalayas from Nepal eastwards; Khasya; 2,000–5,000 feet elevation.

(Also in Central America.)




OLEANDRA NERIIIFORMIS. (Cav.)

2. *OLEANDRA MUSÆFOLIA*. (*Kunze.*) Shoots firm, wide-creeping, clothed with long golden scales which are not deciduous; fronds generally single, scattered, or more rarely in tufts, 12-20 inches long, by about $1\frac{1}{2}$ inch broad, narrow oblong acuminate on stipes which are 1-2 inches long and jointed close to the base; texture thin, a little hairy on both sides and the midrib scaly below; sori in two irregular rows near the midrib. *Mett. Fil. Ind. p. 240.* *Hook. Syn. Fil. 302.* *Oleandra neriiformis*, *Bedd. F. S. I. 91.*

South India on the Western mountains, particularly abundant

in Coorg, rare elsewhere; Ceylon, central provinces. Mr. Clarke considers this species not distinct from the last; they appear to me as distinct as the others; all four, however, are so closely allied that they might be all considered varieties of one species.



OLEANDRA WALLICHII. (*Presl.*)

times paleaceous, jointed close to the base, so that the very short lower articulation is concealed among the scales; fronds 6-14 inches long, membranaceous soft, more or less hairy, subelliptical-oblong, the sides parallel, the base often obtuse and rounded, the apex suddenly and sharply acuminate; sori compact, almost imbricated in a continuous line close to and parallel with the costa on each side; involucres reniform, ciliated, opening towards the margin.

3. *OLEANDRA WALLICHII.* (*Presl.*) Caudex creeping, rooting, thicker than a goose-quill, densely clothed with subulate crisped ferruginous, spreading and often reflexed scales thickly tufted at the extremity of the branches; stipes distant $\frac{1}{4}$ -3 inches long and some-

Presl. Tent. Pter. p. 78. Hook. Sp. Fil. iv. 158. Aspid. Wallichii, Hook. Ex. Fl. i. p. 6. t. 5. Bedd. F. B. I. t. 265.

Birma and the Malay Peninsula.

4. OLEANDRA CUMINGII. (*J. Sm.*) Caudex creeping, scarcely so thick as a writing pen, clothed with close-pressed subulate imbricated scales, stipes subterminal and subaggregated 2–7 inches long slightly downy, articulated within $\frac{1}{2}$ –3 inches from the base; fronds 1–1 $\frac{1}{2}$ foot long, firm membranaceous, elongate lanceolate acuminate, very much attenuated and gradually decurrent at the base, pubescent-villous on the costa and veins, sori rather large, forming a single and scarcely interrupted line very near to and on each side the costa. *J. Sm. in Hook. Jour. Bot. iii. 413. Hook. Sp. Fil. iv. 158. Bedd. F. B. I. t. 135.*

Birma and the Malay Peninsula. The Assam locality is doubted by Mr. Clarke, who thinks the specimen in the Kew Herbarium of Griffith's was collected at Malacca. The Nilgiri locality is an error, the specimen being musæfolia (*Kze.*).

(Also in the Philippines and Canton.)

B. EXINVOLUCRATÆ. — Sori without an indusium. Tribes Polypodiæ to Acrosticheæ.

TRIBE X.—POLYPODIEÆ.

Sori on the back of the lobes, round or rarely short-oblong.


* *Desmobryoid series, habit and mode of growth of Aspidiæ, the stems continuous with the caudex; sori generally medial on the veins.*

GENUS LVII.—PHEGOPTERIS. (*Fée.*)

(*Phegos*, beech—the beech fern.)

Sori round, veins forked or pinnate, veinlets free; fronds from pinnate to decomposed. Habit of *Lastrea*, and only differing from that genus in wanting an indusium.

PHEGOPTERIS SCOTTII. (*Bedd.*) Caudex suberect, stout clothed with brownish-black subulate scales as is the base of the stipe; fronds pinnate, oblong, with the stipe 10–12 inches long, but 3 inches broad, rachis furnished with hair-like scales; alternate 6–7 approximated pairs, with a terminal one, lower one much decreased in size, all short oblong or ovate obtuse with a square unequal base about $1\frac{1}{2}$ inches long, by a little less than 1 inch broad, and furnished with falcate acute serratures, glabrous on both sides, but with a few weak setæ or scales on the costae, especially beneath; veins pinnate, the lower veinlet two lower not reaching more than half-way to the margin, upper ones terminating in a dot within the margin; sori medial on the lower veinlets. *Bedd. F. B. I. t.* 345. Near Darjeeling, Valley of the Teesta, 2,000 feet elevation (perhaps an abnormal form of *Lastrea* or *hirtipes*).



Nº148.

PHEGOPTERIS SCOTTII. (*Bedd.*)

PHEGOPTERIS ERUBESCENS. (*under Polypodium.*) Stipes 2–3 feet and more long, stout, as the rachis and costa more or less purplish-tawny, fronds in length ample but varying from 1 to 4 feet in length and from 6 to more than 2 feet in breadth, firm, sub-coriaceous, broad-cuminate, pinnate, pinnæ 3–16 inches long, $\frac{1}{2}$ – $1\frac{1}{2}$ inch wide, alternate, sessile, elongate-oblong, the sides parallel for a long distance and then gradually acuminate to a serrated apex, deeply nearly costate pinnatifid, segments oblong, subfalcate, rather acute, or obscurely serrated, glabrous above, beneath sparingly furnished with longish white needle-like hairs on the rachis and costae sometimes on the veins; veins approximate, simple free, two opposite pair meeting but scarcely uniting at the sinus,

soriferous always at the very base, so as to form two lines or series (at length confluent) one on each side and close to the costa, not extending to the apex of the segment; main rachis stout with a broad groove on the upper side (when dry.) *Wall. Cat. n. 330. Hook. Sp. Fil. iv. 236. Bedd. F. B. I. t. 213. Hook. Syn. Fil. 306.*

Himalayas, Kashmir to Bhotan, Khasya, 3,000–7,000 feet elevation; Malay Peninsula. It is very like *Lastrea tylodes*.


(Also in Malay Islands.)

3. *PHEGOPTERIS AURICULATA.* (*Wall. under Polypodium.*) Stipes stout, densely villous, with white matted hairs (as is the rachis), furnished below with numerous ovate black shining scales; fronds ample, firm, membranaceous, 3 feet and more long, 16 inches broad, oblong-lanceolate acuminate, generally gradually attenuated below by the dwarfing of the pinnæ, pinnated, pinnæ numerous, about 8 inches long by 1 inch broad, sessile and furnished at the base on the under side with a curious hooked gland, sometimes nearly quarter inch long, pinnatifid nearly to the rachis, segments broad oblong, very obtuse, entire, subfalcate, costa costules and veins villous, and the segments sparingly hairy on both sides between the veins; veins pinnate, simple, extending to the margin, lowest pair entering the margin above the sinus, soriferous below the middle, spore cases furnished with a few long weak hairs. *Polypodium auriculatum, Wall. Cat. 314. Hook. Syn. Fil. 306. Bedd. F. B. I. t. 203.*

Himalayas, from Gurwhal to Bhotan, 5,000–8,000 feet elevation, common about Darjeeling, Khasya, Myrung wood, 5,000 feet elevation. *Polypodium appendiculatum, Bedd. F. B. I. 256*, is only a state of this with the pinnæ not reduced at the base.

(Also in Java.)

4. *PHEGOPTERIS VULGARIS.* (*Mett.*) Rhizome slender creeping;



CHAM.

PHIEGOPTERIS AURICULATA (Walt.)

stipe with large lanceolate yellowish-brown scales near the base fronds 6–9 inches long by 4–6 inches broad, subdeltoid, lowest pinnæ as long or nearly as long as any above them, main rachis beneath, with small ovate scales, pinnæ $\frac{1}{2}$ – $\frac{3}{4}$ inch broad, cut down nearly to the main rachis, lobes oblong blunt or scarcely acute, entire or slightly crenate, $1\frac{1}{2}$ –2 lines broad, more or less pilose on both sides; veinlets 6–8 on a side, the lower ones forked; sori nearer the margin than the midrib. *Mett. Fil. Hort. Lips.* 83. *Polypodium Phegopteris*, *Hook. Syn. Fil.* 308. *Clarke, F. N. I.* 544.

Cashmir, above Sonamurg, 11,000 feet elevation.

(Also in North Europe, Asia, and America; Caucasus and Japan.)

5. *PHEGOPTERIS DISTANS*. (*Don. under Polypodium.*) Stipe tufted, squamose near the base up to 2 feet long, yellowish or dark purplish-brown, glossy; fronds up to about 3 feet long, the lower pinnæ generally reduced and distant, sometimes not at all reduced, pinnæ 6–8 inches long, $1\frac{1}{2}$ inch broad, cut down nearly or quite to the rachis into deeply pinnatifid pinnules $\frac{1}{4}$ inch broad, with blunt or acute toothed or subentire lobes, the base dilated, texture herbaceous, rachis villous or rarely glabrous, undersurface a little hairy; veinlets pinnate in the ultimate lobes, pellucid; sori generally on the lower veinlets below the apex, or rarely apical, the veinlet not being continued beyond the sorus. *Polypodium*, *Don. Prod., Fl. Nep.* 2. *Polypodium paludosum* (*Ll.*), *Bedd. F. S. I. t.* 168. *P. Griffithii*, *Hook. Sp. Fil.* iv. 236. *P. longipes*, *Wall. Cat.* 316. *P. adnatum*, *Wall. Cat.* 328. *P. brunneum*, *Wall. Cat.* 333. This variety “adnata,” *Clarke, F. N. I.* p. 544 (which is *Lastrea microstegia*, *Hook. Sp. Fil.* iv. 119 and *Bedd. F. B. I. t.* 39) only differs in being a little more compound and generally bipinnate, but it runs into the type. Var. “glabrata” of Clarke only differs in being somewhat more glabrous. Var. “minor” of Clarke is a smaller, less cut form, pinnæ only 1–2 inches long, pinnatifid only one-third down, but it runs into the type.

North India, from Kashmir to Bhotan, 3,000–8,000 feet elevation, very common; Khasya, 3,000–5,000; South India, on the Western

mountains, 5,000–8,000 feet, very abundant; Ceylon; Malay Peninsula.

6. *PHEGOPTERIS RUFESCENS*. (*Blume, under Polypodium.*) Rhizome creeping, furnished with appressed scales and wiry roots, stipe 6 inches long and together with the main and partial rachises highly asperous; fronds 12–18 inches long by 9–12 inches broad, deltoid, subcoriaceous, puberulous beneath, with minute yellow spots, bi-tripinnatifid, *i. e.* lower pair of pinnæ often bipinnatifid, the other ones pinnatifid, pinnules long obtuse, spinulose-seriate; veins pinnate, 3–4 on a side, veinlets simple or forked, reaching to the margin; sori circular on the veinlets. *Polypodium, Bl. Fil. Jav. p. 194, t. 1. Hook. Sp. Fil. iv. 257. Syn. t. 309. Bedd. F. S. I. t. 236.*

Ceylon, central provinces, 300–5,000 feet elevation.

(Also in Java, New Caledonia and Queensland.)

7. *PHEGOPTERIS DRYOPTERIS*. (*Linn. under Polypodium.*) Rhizome creeping; stipe with lanceolate subulate scales near the base; fronds

8 inches long and broad, deltoid, 3-pinnate, thin green, nearly smooth, often glandulose, lower pinnæ largest, pinnules lanceolate, the lowest free, oblong slightly crenate, texture thin herbaceous, this glabrous; veins pinnate in the lobes, veinlets forked or simple; a submarginal a little below the apex of the veinlet. *Polypodium dryopteris, Linn. Sp. Pl. 1555. Hook. Syn. Fil. 309. Bedd. F. B. I. t. 74. Polypodium Robertianum, Milde, Fil. Eur. 99.*




PHEGOPTERIS RUFESCENS. (Blume.)

Kashmir, 7,000—11,000 feet elevation, and eastwards to Kumaon.
(Also in Arctic and Alpine Europe, Asia and America.)

8. *PHEGopteris ornata*. (*Wall. under Polypodium*.) Caudex erect; stipe, main and partial rachises with lanceolate-linear scales and muricate or scabrous from their persistent bases, not pilose, viscous or glaucous; fronds very large up to 15–20 feet long, tripinnate, except that there is a very regular wing to the rachis, pinnæ generally 2 feet or more long and about 1 foot broad, secondary pinnæ 1–2 inches broad,



№151.
PHEGopteris dryopteris. (*Linn.*)¹



№152.
PHEGopteris ornata. (*Wall.*)

their rachis also scabrous and scaly and furnished with long white needle-like hairs, tertiary pinnules up to $\frac{1}{4}$ inch broad, linear-oblong from a sessile square base, resting on the wing of the tertiary rachis, rather distant from each other, acute or blunt at the apex, deeply often nearly to the costa pinnatifid, the costa and often the veins furnished with needle-like hairs, texture thin herbaceous; veinlets simple or forked; sori medial. *Wall. Cat.* 327. *Bedd. F. S. I. t.* 171. *Lastrea tenericaulis*, *Hook. Sp. Fil.* iv. t. 269, not the description which

Sp. refers to *Lastrea tenericaulis*. *Phegopteris pallida* (*Brack.*), *Sp. Fil.* iv. 266.

Himalayas, from Kumaon to Bhotan, in tropical valleys up to 5000 feet elevation, common in Chittagong Hills, 500 feet elevation ; in India, Carccor ghat, Malabar and elsewhere along the Western Ghats, but not common ; Malay Peninsula.

One of the handsomest of Indian ferns, and much in cultivation, may be a *Lastrea* (if *Phegopteris* is to be kept distinct from that genus, the propriety of which I doubt), but no one has yet detected a difference, in any case it is quite distinct from *Lastrea tenericaulis*.

(Also in North Australia and Polynesia.)

9. *PHEGOPTERIS PUNCTATA*. (*Thunb. under Polypodium*.) Rhizome firm, wide-creeping, villous ; stipes scattered 1-2 feet long and with the lower part of the main rachis more or less viscidulous ; fronds 1-4 feet long, tripinnate, pinnæ up to about 1 foot long, lanceolate to deltoid, secondary pinnæ 4-6 inches long, tertiary pinnæ sessile on the rachis, which is not winged, blunt at the apex, and pinnated two-thirds down into short rounded crenated lobes, more or less furnished on both sides with scurfy hair-like scales, the margins often somewhat reflexed, as in *Hypolepis* ; sori numerous towards the apex of the veinlets and near the margin. *Thunb.*

Jap. 337. *Phegopteris punctata*, *Hook. Syn. Fil.* p. 312. *Polypodium rugulosum*, *Hook. Sp. Fil.* iv. 272. *Bedd. F. S. I. t. 1.* *Polyp. rugulosum*, *Labill. Fl. Nov. Hol.* ii. 92 t. 241. *Hypolepis hostilis*, *Presl. (as to the Nilgiri plant.)*

Himalayas, from Chumba to Bhotan and Chittagong, 1,000-5,000 feet elevation ; South India, common on the Western mountains at the higher elevations ; Ceylon, about Newera Elya ; Malay Peninsula.

(Also almost throughout the tropics and south temperate zone, extending to Japan, New Zealand, St. Helena, and Chili ; not from Continental Africa.)

10 *PHEGOPTERIS SUBDIGITATA*. (*Blume, under Polypodium*.) Stipes tufted, firm, continuous with the rachis, furfuraceous, or

drying quite green, glabrous; veinlets forking well within the margin and clavate at the apex of the veinlet. *Polypodium Fil.* 196. *t.* 93. *Bedd. F. B. I. t.* : *Phegopteris davallioides* (Mett.), *Hooker's coniiifolium*, *Wall. Cat.* 326.

Himalayas, from Nepal to Bhotan common ; Malay Peninsula.


(Also in the Malay Islands.)

GENUS. LVIII.—GONIOPTERIS

(*Gonia*, angle ; *pteris*, fern—the veinlets at an angle)

Habit and venation of *Nephrodium*. Lowest or several pairs of veinlets of cordate shape at an angle from which proceeds an excurrent veinlet continuous with the rhizome ; fronds pinnate as in *Nephrodium*, except that there is no indusium.

All the supposed Indian species except *Gonia* proved to possess an indusium in a young condition, so that they have been transferred to the following species deviates somewhat from the



GONIOPTERIS PROLIFERA. (Roxb.)

Cat. 312, in *Calc. Journ. Nat. Hist.*
315. *Bedd. F. S. I. t.* 172. *Menis*

Throughout the Indian region g
ditches in the plains or low down on
(Also in North Australia, Trop
Islands ; Philippines ; New Caledoni

GENUS LIX.—DICTYC


(*Diktyon*, a net ; *ptēris*, fern

Habit and venation of *Aspidium*
mosing with or without free included
Aspidium in wanting an indusium.

1. DICTYOPTERIS BARBERI. (*Ho*
long, slightly scaly at the base ; frond
usually pinnate with a large terminal se
the upper ones oblong-lanceolate, 4-6
nearly entire, the lowest pair with a de
on the lower side, texture subcoriaceous
rather large and regular, with conio

der, glossy, stramineous, the base very tomentose with a few
 rse scales partly subterraneous, attached to the caudex by a small
 nt; fronds thin, membranaceous pale-green, glabrous, subpuber-
 us on the veins, 6-9 inches long, quite as broad as long, cordate-
 toid, subternate or pinnated with 3-9 subfalcate pinnæ, lateral
 s nearly opposite, petiolate, lowest pair half ovate, acuminate,
 1 inches long, lobate-pinnatifid, those of the superior margin with
 ad ovate nearly equal subdentate blunt lobes, the inferior
 s much elongated and acuminate, the basal ones again lobate
 natifid, intermediate pinnæ ovate-lanceolate, pinnatifid, terminal
 na long petiolate, broad-ovate
 minate, deeply pinnatifid
 ecially at the base; veinlets
 formly reticulated, areoles
 a no included free veinlets,
 tular areoles large; sori rather
 gularly scattered, dorsal or
 apital. *Hook. Sp. Fil.* v. 104.
F. B. I. t. 4.

Moulmein; very like small
 imens of *Aspidium cicatarium*
 habit and outline, but there
 no free veins, and the sori
 more scattered and larger,
 show no signs of an in-
 ium.




N°154.

DICTYOPTERIS BARBERI. (*Hook.*)

3. DICTYOPTERIS CHATTAGRAMICA. (*Clarke.*) Rhizome short,
 t; stipes 10-12 inches long, tufted, long, black with small scales
 the base; fronds deltoid, distinctly dimorphic, pinnate below,
 natifid upwards about 10-12 inches long by 6-7 inches broad,
 r pinnæ deltoid with the lower basal lobes much enlarged, mar-
 pinnatifid, surfaces pubescent, the margin with minute golden
 cellular hairs; sori naked, medial or terminal on the veins.
Clarke, F. N. I. p. 548, t. 81.

Chittagong at no elevation, very general.

half ovate, unequally bipartite, lo
times free, and 5-6 inches long, a
often to within one-third of the c




Nº155.

DICTYOPTERIS CHATTAGRAMICA. (*Clarke.*)

Polypodium confluens, *Wall. Cat.* 3

The Malay Peninsula, Tenasse

/A1--



Л. С. ЧЕЛОВЕКОВ

DICTYOPTERIS DIFFORMIS. (Bl.)

texture membranaceous, surfaces glabrous; main veins distinct to margin; areoles with free included veinlets; sori copiously scattered, terminal or compital. *Mett. Kuhn. Lin. 36, p. 124. Baker, Hook. Syn. Fil. 506.*

Malacca.

6. *DICTYOPTERIS HETEROSORA. (Baker.)* Stipe short brown, with spreading linear scales; fronds oblong-spathulate, 2–3 feet long, simply pinnatifid, the lowest foot only a narrow wing to the rachis, above this 3–4 pairs of distant oblong-lanceolate acuminate pinnae, 1½–2 inches broad at middle, confluent in a narrow wing, the lowest simple or bipartite, texture thin but firm, surfaces glabrous, ribs naked brownish; main veins distinct to edge, 3–4 lines apart with distinct arches between them, and copious small areoles with free included veinlets; sori very abundant, small irregular, often confluent. *Baker, Hook. Syn. Fil. 506.*

Malacca, (habit of *Aspidium vastum*.)

** *Eremobryoid series.* Stems articulated at the point of junction with the rhizome; sori generally, not always, terminal on the veins.

GENUS LX.—POLYPODIUM. (*Linn.*)

(*Polys*, many; *pous*, foot; *polypus*,—the rhizome when destitute of fronds, having the appearance of some kinds of sea polypus.)

Veins all free; sori naked, generally punctiform, round, rarely oblong, generally terminal on the veinlets, or more rarely below the apex; fronds simple or pinnatifid or rarely pinnate or more compound.

* *Fronds entire.*

1. *POLYPODIUM PARASITICUM. (Mett.)* Rhizome erect, rarely creeping; stipes tufted, sessile, hairy or fuscous, fronds 2–3 inches long, ¼–¾ inch broad (rarely more), narrow linear (rarely abnormally pinnatifid), blunt at the apex, gradually narrowed at the


re, the margin entire or slightly undulated, texture herbaceous to coriaceous, both sides more or less clothed with long hairs; veins more or less immersed, forked; sori often mixed with copious hairs, round or linear, at the apex of the superior veinlet forming a single row on each side of the costa. *Hook. Syn. Fil. p. 319. Bedd. S. I. t. 165. P. mediale (Baker), Hook. Syn. Fil. p. 507.*

South India, common on the Nilgiris and other Western mountains at the highest elevations; Ceylon. (This includes *P. mediale* of Baker, the sori often being linear and globose on different fronds of the same plant, as in figure quoted above).

Dr. Wall sends some specimens from Ceylon with a creeping root, but I never saw it creeping in the Nilgiri plant.

2. *POLYPODIUM BEVENOSUM. (Baker.)*

Plants tufted, very short, black wiry; fronds 3-4 inches long, $\frac{1}{2}$ inch broad, ligulate, the point antish, the lower part narrowed very gradually, the edge slightly repand, texture subcoriaceous, colour bright green, the underside black, slightly fur-




Nº157

POLYPODIUM PARASITICUM. (Mett.)

ple, very short and indistinct; sori round, tending slightly towards oblong, in a long row on each side close to the midrib. *Hook. Syn. Fil. p. 320. Bedd. F. B. I. t. 323.*

Penang.

3. *POLYPODIUM ZEYLANICUM. (Mett.)* Rhizome strong, wide-



creeping, clothed with broad grey scales ; stipe 1 inch or more long, clothed with soft spreading hairs ; fronds 8–12 inches long, $\frac{1}{4}$ – $\frac{3}{8}$ inch broad, acuminate, the base narrowed very gradually, the margins slightly undulated ; texture subcoriaceous, both sides naked ; veins forked oblique, rarely simple ; sori round, at the apex of the upper veinlet, or at the apex of the vein when simple, forming a single row on each side of the costa. *Hook. Syn. Fil.* 321. *Bedd. F. S. I. t.* 237.

Ceylon, Newera Elya, on rocks and trees.


4. *POLYPODIUM WALLII*. (*Bedd.*) Rhizome creeping, scaly, furnished with numerous thick black wiry roots ; stipes 3–6 lines long, glabrous and hairy ; fronds linear-lanceolate with a blunt apex 3 inches long by $\frac{1}{2}$ inch broad, thick, coriaceous, the veins quite hidden, more or less clothed on both sides and on the margins with delicate black hairs ; veins thick, forked (or simple towards the apex of the frond) ; sori large, round, terminal on the superior veinlet and forming a regular row on each side about midway between the costa and the margin. *Bedd. Fern Sup. t.* 380. *Polypodium parasiticum*, var. *obtusiusculum*, *Thw. MS. C. P.* 3921.

Ceylon, Adam's Peak ; allied to *parasiticum*, but much larger and of much thicker texture.

5. *POLYPODIUM HIRTELLUM*. (*Bl.*) Caudex small ascending, glabrous above ; stipes tufted $\frac{1}{2}$ – $1\frac{1}{2}$ inch long, slender, filiform, villous, with spreading reddish-brown hairs, fronds firm-membranaceous, linear-oblong to lanceolate, 1–4 or more inches long, $\frac{1}{8}$ – $\frac{1}{3}$ of an inch broad, quite entire at the margin, attenuated at the base, rather densely villous and ciliated with similar hairs to those of the stipes ; veins spreading, simple or often forked near the base ; sori in a single regular series close to the costa ; veins when simple soriferous near their base, when forked, soriferous at the apex of the short fork. *Bl. En. Fil. Jav. p.* 123. *Hook. Sp. Fil.* iv. 166. *Polypodium hirtum*, *Hook. Sp. Fil.* iv. 170, (*not Mett.*). *Bedd. F. B. I.* 212. *P. lasiosorum* (*Hook.*), *Bedd. F. B. I. t.* 172.


Ceylon.

(Also in Java and Luzon.)



Nº15.

POLYPODIUM HII



6. *POLYPODIUM SESSILIFOLIUM*. (*Hook.*) Caudex small, erect ascending, scaly; stipes scarcely any; fronds tufted, membranous, firm, 3-9 inches long, 2-3 lines wide, linear, scarcely minated, obtuse, attenuated at the base; veins oblique, approximate, forked; sori oblong, parallel with the costa in lines or series intermediate between the costa and the margin, medial on the upper let. *Hook. Fil.* iv. 168. *Syn. Fil.* p. 322. *Bedd. F. B. I. t.* 119. Penang.

(Also in the Philippines.)

** *Fronds pinnatifid.*

7. *POLYPODIUM CORNIGERUM*. (*Hook.*) Fronds tufted, subsericeous, erect ligulate, 3-4 inches long, 2½ lines broad, pinnatifid or bipinnate, lobes 20-30 jugate, diminishing in size from middle of and both ways, almost triangular, subacute, repand on the upper edge; texture subcoriaceous, surfaces glabrous; veins one to each lobe, forked, falling far short of the margin; sorus large round, solitary at fork close to main rachis, and medial on the vein. *Hook. Fil.* 503. *Bedd. Fern Sup.* 381.

Ceylon, Horton plains, rare, (P. 4005.)




8. *POLYPODIUM CUCULLATUM*. (*Nees.*) Caudex very small, ascending; stipes densely tufted 3 lines long, naked; fronds subcoriaceous or somewhat membranous, 3-5 inches long, 2-3 lines broad, flaccid, linear-lanceolate uninate, deeply nearly to the costa pinnatifid; the lobes linear-

oblong $\frac{1}{4}$ - $\frac{1}{2}$ line broad, those of the upper half fertile, broader and with the edges upcurved so as to clasp the large solitary sorus; costa and sometimes the frond deciduously pilose. *Hook. Syn. Fil.* 324. *Calymmodon cucullatus* (*Presl.*), *Bedd. F. S. I. t.* 233.

Ceylon, on rocks and trees, central and southern provinces, 3,000-5,000 feet elevation.

9. *POLYPODIUM KHASVANUM.* (*Hook.*) Stipes tufted, very short.



N°162.

POLYPODIUM TRICHOMANOIDES. (*Sw.*)

midrib. *Hook. Syn. Fil.* 325. *Bedd. F. B. I. t.* 173.

Khasya 3,000-4,000 feet, near Cherra; Assam, from Cherra to Jowye and Jarain in Jaintea, 3,000-4,000 feet elevation, not plentiful anywhere.

10. *POLYPODIUM TRICHOMANOIDES.* (*Sw.*) Caudex short, erect, paleaceous above, densely rooting with wiry fibres; stipes short, $\frac{1}{4}$ - $\frac{1}{2}$ inch long, densely tufted, patent-villous with long fibrous hairs, fronds 3-4 inches to a span long, 2-4 lines wide, rigid-membranaceous,

fringed with soft hairs. fronds up to about 14 inches long, by $1\frac{1}{2}$ inch broad, flaccid, cut down nearly to the rachis into linear-oblong entire or slightly undulated blunt lobes 2-3 lines broad, texture thick subcoriaceous; rachis and both sides sparingly hairy; veinlets simple; sori apical on the veinlets, sunk in an oval cavity (having an elevation on the upper side of the frond) in rows of 4-6 on each side of the

tenuated at both extremities, deeply nearly to the costa or almost pinnate (quite so at the very base), lobes horizontal from a broad base, ovate or oblong entire, sub-concave sub-deciduously ciliated with long fibrous or blackish hairs; veins solitary in each lobe, simple or forked, each lobe with a single sorus near the base of the veins and the costa. *Hook. Bot. Beechey.* t. 178; *Syn. Fil.* 326. *Bedd. F. B. I. t. 2.*

Common, 9,000–12,000 feet elevation.

Found in South America, Brazil and Ecuador; and Cuba.)

POLYPODIUM GLANDULOSUM. (*Hook.*) Caudex small, creeping, clinging to the bark of trees by copious glandular scales, the rest of the plant all over glandulose, most so beneath; stipes 1–2 lines long, fronds 2–4 inches long, 1–1½ linear obtuse, scarcely attenuated to the extremity, rather firm-membranaceous-succulent, deeply nearly to the rachis pinnatifid; segments ovate, sub-acute, erect-patent, decurrent at the base, the upper ones free, all of them serrate-pinnatifid, and rather distant few and obsolete veins indistinct; sori 1–6 to each segment globose. *Hook. Sp. Fil.* iv. 193. 327. *Bedd. F. S. I. t. 238B.*

Common, in the central provinces, Ramdoo summit of Wattakelia Hill.



Nº163.
POLYPODIUM
GLANDULOSUM. (*Hook.*)

POLYPODIUM THWAITESII. (*Bedd.*) Caudex creeping, with lanceolate scales; fronds sessile or subsessile, sub-linear, perfectly glabrous, 2–4 inches long by about 1 inch broad, lanceolate or oblanceolate, deeply almost to the rachis pinnatifid, the lower reduced to a narrow wing; veinlets distant, erect-patent, simple or rarely forked, where simple soriferous at the apex, where

forked the superior veinlet only is soriferous; sori round, slightly immersed. *Bedd. F. B. I. t. 188. Baker, in Hook. Syn. Fil. p. 508.*

Ceylon, about Dickoya, on trees, very sweet scented (C.P. 3,900).



Nº164.

POLYPODIUM THWAITESII. (Bedd.)

Exp. p. 7, t. 2, f. 2. Bedd. F. B. I. t. 238A. P. serra, Wall. Cat. t. 313.

13. *POLYPODIUM DECORUM. (Brack.)* ♂ Caudex short, rather thick, creeping, densely ferruginous-squamose, stipes approximate, sub-terminal on the caudex, 2-4 lines long; fronds coriaceous, glabrous, 6-12 inches long, 1-1 inch broad, narrow-lanceolate, much and almost caudate-acuminate, the base very gradually attenuated into the short stipe, deeply and nearly to the rachis pinnatifid, segments horizontally patent, narrow-oblong obtuse, quite entire, below gradually becoming shorter and broader and forming shallow elongated lobed wings at the base; costa glabrous or pilosulous, costule and veins quite sunk and inconspicuous; sori oblong, 2-8 in a row, on each side of the costule, and apical on the short veins, which do not nearly reach the margin, partially sunk in a hairy cavity (but with no raised border). *Hook. Sp. Fil. iv. 179. Brack. Fil. Un. St.*

South India, on the Tinnevely Mountains, near Courtallum; Ceylon, Singhe Rajah Forest; Singapore and Mount Ophir; North-east India, near Mandal (Edgeworth).

(Also in Borneo, Luzon, Tahiti.)

14. *POLYPODIUM FUSCATUM*. (BL) Stipes tufted, 1-3 inches long, densely clothed with soft spreading hairs; fronds 3-6 inches long, $\frac{1}{2}$ -1 inch broad, cut down nearly to the rachis throughout into loose entire obtuse pinnæ, line broad, the lower ones shorter and broader, texture subcoriaceous, rachis and both sides thinly clothed with soft hairs; veins obscure, simple; sori forming a close row on each side the midrib, at last filling nearly the whole surface.

Hook. Syn. Fil. p. 331.

Vedd. F. B. I. t. 324.

Ceylon.

(Also in Java.)


*** *Fronds pinnate.*

15. *POLYPODIUM BILLOQUATUM*. (BL) Stipes tufted, 1 inch or more long, rigid, naked or vil-

lous; fronds up to 15 inches long and 2 inches broad, pinnate, being cut down to the rachis throughout into entire close subhorizontal near pinnæ, 1-1 $\frac{1}{2}$ lines broad, dilated at the base, the lower ones warped, texture subcoriaceous; rachis naked or hispid, both sides naked; veinlets simple, not reaching the margin; sori sunk in a cavity 4-6 on each side, apical on the veinlets; the edges of the



POLYPODIUM DECORUM. (Brack.)



fertile pinnæ sometimes undulated. *Hook. Syn. Fil.* 328. *Bedd. F. S. I.* 167. *Cryptosorus, Fée. Cœlopteris, A. Br.*

South India, Anamallay Mountains, up the Toracadu River, 4,000-5,000 feet elevation, Tinnevelly and Travancore Hills; Ceylon, common on trees about Newera Elya.

16. POLYPODIUM

REPANDULUM. (*Mett.*)

Caudex stout, erect, with densely matted roots; stipes tufted, very short, naked or hairy; fronds 4-6 inches long, about 1 inch broad, cut down to the rachis throughout, into erect-patent blunt crenated pinnæ $\frac{1}{4}$ - $\frac{1}{2}$ lines broad; texture subcoriaceous; rachis naked, or with a few very long hairs; surfaces naked, or with long hairs beneath on the costa; veinlets simple, falling short of the margin; sori slightly immersed, filling nearly the whole space between the



Nº 167

POLYPODIUM REPANDULUM. (*Mett.*)

midrib and the margin, and apical on the veinlets. *Hook. Syn. Fil.* 328. *Polypodium subfalcatum*, var. β , *Bedd. F. B. I. t.* 189B. *Polyp. minutum (Baker)*, *Hook. Syn. Fil.* 328, as far as the Ceylon plant is concerned. *C. P.* 3073.

Ceylon, higher parts of central provinces, on trunks of trees.

17. *POLYPODIUM SUBFALCATUM*. (Bl.) Rhizome erect; stipes densely tufted, clothed with soft spreading hairs; fronds 2-10 inches long, by $\frac{1}{4}$ -1 inch broad pinnate, *i.e.* cut down to the rachis into close spreading pinnæ; pinnæ hairy on both sides, soft in texture, rather sharply toothed sometimes one-third or half-down, decurrent at the base, the lower ones gradually reduced; veinlets simple, not reaching the margin; sori apical on the veinlets one to each tooth or lobe of the pinnæ. *Bl. Fil. Jav.* 186, t. 87. *A. B. Hook. Syn. Fil.* 328.



POLYPODIUM SUBFALCATUM. (Bl.)

Polyp. parvulum, *Becc.* *F. S. I.* t. 166, not *Becc.* *P. subfalcatum*, *Becc.* *F. B. I.* 189, fig. A. (fig. B.) *P. parvulum* *Thw. En. Pl. Zey.* 394 in part, *C. P.* 129 not 3073. It differs from the last in being soft in texture instead of coriaceous, in being much more hairy (*repandulum* being generally quite glabrous), in being serrate instead of crenulate, and in the sori not being immersed. Mr. Thwaites included them both under his *parvulum*, and Mr. Baker has the former both under *repandulum* and *minutum* in the *Synopsis Filicum*.

South India, Nilgiris Lamb's Rock near Conoor, Anamallays banks of Toracadu river, 4,500 feet; Ceylon, central provinces; North India, Himalayas from Gurwhal to Bhotan, 5,000-9,000 feet elevation; Khasya. Mr. Clarke says that some of the Khasya specimens are as small as *P. trichomanoides*, but that species has only one sorus to each pinna, whilst this has the pinna polysorus, or



POLYPODIUM DARCEIFORME. (Hook.)

one to each lobe of the pinna, besides the sori are differently situated.

* * * * *Fronde compound.*

18. POLYPODIUM DAREÆFORME. (*Hook.*) Rhizome ~~stolon~~ creeping, clothed densely with filiform golden scales; stipes 6–9 inches long, articulated on the rhizome, naked, glossy; fronds 12–18 inches long, 8–12 inches broad, sub-deltoid, 3–4-pinnate; Darea-like; pinnae 1–2 inches broad, lanceolate deltoid; secondary pinnae $\frac{3}{4}$ inch long, tertiary pinnae small, forked or pinnate; the ultimate segments narrow, almost filiform bluntish, texture herbaceous, rachis glossy, both sides quite naked; veins one to each ultimate segment, not reaching the margin, and thickened at the apex, and bearing the sorus low down; sori extending beyond the margins of the segments; indusium none. *Hook. Sp. Fil.* iv. 256, *Syn. Fil.* 339. *Acrophorus Hookeri*, *Bedd. F. B. I.* 95 (*not Moorei*, which is *Leucostegia Hookeri*.)

Himalayas and Khasya, 4,000–5,000 feet elevation. This cannot belong to *Phegopteris* as the stipes are articulated with the rhizome; it is very probably a *Leucostegia*, but no indusium is to be detected in the many examples that I have examined; it has long been confused with *Leucostegia Hookeri*, which it closely resembles in habit.

GENUS LXI.—GONIOPHLEBIUM. (*Presl.*)

(*Gonia*, angle; *phebes*, veins—the veins forming angles.)

Veins forming ample regular areoles, each with a simple or rarely forked free included veinlet, on which the sorus is terminal, often in the costal areoles only, but sometimes also in the second or even the third row, marginal veinlets free; sori non-indusiate, globose or rarely oblong; fronds pinnatifid or pinnate (simple in some species not Indian); rhizome wide-creeping, the stipe articulate on the rhizome; pinnae articulate with the rachis.

* *Fronde pinnatifid or somewhat pinnate at the base only.*


1. *GONIOPHLEBIUM AMÆNUM.* (*Wall.*) Caudex creeping, out, densely paleaceous, with ferruginous subulate scales, which are sub-adpressed, never hair-pointed; stipes a span to a foot or more long, stramineous or brown; fronds 1 to nearly 2 feet long, 10 inches wide, glabrous or sub-pubescent, ovate, terminating in a lanceolate acuminate subentire segment, deeply pinnatifid to within 3 lines of the costa, segments horizontal, more or less approximate, 8 inches long, $\frac{1}{4}$ – $\frac{1}{3}$ inch wide, subfalcate from a broad base, cuneiform gradually acuminate, entire, or coarsely dentate-serrate, lowest pair deflexed, veins forming one costal series of moderately sized soriferous areoles, and sometimes two (the second not soriferous), marginal veinlets free, soriferous veinlet always arising from the vein at a distance from the costa; sori subglobose sunk (papillate on the upper side.) *Hook. Sp. Fil.* v. 24. *Wall. Cat. n.* 290. *edd. F. B. I. t.* 5.

Himalayas, from Gurwhal to Bhotan, 4,000–11,000 feet elevation, common; Khasya 3,000–6,000 feet. (Clarke's var. *anglensis* from Tonglu near Darjeeling, appears only to differ in the pinnæ being blunt at the apex, there is, however, only one poor specimen known of it.)

2. *GONIOPHLEBIUM SUBAMÆNUM.* (*Clarke.*) Rhizome slender, densely clothed with grey-brown lanceolate-subulate scales, which are adpressed, never hair-pointed, near the base of the main rachis are sometimes ovate or lanceolate scales; stipes about 2 inches long; fronds 6 inches long, deeply pinnatifid nearly to the rachis, lower segments much abbreviated and deflexed, all serrated at the margins; costal arches of the main rachis continued nearly or quite to the base of the frond; main rachis above glabrous or puberulous. *Clarke, N. I. p.* 550, *t.* 82, *f.* 2.

Very nearly allied to *amænum*, and perhaps only a form of it, differs in its slender rhizome and small size.

Himalayas, on the ridge between Sikkim and Nepal, 11,000–15,000 feet elevation, very abundant.




A.S. CATTELL & C. H. HARRIS

GONIOPHLEBIUM AMENUM. (Wall.)

3. *GONIOPHLEBIUM LACHNOPUS*. (*Wall.*) Rhizome more slender than in *amænum*, densely clothed with hair-pointed brown-black scales spreading from small bases; base of the rachis often with a few larger scales; stipe 2-4 inches long, slender naked; fronds flaccid, 18 inches long, 3-4 inches broad, cut down nearly to the rachis spreading slightly-lanched segments $\frac{1}{4}$ inch broad, texture herbaceous, both sides naked, the rachis beneath minutely scaly, and pubescent above; costal areoles in single series (never as sometimes occurs in *amænum*), or rarely veins are all free (not forming areoles); sori in single series in the areoles, or at the apex of the naked veinlet when there is no anastomosis. *Wall.* t. 310. *Hook. Syn.* p. 342. *Bedd. F. B. I.* t. 63. *Clarke, F. N. I.* 551.

Himalayas, from Kashmir to Bhotan, 1000-3,000 feet elevation, very common. Sikkim, 3,000-5,000 feet elevation.



Nº 171.

GONIOPHLEBIUM ERYTHROCARPUM. (*Mett.*)

4. *GONIOPHLEBIUM ERYTHROCARPUM*. (*Mett.*) Rhizome $\frac{1}{4}$ line thick, glaucous, firm, wide-creeping, scales lanceolate, dark-castaneous,


minute, not hair-pointed ; stipe 3-4 inches long, naked, stramineous, very slender ; frond oblong-deltoid, 3-4 inches long, $1\frac{1}{2}$ -2 inches broad, pinnatifid nearly to the rachis, lobes 5-10 jugate, ligulate blunt, 3-4 lines broad, obscurely repand, adnate, only the lowest narrowed at the base, on the lower side scarcely shortened, texture rigidly subcoriaceous ; rachis and costas below and under surface pilose ; main veins raised, very distinct to nearly the edge, costal areoles in one or two series, the second series when present generally without a free veinlet in lower areole, sometimes forked, or rarely netted and graduating into the venation of *Pleopeltis* ; sori in a single series in the lower areoles. *Mett. Kuhn. Linn.* 36, p. 135. *Hook. Syn. Fil.* 511. *Bedd. F. B. I. Sup.* 382.

All my specimens from Messrs. Clarke and Atkinson have typical *Goniophlebium* venation, so also have nearly all the specimens at Kew. Mr. Clarke, however, has removed the species to *Pleopeltis*, because some specimens gathered by him have the costal areoles less defined and the veinlets somewhat netted and graduating into the venation of *Pleopeltis*. I look upon these as abnormal, the whole habit of the plant being that of *Goniophlebium*.

Sikkim, Lacheen, 9,000-11,000 feet elevation, Yakla valley, 8,000 feet.

5. *GONIOPHLEBIUM HENDERSONI*. (*Atkinson, MS.*) Rhizome very slender, glaucous, wide-creeping, clothed with hair-pointed scales ; stipes 3-4 inches long, naked or scaly, very slender, grey, stramineous ; fronds narrow-lanceolate 8-10 inches long, $1\frac{1}{2}$ -2 inches broad, deeply nearly to the rachis pinnatifid, or the lower lobes or pinnæ quite free ; pinnæ or lobes 20-30 pair, patent lanceolate acute, 3-4 lines broad, distantly incised, all except the lowest dilated at their base, texture moderately firm, surfaces green, naked ; areoles and sori uniserial, the latter near the midrib, not crowded, occasionally the veins are all or nearly all quite free and not anastomosing, or forming areoles, costal arches of the main rachis prominent, very long and narrow, absent of course at the base of the larger fronds, where it is pinnate, not pinnatifid. *Hook. Syn. Fil.* p. 511. *Bedd. F. B. I. t.* 383.

Nº172.



ACORONAS

GONIOPHLEBIUM MOLLE. (*Bedd.*)

Himalayas, dividing ridge between Sikkim and Nepal, and north of the Jongri, 11,000 to 13,000 feet elevation, very abundant.

6. GONIOPHLEBIUM MICRORHIZOMA. (*Clarke*.) Rhizome 1 line thick, firm wide-creeping, clothed with grey-brown ovate or lanceolate scales, not hair-pointed; base of the rachis often with a few similar scales; stipes 4–5 inches long; fronds lanceolate, often caudate at the apex, about 1 foot long by 3–4 inches wide, deeply nearly to the rachis pinnatifid, the lowest pinnæ sometimes free; pinnæ 20–30 pair, patent lanceolate acute, 3–4 lines broad, distantly incised, texture herbaceous, rachis glabrous above, castaneous beneath; areoles and sori uniserial, or all the veins often quite free, and not forming areoles. *Clarke, F. N. I.* 551. *Hook. Syn. Fil.* p. 511. *Bedd. F. B. I. t.* 384.

Kashmir to Bhotan, 5,000–8,000 feet elevation, very common.


* * *Fronds distinctly pinnate throughout.*

7. GONIOPHLEBIUM MOLLE. (*Bedd.*) Rhizome creeping, size of the stem of a tobacco pipe, densely covered with reddish brown subulate scales; fronds 1½ foot long, 7–8 inches broad, oblong acuminate membranaceous, pinnæ numerous, 3–4 inches long, ½ an inch wide, furnished on both sides as is the rachis with soft weak whitish hairs, sinuate-crenate, superior base often subauricled, superior ones decurrent on the rachis, remainder sessile from a broad base, or inferior ones subpetiolate; veins forming two series of areoles, the lower of which has the soriferous veinlet arising from the side of the areole distant from the costa, marginal veinlets free or rarely uniting, simple or forked. *Bedd. F. B. I. t.* 206. *Polypodium Beddomei* (*Baker*), *Hook. Syn. Fil. 2nd Edit.* 344.

Malay Peninsula, in Tenasserim.

8. GONIOPHLEBIUM SUBAURICULATUM. (*Bl. under Polypodium*). Rhizome creeping, stoloniferous, densely clothed with narrow acuminate, often hair-pointed scales, subpruinose; stipes a span to a foot and more long, sparingly paleaceous, rufous-brown; fronds firm, subcoriaceous, glabrous, 2–6 feet long, a foot and more wide, pinnated, pinnæ horizontal, numerous yet remote, 3–10 inches long, ¼ to nearly ½ an inch wide, shortly petioled and jointed on the rachis from a

slightly cordate or truncated and occasionally auricled base, linear-ovate, finely and gradually acuminate, serrated; veins forming a distinct costular series of areoles, including a free venule, of which the outer series is soriferous, marginal veinlets free; sori in a single series forming slightly elevated pustules on the superiorside. *Bl. Fil. Jav. p. 177, t. 1.*
Hook. Sp. Fil. v. 32.
Add. F. B. I. t. 78.



Nº 173.

Khasya, 3,000-5,000 feet elevation; Malay Peninsula. *GONIOPHLEBIUM SUBAURICULATUM. (Bl.)*
 (Also in Malay Islands and Queensland.)

9. *GONIOPHLEBIUM ARGUTUM. (Wall. under Polypodium.)*

Rhizome thick as a writing-till, squarrose with ovate-acute short blackish or brown glossy spreading scales, stipes 3-4 inches to a span long, brown glossy; fronds firm-membraceous, glabrous, 1-2 feet long, 8-10 inches broad, pinnated, pinnæ distant subhorizontally patent, 4-6 inches long, $\frac{1}{4}$ - $\frac{1}{2}$ inch wide, from a broader but obtusely and oblique cuneate or subtruncated base (rarely rotundate-auricled) longate-oblong, acuminate, either obtusely serrated, lower ones often opposite, the rest ternate, superior ones more



Nº 174.

GONIOPHLEBIUM ARGUTUM. (Wall.)


less adnate and decurrent, terminal one similar to the rest, veins forming a costal series of large oblong soriferous areoles; marginal veins ending in a thickened apex within the margin all free; sori super-

ficial at the apex of the free veins in the costal areoles. *Wall. Cat. n. 308. Hook. Sp. Fil. v. 32. Bedd. F. B. I. t. 6.*

Himalayas, from Kashmir to Bhotan, 4,000–9,000 feet elevation, very common. Closely allied to the last species.

10. GONIOPHLEBIUM VERRUCOSUM. (*Wall. under Polypodium.*)

Rhizome long, stout creeping, very paleaceous; stipes $1\frac{1}{2}$ foot and more long; fronds 2–3 feet long, oblong-acuminate, firm-membranaceous, drooping; pinnæ numerous but distant, 6–9 inches long, $1-1\frac{1}{2}$ inch broad, oblong costate, articulate upon the rachis, suddenly and shortly cuspidate-acuminate entire or serrated chiefly towards the apex,



N°175.

GONIOPHLEBIUM
VERRUCOSUM. (*Wall.*)

nearly sessile, the base obliquely cuneate; primary veins slender but straight and parallel costuliform, forming with the anastomosing veins four or five series of areoles each with a free included veinlet, but of which the lowest series only is soriferous; sori in a single series next the costa sunk in a deep cavity, having a corresponding pustule on the upper side. *Wall. Cat. n. 296. Hook. Sp. Fil. v. 3¹. Bedd. F. B. I. t. 257.*

Malacca and Penang.

(Also in the Philippines.)

Mr. J. Smith has separated these East Indian species of Goniophlebium from the American because the pinnæ are articulated with the rachis, and he has constituted the genus Schellolepis for them.

GENUS LXII.—NIPHOBOLUS. (*Kaulf.*)

(*Niphos*, of snow; *bolus*, a large pill—the snow-like scales and round sori.)

Fronds on their under surface matted with woolly or cottony tomentum; sori globose or elliptic, superficial or immersed, buried in the tomentum; veins internal, obscure, pinnate, prominent or


form from a central costa, venules anastomosing sometimes transversely parallel, forming parallelogrammoid areoles, with excurrent or occasionally connivent or anastomosed veinlets, sometimes merging in roundish or oblong hexagonal unequal oblique areoles, with uniformly directed simple or divaricately forked veinlets, the veins of the fertile frond when contracted less developed; fronds simple or rarely lobed, rigid, coriaceous, opaque, the fertile often contracted.

1. NIPHOBOLUS ADNASCENS. (*Sw. under Polypodium.*) Rhizome wide-creeping, paleaceous with lanceolate-setaceous scales; stipes distant, 1-2 inches long, furnished at the base with lanceolate-linear hairy scales; fronds dimorphous, carnose-coriaceous, dark green above, but hoary with sparse stellated pubescence beneath, and under and even white with more copious compact hairs; sterile fronds, 4 inches long, spatulate or elliptical-lanceolate, obtuse; fertile ones 8 inches long, linear or oblong obtuse or acute, both kinds tapering slowly into the stipe, costa sub-carinate, costules sunk obscure, their areoles including four (sometimes only three or two) veinlets, which are generally free with clavate apices, though they sometimes anastomose; the deep sunk in the tomentum and in the substance of the frond on each side of the costa, arranged 5-6 in obliquely transverse series, costules long stalked, mixed with long stalked stellated scales.
v. Syn. Fil. pp. 25 and 228. Hooker. Sp. Fil. v. 47. Bedd. F. S. t. 184. Wall. Cat. 268. Polyp. verrucosum, Wall. Cat. n. 267. vittarioides, Wall. Cat. p. 270. P. pertusum, Wall. Cat. 267.

Throughout India from the plains up to 4,500 elevation; Ceylon; Malay Peninsula.

(Also in China; Fiji; Mascareen Islands, Cameroon Mountains.)

2. NIPHOBOLUS LÆVIS. (*J. Sm.*) Rhizome slender, wide-creeping, clothed with linear setaceous scales; stipes distant, somewhat hairy $\frac{1}{4}$ inch to nearly 2 inches long, furnished with a tuft of scales at their base; fronds coriaceous, linear-lanceolate, much narrowed towards the apex, 2 $\frac{1}{2}$ -6 inches long, $\frac{1}{4}$ - $\frac{3}{4}$ inch broad, glabrous above, hairy with stellate pubescence beneath; venation obscure, costules in areoles in three series between the costa and margin, with one,



NIPHOROLUS ADNASCENS. (Ste.)

0 or three clavate veinlets in each, which are variously directed, or rarely joined to the base of the areole above; sori large, partially sunk in the tomentum, often confined to the apex of the frond, or scattered, or covering nearly all the under surface. *Bedd. F. I. t.* 161. *P. Niphobolus jaintense*, *Clarke, F. N. I. p.* 552, *t.* 82.

Khasya hills, Jaintea, Jarain, 3,500 feet elevation. Considered a variety of *adnascens* by Mr. Baker, but sufficiently distinct in its constitution, I think, to rank as a species.


3. *NIPHOBOLUS ACROSTICHOIDES*. (*Sw.*) Rhizome wide-creeping, branched, younger portions squarrose, with rather large bright ferruginous lanceolate scales; stipes 1–3 inches long furnished at the base with obtuse scales; frond hard coriaceous, 1–2 feet and more long, 1 inch wide, glabrous above, beneath whitish or tawny, stellate-tomentose (tomentum very deciduous) elongate-lanceolate or linear use; sterile fronds generally the broadest; areoles 5–7 between the costa and margin, each with 3–6 veinlets free and simple, or variously forked and anastomosing; sori rather small, prominent, very compact, generally occupying the upper part of the frond, arranged in 6–8 regular very close series between the costules. *Sw. Syn. Fil. p.* 297, *t.* 225. *Hook. Sp. Fil. v.* 44. *Bedd. F. B. I. t.* 81. *Polyp furfuraceum*, *Wall. Cat. n.* 278.

Birma and the Malay Peninsula generally. I have specimens from Tenasserim 2 feet long and only $\frac{1}{4}$ inch wide.

4. *NIPHOBOLUS HETERACTIS*. (*Mett.*) Rhizome wide-creeping, scales copious lanceolate-linear, not adpressed golden; stipes persistent up to 7 inches long, more or less scaly; fronds oblong, with a long acumination about 6 inches long by $1\frac{3}{4}$ inches broad, coriaceous, the fertile not generally contracted, upper surface glabrous, lower clothed with close compact stellate scales, the rays of which are broad, with filamentous hairs admixed; areoles in 8–9 series between the costa and margin, each including 2–3 veinlets, which are generally forked and more or less anastomosing; sori rather large and prominent. *Mett. Kuhn in Linn.* 36, 140. *Clarke, F. N. I. p.* 35. *N. Lingua*, *Hook. Syn. Fil.* 350, *in part*, and *p.* 512. *Bedd. F. Supl. t.* 385.

Sikkim and Bhotan, 4,000–6,000 feet elevation ; Kharya, 3,000–5,000 feet. This is very near the Japan Lingua, but it appears to differ sufficiently to be considered a species, besides the filamentous hairs radiating round the scales, which it requires a magnifying glass to detect, the habit is somewhat different and the scales of the rhizome more free and copious.

5. *NIPHOBOLUS PANNOSUS*. (*Mett.*) Rhizome wide-creeping, slender, scales linear, dark coloured, closely adpressed ; stipes up to 6 inches long, stellately scaly ; fronds lanceolate up to 6 inches long by $1\frac{1}{2}$ inch broad, narrowed at both ends, coriaceous, glabrous above, at least in age, beneath densely matted with stellate scales, the rays of which are hair-like ; areoles in about seven series between the costa and margin, each including 2–4 veinlets which are simple, forked, free, or variously anastomosing ; sori crowded, rather small, 5–6 in a row between the main veins, not reaching the margins. *Mett. Kuhn.*



N^o 177.

NIPHOBOLUS PANNOSUS. (*Mett.*) *Linn.* 36, p. 141. *Hook. Syn. Fil.* 512. *Niphobolus Lingua*, *Bedd. F. S. I. t.* 240. Ceylon, up to 3,000 feet elevation ; Tenasserim.

6. *NIPHOBOLUS STIGMOSUS*. (*Sw.*) Rhizome wide-creeping, stout, rufous-paleaceous with slender subulate scales ; stipes subaggregate, $1-6\frac{1}{2}$ inches long, angled ; fronds a span to $1\frac{1}{2}-2$ feet and more long, $1-2-3$ inches and more broad, firm, coriaceous, in age glabrous above, beneath densely stellate-tomentose and subfurfuraceous (tomentum forming a close coat), lanceolate or suboblanceolate acuminate.

ally tapering below and decurrent upon the stipes, costules erect, elevated on the under side; sori immersed, very minute and numerous, arranged in compact closely placed lines or series between costules and in equally-compact series transversely with them; veins in about 10 series between the costa and margin, each including many veinlets which are much branched and anastomosing.

Syn. Fil. 29, 226. *N. costatus*, *Hook. Sp. Fil.* v. 50. *Wall. p.* 265. *Bedd. F. B. I. t.* 120.

Himalayas, from Gurwhal to Sikkim, 2,000–6,000 feet elevation, Assam, 2,000–3,000 feet, Parasurama, 2,500 feet; Tenasserim; Kondah Hills West of Vizagapatnam. My Birma specimens have long stipes and the frond scarcely decurrent, my Himalaya and Vizagapatnam specimens have the frond very short and decurrent, and the stipe very short.

7. *NIPHOBOLUS SUBFURFURACEUS*. (*Hook.*) Rhizome short, creeping, the younger fronds paleaceous with ferruginous late scales, stipes subaggregated, 1–2 inches long, nearly glabrous, fronds 24–30 inches long, 4–5 inches

broad-lanceolate or oblanceolate, sharply acuminate below, usually and much attenuated upon the stipes, glabrous above, densely sellate-tomentose beneath, the tomentum thin, firm and so as to appear subfurfuraceous or compactly pannose of a greyish colour, costa stout, costules evident but not elevated; venation indistinct, quite that of true *Campyloneuron*, the areoles from 10 to 15 in a series between the costa and the margin, each with 10–15 (8–9) included free or scarcely anastomosing soriferous veins; sori rather small, elevated, subglobose (not in the least sunk),



Fig 178.

NIPHOBOLUS STIGMOSUS. (*Steud.*)

forming as many arched series between the costules as there are areoles. *Hook. Sp. Fil.* v. 52. *Bedd. F. B. I. t.* 259.

Bhotan and Mishmee. The areoles are much more numerous than in *stigmatus*, and the included veinlets also more numerous, but much less anastomosing, the fronds in my specimens are of thin texture, the veinlets being visible (without soaking the frond).



Nº 179.

NIPHOBOLUS FISSUS. (Bl.)

Rhizome creeping, paleaceous with ferruginous scales, stipes aggregated short scarcely any or 3—5 inches long, and then winged margined to the base; fronds coriaceous 6 inches to a foot and more long, $\frac{1}{4}$ an inch to 1—1 inch wide, lanceolate or linear-lanceolate or even linear, often finely acuminate, the margin quite entire (or rarely spuriously and very unequally pinnatifid with remote long and narrow segments), from near the middle gradually attenuate downwards to the caudex, at first wholly tomentose with ferruginous stellated hairs, at length glabrous on the upper side and depressed punctate, the dots corresponding with the sori; venation quite sunk, veins pinnate but not prominent, venules transversely parallel and forming with the veins parallelogrammoid areoles (about five between costa and margin), with two, sometimes only one, excurrent free clavate veinlets in each areole; sori copious and at first quite sunk among the tomentum appearing in the form of small tubercles, at length a circular opening appears, but the sori scarcely rise above the surface of the tomentum, arranged in series. *Bl. Fil. Jan. p.* 58. *t.* 24. *Niph. porosus*, *Hook. Sp. Fil.* v. 48. *Bedd. F. S. I. t.* 183.

Adiantum porosum, Wall. Cat. n. 266. Niph. Schmidianus, Kze. Zeit. vi. p. 121. Polyp. mysurense (Heyne), Wall. Cat. 269. floccigerus (Mett.), Bedd. F. Sup. t. 386.

South India, Western mountains, 3,000–8,000 feet elevation; N., central provinces; Himalayas and Khasya, 2,000–7,000 feet elevation; Malay Peninsula. (The Khasya floccigerus only in the fronds being narrower.)

. NIPHOBOLUS FLOCCULOSUS.

Rhizome short, stout densely especially at base of stipes, finely subulate golden scales; 2–8 inches long, aggregated, and as are the fronds with whitish or brownish stellate tomentum; fronds 4–16 inches long by $\frac{1}{2}$ –1 $\frac{1}{2}$ broad, coriaceous-lanceolate, finely acuminate, serrate, moderately attenuated pinnate, or broad and unequally hastate at the base, in age green above and blackish punctate-venation immersed, costules distinct, areoles 9–16 each with included veinlets which are usually free, simple or forked,




N° 180.

rarely anastomosing with a distinct line from the top of the areole; sori scarcely sunk, small but very elevated above the tomentum, very copious, arranged in regular rows between the costules. Don. Prod. Fl. Nep. 1. N. gibbilis, Hook. Sp. Fil. v. 49. Bedd. F. R. I t. 162. P. costalis, Wall. Cat. 265, partly.

Himalayas, Gurwhal to Bhotan, up to 5,000 feet elevation, Khasya, extending into the Sylhet plains.


10. NIPHOBOLUS GARDNERI. (Mett.) Rhizome short-creeper.

ing, the younger portions densely ferrugineous-paleaceous; stipes approximate, arising from a scaly branch of the caudex, 2-4 inches long; fronds about a foot long, carnose-coriaceous lanceolate, ob-



N°181.

NIPHOBOLUS GARDNERI. (Mett.)



tusely acuminate, gradually attenuated upon the stipes, densely clothed with a very compact firm sub-furfuraceous mass of whitish or ferruginous stellated tomentum, costa and primary veins or costules slightly elevated beneath, venation of *Campyloneurum*, secondary transverse veins more obscure, areoles about 7 between

the main costa and margin, each including 2-4 veinlets, which are generally simple with a clavate apex, rarely forked and rarely anastomosing with adjoining veinlets; sori superficial (not sunk) in

3-4 series, parallel with the costules and 10-12 transverse series between the costa and the margin. *Mett. Polypod. p. 129. Hook. Sp. Fil. v. 51. Bedd. F. S. l. t. 241.*

Ceylon, rocks and trees in central and southern provinces up to 3,000 feet; South India, Annamallay Hills, 5,000 feet elevation.



N°182.

NIPHOBOLUS PENANGIANUS. (Hook.)

ob lanceolate finely acuminate, the base much and gradually attenuated, sessile, the margin irregularly sinuated, glabrous above (at least in maturity), beneath thinly clothed with fuscous stellated hairs, venation internal, but manifest

II. *NIPHOBOLUS PENANGIANUS.* (Hook.) Rhizome short creeping; stipes scarcely any; fronds submembranaceous, 1 foot and more long, 2½ inches wide,

n viewed between the eye and the light, costules not elevated, les 10-12 between the costa and the margin, including 2-4 ple or forked clavate soriferous veinlets; sori prominent (not c), forming a broad mass in the disk of the upper half of the d, not reaching the margin; capsules mixed with long stipitate ate stellated scales. *Hook. Sp. Fil.* v. 52. *Hook. Ic. Pl.* t. 203. . *Fil.* 83, *Polycarpium*, *Pr.* *Bedd. F. B. I.* t. 121.

Penang, Tenasserim (Parish).


12. NIPHOBOLUS BOOTHII.

ok.) Rhizome short with lan- late-linear golden scales; stipes 6 inches long, and stout in pro- tion, tawny brown, paleaceous, b imbricating ferruginous lanceo- : scales only at the very base; ad carnose-coriaceous, 16-24 bes long, 3-4 inches wide, el- ic-lanceolate, obtusely acuminate, clerately attenuated, glabrous, and ctated above with minute black-

dots (probably corresponding a the receptacles of the sori), eath covered with a dense vel- mass of ferruginous stellated entum; primary costular veins lent but not prominent, united by sverse arched veins, forming ay series of areoles, each with ncluded veinlets, which are much nched or anastomosing; sori

all, partially sunk in the tomentum ming transverse lines between the costules and as many as there are oles. *Hook. Sp. Fil.* v. 53. *Bedd. F. B. I.* t. 258.

Bhotan.



NIPHOBOLUS MUMMULARIFOLIUS.
(Mett.)

13. *NIPHOBOLUS NUMMULARIÆFOLIUS*. (*Sw. under Acrostichum*). Rhizome slender wide-creeping, filiform, clothed with bright ferruginous lanceolate fibrillose scales; fronds dimorphous, carnosé-coriaceous, the barren ones roundish or elliptical, subsessile $\frac{1}{2}$ –1 inch each way, the fertile ones longer and narrower, $1\frac{1}{2}$ –2 inches long. $\frac{1}{2}$ – $\frac{1}{3}$ inch broad, upper surface naked, at least in age, lower densely coated with loose ferruginous woolly tomentum; veins forming areoles irregular in size and shape, empty or with one free veinlet, marginal veinlets free; sori close, scattered, sometimes covering the whole surface. *Sw. Syn. Fil.* 191, 419 to 2. *Polyp. nummulariæfolium. Mett. Farngatt Polyp.* 122, t. 3. *Hook. Sp. Fil.* v. 54. *Syn. Fil.* 351. *Bedd. F. B. I.* t. 320. *Galeoglossa, Presl.*

East Bengal, Bhotan, Assam, Khasya, Cachar, up to 2,500 feet elevation. There is a variety from Khasya (*obovatum*, Mett.) with the sterile fronds obovate on a $\frac{3}{4}$ inch petiole. This species is abnormal in the genus, I do not think it is a *Niphobolus* at all, but probably a *Drymoglossum*.

GENUS LXIII.—DIPTERIS. (*Reinw.*)

(*Di*, two; *ptēris*, fern—the fan-like fronds in two parts.)

Fronds adherent to the caudex, flabellate in two halves which are deeply lobed from their circumference in the direction of the base, secondary veins forming many areoles, including free or netted veinlets; sori punctiform, numerous, scattered small, or in a single series.

1. *DIPTERIS WALLICHII*. (*Br.*) Rhizome stout, creeping, clothed with appressed copious black hard subulate setaceous scales wrinkled at the back; stipes 1–2 and more feet long, smooth and polished, distant erect; fronds coriaceous, $1-2\frac{1}{2}$ feet long and much wider than long, dark-green above, whitish or ferruginous beneath, flabelliform bipartite into two nearly equal broad-cuneate portions which are palmately and dichotomously divided, ultimate segments 6–10 inches long, $1\frac{1}{2}$ –2 inches wide, oblong acuminate; costæ from the summit of the stipes dichotomously branched through the



ACTEPM.

DIPTERIS WALLICHI. (Br.)

disk of the frond, a single pair runs through each terminal segment connivent at the apices; venation manifest prominent beneath, all the costæ are united by transverse flexuose costules, these by a longitudinal flexuose vein through the middle into two nearly equal series of costular areoles, then again into lesser ones all subquadrangular; sori small, superficial copious, scattered in the areoles, but generally in series more or less perfect and accompanied by a resinous or gummy substance. *R. Br. in Hook. and Grev. Ic. Fil. t. 168-9. Hook. Sp. Fil. v. 99. Wall. Cat. n. 287. Bedd. F. B. I. t. 80.*



DIPTERIS HORSFIELDII. (*Br.*)

main veins, confluent, not gummy. *R. Br. Hook. Sp. Fil. v. p. 99. Bedd. F. B. I. t. 321. Polyp. conjugatum, Kaulf. (not Lam.) Polyp. Dipteris (Bl.), Hook. Syn. Fil. 362.*

Malacca, Penang and Singapore.


(Also in the Malay and Pacific Islands.)

Khasya, up to 4,000 feet, Borpani, 2,000 feet, and elsewhere, Cachar, at Luckipoor, 250 feet elevation, Jaintea, Sylhet station in the plains.

2. DIPTERIS HORSFIELDII.

(*Br.*) Stipe strong, 3-5 feet long, frond 1-3 feet long, and much broader, the main lobes reaching three-quarters of the way down, the edges sharply toothed, texture subcoriaceous, upper surface dark-green, lower very glaucous, main veins very prominent, dichotomously forked; areoles copious; sori very minute, scattered irregularly, especially plentiful near the

3. DIPTERIS LOBBIANA. (*Hook.*) Stipes 1 foot and more long; fronds coriaceous, 8-12 inches long, flabelliform, glabrous, 3-4 times



Nº186.
DIPTERIS LOBBIANA. (*Hook.*)



Nº187.
DRYNARIA CORONANS. (*Wall.*)

digitately dichotomous, dark brownish-green above, pale and tawny beneath, segments 4–8 inches long, scarcely half an inch wide, linear-sub lanceolate, finely acuminate costate entire; venation manifest, the costules form a costal series of large soriferous areoles, the rest of the veins unite, constituting lesser areoles and include free veinlets; sori 1–2 in each primary areole (sometimes confluent) in a single series very near the costa. *Hook. Sp. Fil.* v. 100. *Hook. in Kew Gard. Misc.* v. p. 300, t. xi. *Bedd. F. B. I.* t. 233. *Polypodium bifurcatum*, *Baker in Syn. Fil.* p. 362.

Malacca, on Mount Ophir.

GENUS LXIV.—DRYNARIA. (*Bory.*)

(*Dryads*, the sterile fronds being like oak-leaves, a tree sacred to the Dryads.)

Fronds articulate with the caudex, with either a separate sterile frond like an oak leaf, or the base of the frond pinnatifid and oak-leaf-like; veins copiously anastomosing, forming quadrate or hexagonal areoles; sori small, round or oval, numerous


1. *DRYNARIA CORONANS.* (*Wall.*) Rhizome thick, creeping, branched and interwoven, villous-paleaceous with golden scales; fronds large uniform, when growing arranged in a circle, firm, coriaceous, but rather thin and translucent, two and more feet long, from a broad sessile cordate, lobate-pinnatifid base (then suddenly contracted), lanceolate, deeply almost to the rachis pinnatifid, segments 8–12 inches long, $\frac{3}{4}$ –1 inch and more wide, oblong-lanceolate, acuminate, lower ones gradually shorter, entire, incrassated at the margin; venation very manifest, costules parallel extending to the margin, these are connected by transverse veins forming 5–6 soriferous areoles and then again generally two or three others which include free veinlets; sori oval, often confluent longitudinally, solitary in the primary areoles, forming an eccentric series always nearer one costule than the opposite one. *Wall. Cat.* n. 288. *Hook. Sp. Fil.* v. 94

edd. F. B. I. t. 13. D. conjugata, Hook. Syn. Fil. 366 (not m.)

Himalayas, Nepal to Bhotan, 1,000–4,000 feet elevation, rare, Siam, 1,000–3,000 feet common, Chittagong Hills, 500 feet elevation; Tenasserim and Malay Peninsula.

(Also in Hong Kong and Formosa.)

2. *DRYNARIA SPLENDENS.* (*Hook.*) Rhizome woody, the scales
 dry, fibrillose, adpressed;
 fronds subsessile, 2–3 feet long,
 root or more broad, the lower
 barren, cut down nearly
 quite to the rachis into
 obtuse entire acute lobes
 3 inches long, 1–2 inches
 broad, the upper fertile pinnæ
 distant, ligulate, 4–8 inches
 long, $\frac{1}{2}$ – $\frac{3}{4}$ inch broad; main
 veins distinct, with distinct
 transverse veinlets and copious
 intermediate ones and free
 veinlets; sori oblong or sub-
 quadrangular, filling up the
 middle of the fertile pinnæ ex-
 cept the midrib. *Hook. Sp.*
il. v. 96. Syn. Fil. 367.




DRYNARIA SPLENDENS. (*Hook.*)

edd. F. B. I. t. 316. Dryostachyum, J. Sm.

Singapore.

(Also in the Philippines.)

3. *DRYNARIA PROPINQUA.* (*Wall.*) Rhizome stout, creeping,
 the scales lanceolate-linear pubescent; fronds glabrous dimorphous,
 the barren ones 4–9 inches long, 3–4 inches broad, cut half or three-
 quarters down into bluntish or acute lobes, the fertile ones 1½–3 feet
 long, often 1 foot broad, with a distinct stem, the lobes 4–6 inches
 long, $\frac{1}{2}$ – $\frac{3}{4}$ inch broad, reaching down nearly or quite to the rachis,



A. & C. TILLACOMERAS

DRYNARIA PROPINQUA. (Wall.)

scurely serrated, texture subcoriaceous; main veins of the fertile fronds ramifying, not carried in distinct parallel lines to the margin, areoles copious, the free veinlets few and variously directed; sori in a single row near the midrib, placed at the point of union of several veinlets. *Wall. Cat.* 293. *Hook. Syn. Fil.* 367. *Bedd. F. B. I.* 60. *Pleopeltis Parishii*, *Bedd. F. B. I. t.* 125.


Himalayas, from Gurwhal to Bhotan, 2,000–7,000 feet elevation, Asia, very common; Birma and Malay Peninsula.

(Also in Java.)

4. *DRYNARIA MOLLIS*. (*Bedd.*) Rhizome creeping, clothed with subulate golden transparent ciliated scales; fronds firm-membranaceous dimorphous; sterile ones 4–6 inches long, sessile ovate or oblong, deeply pinnatifid with the sinuses very narrow and the pinnae sometimes overlapping each other, fertile fronds very short, stipitate (the dwarfed lower segments often extending nearly to the base of the rachis), ovate-lanceolate up to 1½ feet long, 2–4 inches broad, furnished with soft hairs on both sides and ciliated, the fertile fronds almost to the rachis pinnatifid, segments lanceolate 1–2 inches long, ¼–½ inch broad, nearly horizontal, rather distant, the sinus being very broad, the lower ones dwarfed and gradually reduced to a decurrent wing; venation very prominent in the sterile fronds, but much less so in the fertile ones than in *D. propinqua*; veins forming three or four series of rather regular areoles in which are sometimes included free veinlets; sori forming only a single series close to the costa, each sorus being on the vein between the first and second areole. *Bedd. F. B. I. t.* 216. Polyp. (*Drynaria*) *rivale*, *Mett.* *Hook. Syn. Fil. p.* 367, *a later name*.

Himalayas, Nynee Tal, Gurwhal, Kumaon, 6,000–9,000 feet elevation.

5. *DRYNARIA QUERCIFOLIA*. (*L.*) Rhizome creeping, short, stout, densely clothed with red-brown satiny lanceolate-subulate soft scales, which have a cordate base, and are ¼–½ inch long; fronds coriaceous or subcoriaceous of two kinds, sterile ones varying in




DRYNARIA MOLLIS. (*Bedd.*)

from 3-12 inches and more long, and 7-8 inches wide, green when very young, but soon turning dark-brown, glossy, cordate-ovate deeply lobate-pinnatifid, sometimes half-way down to the costa; fertile fronds 2-3 feet long, long-petiolate broad-ovate deeply nearly to the rachis pinnatifid, segments 5-9 inches long, 1-1½ inch wide, oblong acuminate, entire; venation manifest, costules distinct rather distant, united by transverse veins forming 4-6 primary soriferous areoles filled up with a network of small quadrangular areoles with or without free veins; sori apical small, numerous, two in each primary areole, consequently in series between and parallel to the costules. *Polypodium quercifolium*, *Lin. Sp. Pl. p. 1547.* *Hook. Sp. Fil. v. p. 96. Bedd. S. I. t. 187.*

Throughout the Indian region, in the plains, or very low down on mountains, on trees or rocks.

6. DRYNARIA LINNÆI. (Bory)
Rhizome stout, the scales 1-2 inches long from a peltate base, broadly ovate obtuse, with a deciduous acumen; fronds dimorphic the barren ones sessile, when rigid, bluntly lobed, the fertile ones long-stalked 2-3 feet long, 6-12 inches broad, cut




DRYNARIA QUERCIFOLIA. (Linn.)

cut nearly to the rachis, into entire erect-patent lanceolate lobes, texture rigid, both sides naked; main veins distinct to the edge with numerous irregular areoles between them, with copious small scattered sori. *Bory. Ann. Sc. Nat. i. v. p. 464, t. 12. Hook. Syn. Fil. 368. Bedd. F. B. I. 315.*

Ceylon; Malay Peninsula. Very like *quercifolia*, but the scales on the rhizome are quite different and the sori smaller and more scattered, it may be only a variety and is united with it by Bentham.

I have seen no specimen from South India, but without critical examination it might be passed by in the field or in the herbarium as *quercifolia*.

7. *DRYNARIA RIGIDULA*. (Sw.) Rhizome stout creeping, the



Nº192

DRYNARIA RIGIDULA. (Sw.)

the scales fibrillose dark-brown, fronds dimorphous, the barren ones sessile, 6-9 inches long, rarely more, 3-4 inches broad, cut down $\frac{1}{3}$ - $\frac{1}{2}$ towards the rachis into blunt lobes, or rarely quite pinnate at the apex, the fertile 2-4 feet long, 12-18 inches broad, long-stalked pinnate, pinnæ one inch or more apart, narrowed or stalked at the base, 6-12 inches long, $\frac{1}{4}$ - $\frac{3}{8}$ inch broad, the point acuminate, the edge more or less deeply incised, crenate, texture rather coriaceous; rachis and both surfaces naked, main veins not distinct to the edge, areoles copious, free veinlets few; sori immersed in a single row half-way between the edge and midrib. Sw. Syn. Fil. 130. Hook. Syn. Fil. 368. *Drynaria diversifolia* (R. Br.), Hook. Sp. Fil. v. 98. Bedd. F. S. I. t. 314.

Malacca.

(Also in tropical Australia, Malay Islands, the tropical Pacific Islands.)

GENUS LXV.—PLEOPELTIS. (H. & B.)

(*Pleos*, full; *peltis*, shield—the sori often furnished with round scales.)

Veins copiously anastomosing, forming copious irregular areoles, with generally free included veinlets spreading in various directions,

sori various in position, generally on the back of united veinlets ; fronds simple, pinnatifid or pinnate, articulate with the caudex.

In the *Synopsis Filicum* Phymatodes is the name adopted for this genus (or subgenus only in that work) ; this name, however, is of much later date than *Pleopeltis*, having been instituted by Presl.

In 1836, *Pleopeltis* by Humboldt and Bonpland in 1810. Some authors split the genus up into many genera.

* *Fronds simple (also 3-5 lobed in pteropus.)*

A. *Main veins not distinct to the edge.*

1. *PLEOPELTIS ACCEDENS.* (Bl.) Rhizome wide-creeping, slender filiform, scarcely paleaceous, attached to the bark of trees by numerous woolly radic'les, stipes scattered 2-5 lines long ; fronds sub-paleaceous, opaque, glabrous entire, sterile ones 1-1½ inch long, oblong-lanceolate obtuse, fertile ones 2-3½ inches long, from a broad cuneate base, long-acuminate ; venation obscure, costules indistinct, veins forming three large areoles (in each of which is one large forked veinlet) and a smaller marginal areole ; sori large for the size of the frond on the acuminate portion in a single series on each side of the costa. *Bl. En. Fil. Jav. p. 121. Hook. Sp. Fil. v. 66. Bedd. Bot. I. t. 215.*


Malacca.

(Also in the Philippines, the Malay Islands, Polynesia.)

2. *PLEOPELTIS ROSTRATA.* (Hook.) Rhizome very slender, wide-creeping, furnished with lanceolate-linear scales, stipes distant 2 inches long, naked ; fronds lanceolate-acuminate, somewhat dimorphic, the barren often being broader, coriaceous glabrous, the margin entire ; veins immersed, main veins indistinct, areoles irregular, including simple or forked free veinlets ; sori in a single row on each side close to the midrib. *Hook. Syn. Fil. 353. Bedd. F. B. I. 159.*

Sikkim and Bhotan, 6,000 feet elevation, abundant ; Khasya, 4,000-5,000 feet elevation.

3. *PLEOPELTIS LINEARIS*. (*Thunb.*) Rhizome wide-creeping, woody, the scales lanceolate dull-brown, never with hairs, stipes from a few lines to 2 inches long, rarely with any scales; fronds 6-12 inches long, rarely more, $\frac{1}{4}$ - $\frac{1}{2}$ inch broad, entire, narrowed gradually at both ends, texture coriaceous, both sides naked or nearly so; areoles



N°193

PLEOPELTIS ROSTRATA. (*Hook*)

irregular with forked free veinlets; sori superficial or scarcely immersed, never sunk in a cyst, large prominent, forming a single row half-way between the costa and margin, when young covered with copious thin peltate scales. *Thunb. Fl. Jap.* 335. *Hook. Syn. Fil.* 354. Polyp. loriforme, *Wall. Cat.* 271, 2nd sheet (typical) *P. Wightianum*, *Wall. Cat.* 2222, acutissimum, 4727, gladiatum, 279.

Peltis nuda, *Hook. Exot. Fl.* t. 63. *P. Wightianum*, *Bedd. J.* 180. (*P. loriforme*, *Wall. Cat.* 271, *type-sheet*, belongs I believe, and not to *simplex*, it has larger fronds, but they are low and coriaceous and the veins hidden.)

Himalayas, 1,000–10,000 feet elevation, very common, *Sya*; throughout South India; Ceylon; Malay Peninsula.

(Also in the Malay Islands, China, Japan, Central and South America and its Islands.)

VAR. STENISTE. (*Clarke.*) With the fronds only 1–1½ lines wide and the sori projecting beyond the margin, but always superficial, only a form of *linearis*, and it graduates into the type, his “*polymorpha*” is an abnormal form of *linearis* with the fronds pinnate (such as occurs in many ferns.)

4. PLEOPELTIS SIMPLEX. (*Sw.*)

Like *linearis*, only the fronds are thinner in texture and much larger, to 2 feet long by 1–2 inches wide, the veins more distinct, sori always more or less deeply sunken and forming pustules on the upper side of the fronds, sometimes



Nº 194

PLEOPELTIS LINEARIS. (*Thunb.*)

are sunk in a cyst with visible margins, but never nearly so prominent as in *stenophylla*, which also differs in its smaller size, much more rigid texture and fewer veins. *Sw. Syn. Fil.* 27. *Polyp. avatum* (*Bory.*), *Willd. Sp. Pl.* v. 158. *P. Grevilleanum*, *Wall. f.* 5169. *P. sesquipedale*, *Wall. Cat.* 275. *P. lineare*, var. β *simplex*, *Hook, Syn. Fil.* 354.

Himalayas, confounded by *Hooker* (in his *Sp. Fil.*) and by *Clarke* with *linearis*, but kept distinct by all other botanists, and

made a distinct variety of linearis by Mr. Baker in the *Syn. Fil.*, but Mr. Baker now agrees that it should be made a distinct species; the scales of the rhizome differ from those of "normalis," but the fronds are almost exactly similar to that species, and I had previously confounded it with that (not with linearis.)

5. PLEOPELTIS CLATHRATA. (*Clarke.*) Rhizome short creeping, with ovate acute, often hair-pointed scales; fronds small, stipe often as long as the frond; sori mixed with sessile irregularly peltate and lacerate clathrate scales. *Clarke, F. N. I. p. 559, t. 82, fig. 1.*

Kashmir, Pir Pinjul, 11,000–12,000 feet elevation.

(Also in Afghanistan.)

Mr. Clarke thinks it may be high level form of linearis, but that the scales are peculiar, the texture thinner, the venation consequently more conspicuous, and the stipe longer.


6. PLEOPELTIS STENOPHYLLA. (*Bl.*) Rhizome creeping, paleaceous, with glossy subulate pale imbricated ferruginous scales, stipes 1–2 inches long, distant; fronds firm-coriaceous, very rigid, glabrous polished, 4–9 inches long, $\frac{1}{4}$ – $\frac{1}{2}$ inch, or rarely more wide, linear-oblong obtuse, the margin thickened and subreflexed, broadly crenate-sinuate, rarely narrowing upwards, but much and gradually attenuated below on the stipes; costa prominent beneath, costules none, veins anastomosing so as to form oblong, hexagonal oblique areoles, generally quite empty or with rarely a free included veinlet; sori rather distant oval, arranged in a series close to the margin, deeply sunk in a cyst, forming elevated pustules on the upper side one to each crenature, the margin of the cavity elevated. *Bl. Fil. Jav. p. 135, 55, f. 1. Hook. Sp. Fil. v. 65. Mett. Polyp. p. 99, t. 1, f. 31–32. venation. Bedd. F. B. I. 234.*

Mr. Clarke states that this comes between "linearis" type and his variety "steniste," but both these ferns have different venation, and neither have the sori sunk in cysts; this has the sori always much nearer the margin than the midrib even in the broader forms.

Malacca.

(Also in the Philippines.)

7. *PLEOPELTIS SINUOSA*. (*Wall.*) Rhizome much branched, creeping or scandent on trees, sparingly radiculose, half an inch to two inches thick, almost woody when dry, carnosose when recent, added, as it were, with large conical processes upon which the stipes are or have been articulated, and which appear to increase much in size after the fronds have fallen, the whole densely covered (as it is tessellated) with nearly orbicular appressed whitish peltate scales with a dark spot in the centre, stipes 1-2 inches long, glabrous and scaleless; fronds sub-dimorphic, subcoriaceous glabrous, sterile ones 3-6-7 inches long, 1-1½ inch broad, subelliptical, oblong obtuse entire, fertile ones larger, sometimes 1 foot - 16 inches long, ½-¾ of an inch wide, ear-oblong obtuse, the margins more or less sinuated, the base both attenuate-decurrent; venation copiously anastomosing, forming large very irregular areoles filled with lesser ones (formed by more slender veins), and these include branched or simple veins which occasionally again anastomose; sori remote, large, long, immersed in cysts, which form pustules on the upper side arranged in a single series nearer the margin than the costa. *Wall.*



Nº 185


PLEOPELTIS SINUOSA. (*Wall.*)

Bot. n. 2231. Hook Sp. Fil. v. 61. Bedd. F. B. I. t. 8.
 Malacca.

(Also in the Solomon Islands, Amboyna and New Hebrides.)

8. *PLEOPELTIS LONGIFOLIA*. (*Mett.*) Rhizome horizontal, creeping, thick squamose, stipes approximate, 2-3 inches long; fronds 1-2-3 feet long, ½-1½ inch wide, thick, carnosose-coriaceous,

glabrous, linear-lanceolate, obtusely acuminate, gradually long attenuated below on the stipes, entire, the margins subrevolute, venation internal, very indistinct, costules veniform, united by transverse veins



Nº196.

PLEOPELTIS LONGIFOLIA. (Mett.)

gradually at both ends, the margins entire, 6-12 inches long, by 1-2 inches broad, glabrous; areoles numerous with free veinlets; sori

into rather large areoles then into irregular lesser ones, which include free veinlets; sori corripital oblong, forming a line or series near the margin rather close placed, sunk in cysts, and forming a corresponding not elevated line on the upper surface, capsules mixed with long stipitate scales. *Mett. Polyp. p. 87. Hook. Sp. Fil. v. 60. Bidd. F. B. I. t. 7. P. contiguum, Wall. Cat. 285.*

Birma and the Mal Peninsula generally, Merg Singapore, &c.; North India Kumaon.

(Also in the Philippines and Malay Islands.)

9. PLEOPELTIS SUPERFICIALIS. (Bl.) Rhizome with creeping, thick as a crow-qu the scales ovate and lanceolate, acute, spreading, brown shining; stipes up to six inches long, often curved, sometimes with ovate scales; fronds lanceolate-linear (often forked at the apex), subcoriaceous, narrowed

copious scattered irregularly, compital, large or small. *Polypodium superficiale*, *Bl. En. Pl. Jav. Fil.* 123. *Hook. Syn. Fil.* 355. *Bedd. F. B. I. t.* 75.

Khasya Hills, 4,000–6,000 feet elevation, extending to the Naga country to the east, and to Bhotan and Mishmee; Malay Peninsula.


(Also in South China.)

10. *PLEOPELTIS LANCEOLATA*. (L.) Rhizome long-creeping, paleaceous, with lanceolate ferruginous scales, stipes remote, 1–2–4 inches long; fronds coriaceous, 3–9 inches long, $\frac{1}{4}$ – $\frac{3}{4}$ inch wide, lanceolate, more or less acuminate, long and gradually attenuated at the base, copiously furnished with orbicular ovate, small appressed peltate scales dark in the centre, pale in the circumference and denticulate; veins immersed indistinct, the primary veins form large obliquely elongated areoles, which include very irregular and different sized areoles, and a few free veinlets which are rarely forked; sori generally very large and often exceedingly prominent, pulvinate globose or oval, stalked scales mixed with the spore cases. *Polyp. lanceolatum*, *Linn. Sp. Pl.* 1082. *P. lepidota*, *Hook. Sp. Fil.* v. 56. *Willd. in Schlecht Adunbr.*, p. 17. *Bedd. F. S. I. t.* 181. *P. angustata* var. *depauperata*, *Clarke, F. N. I. p.* 559.

Nilgiris and higher mountains on the West side of the Madras Presidency; Assam (one poor specimen in Kew Herbarium); Ceylon, Ambawalla estate.

(Also in Tropical America and the West Indies, South Africa and its islands; St. Helena; Sandwich Islands.)

11. *PLEOPELTIS ANGUSTATA*. (Sw.) Rhizome long-creeping, branched, paleaceous with falcate subulate-setaceous scales, stipes remote, 1½–4 inches long; fronds 5 inches to a span or more long ½–2 inches wide, tapering below into a petiole, glabrous above, clothed beneath with cottony tomentum which is deciduous, coriaceous, oblong-lanceolate; venation sunk obscure, costular areoles with free or branched and more or less connected veinlets; sori very large, subglobose oval, partially sunk in the frond, very convex,



g a single series on each side between the costa and margin, lly towards the apex only, sometimes longitudinally confluent. *en. Fil. pp. 27 and 224.* *Niphobolus angustatus, Hook. Sp. 43. Bedd. F. S. I. t. 185.* *Niphopsis, J. Sm.* *Pleopeltis ora, Presl.* *Polypodium sphærocephalum, Wall. Cat. n. 272.* Malay Peninsula, Singapore, Malacca, and Penang. South ?) I have never detected it, though I have a specimen said to een collected on the Nilgiris, and Hooker gives Tranquebar as ity.

Also in the Malay Islands and North-east Australia.)

2. *PLEOPELTIS NORMALIS. (Don.)* Rhizome scandent, thick owquill, with ovate or lanceolate almost hair-pointed adpressed which carry erect from their backs tufts of linear stiff black-red ; fronds narrowed much at the base, sometimes to the very the stipe, usually broadest near the middle, 1-2 feet long, by ches broad, submembranaceous glabrous, a few scattered ovate dpressed scales often present on the stipes and near the base main rachis beneath ; venation as in linearis, main veins very ict and areoles with free included veinlets ; sori large, super- r a little immersed, often in several rows, or irregularly in one : rarely regular in one row. *Don. Prod. Fl. Nep. 1. Clarke, t. 558. Hook. Syn. Fil. 358, in part. Bedd. F. B. I. t. 10. gifronds, Woll. Cat. 274.*

Himalayas, Nepal to Bhotan, 4,000-8,000 feet elevation, a, 3,000-5,000 feet ; Tenasserim on Mooleyit 5,000 feet on.


B. *Main veins distinct to the edge, or nearly to the edge.*

3. *PLEOPELTIS RHYNCOPHYLLA. (Hook.)* Rhizome creeping, r, paleaceous, subsquarrose with subulate ferruginous scales, scattered, slender, 1-2 inches long of the sterile frond, 2-4 of tile one ; fronds firm, subcoriaceous, glossy, slightly thickened, ly crenate, sterile ones 1-2 inches long, elliptical, or subovate, , fertile ones 3-6 inches long, lanceolate from near the base,

gradually and long-acuminated, below attenuated; venation distinct, slightly prominent, the costules connected by transverse veins forming areoles which enclose free veinlets, which latter sometimes

unite with opposite veins; the sori mostly confined to the narrow acuminate apex, and when mature very large in a single series on each side of the costa. *Hook. Sp. Fil.* v. 65. *Hook. Ic. Plant.* 655 (or t. 55 of *Cent. of Ferns.*) *Bedd. F. B. I.* t. 9.

Khasya, 4,000–5,500 feet elevation; Tenasserim on Mooleyit, 5,000 feet elevation.



No 198

PLEOPELTIS RHYNCHOPHYLLA.

(Hook.)

feet elevation. Khasya, 4,000–5,000 feet.

14. PLEOPELTIS GRIFFITHIANA. (Hook.) Rhizome wide-creeping, as thick as a crowquill, the scales dense, lanceolate-subulate, brown, spreading; stipes 3–6 inches long, firm, erect; fronds 6–8 inches long, up to 2½ inches broad, coriaceous, lanceolate or ovate-lanceolate, the apex acute, the margin more or less repand, the base rounded or suddenly narrowed; main veins very distinct to the edge, areoles fine, hidden with copious free simple or forked veinlets; sori large, one between each main vein in a single row close to the costa, often extending more than half-way to the base of the frond. *Hook. Syn. Fil.* p. 359. *Bedd. F. B. I.* t. 158.

Sikkim and Bhotan, 6,000–9,000

15. PLEOPELTIS OVATA. (Wall.) Rhizome wide-creeping,


like as a crowquill, scales lanceolate-linear, brown spreading; stipes 1-2 inches long, firm, erect, slightly scaly below; fronds 6-12 inches long, up to 3 inches broad, ovate-lanceolate, the apex acute, the margin entire, the base narrowed suddenly, texture herbaceous, both sides glabrous; main veins distinct to nearly the edge, with copious areoles including free forked veinlets; sori large irregularly scattered on the costa. *Polypodium ovatum*, Wall. Cat. 276, Hook. Syn. 359. Bedd. F. B. I. t. 157.

Sikkim, Chundaghiri, Bhotan, Assam, 4,000 feet elevation, at Hop's Falls, Shillong, rare.

16. *PLEOPELTIS MEMBRANACEA*.


(Don.) Rhizome short-creeping, but the younger portion paleaceous, with blackish-green ovate acuminate scales; stipes distant or aggregated, 1-5 inches long, if younger winged above with the decurrent base of the frond; fronds 1-2 inches to 2-3 feet long, by less than an inch to 6 inches broad, thin membranaceous and translucent, lanceolate or oblong-lanceolate, or oblanceolate, acuminate, the base long attenuated and much gradually decurrent upon the stipes,

glabrous, the margin entire (or rarely more or less deeply sinuate-dentate or pinnatifid, and even fimbriated throughout the whole length with long narrow unequal segments); venation very distinct, main veins horizontal or nearly so, slender, usually remote, connected by transverse veins which form the primary areoles, and these are filled up with irregular net-work, of which the areoles are very unequal and include copious free veinlets; sori compital, rather small, usually in two series between the costules, more or less numerous according



Nº 199

PLEOPELTIS GRIFFITHIANA. (Hook.)



width of the frond, rarely reduced to one sorus near the costa, a single series parallel with the costa is formed, not unfrequently there are 3-4-5 series (not very regular) between the costa, in instance a single series only appears between the costules, and of a very large size and abnormal in form, often oval or oblong (being parallel with the costules). *Don. Prod. Fl. Nep. p. 2.* *Sp. Fil. v. 70.* Polyp. grandifolium, *Wall. Cat. n. 282.* heterocarpum, *Bl. Fil. Jav. p. 167, t. 75.* *Bedd. F. S. I. t. 177.* Himalayas, from Gurwhal and Bhotan, 3,000-8,000 feet elevation; Khasya, 2,000-5,000 feet, Chota Nagpore, Parasnath; India, mountainous tracts on both sides of the Presidency, 5,000 feet elevation; fronds varying from $\frac{1}{2}$ inch to 5 inches in length; Ceylon. It only appears with the rainy season and dies in November.


7. PLEOPELTIS ZIPPELLII. (*Bl.*) Rhizome slender, creeping; scales lanceolate acute; fronds lanceolate, often quite entire at the apex, narrowed gradually at the base into a short stipe (rarely there is a stipe about 2 inches long), subcoriaceous, entire or a little undulate, lower part of the rachis beneath covered with a few scales; primary veins prominent and near the base, areoles copious, with free included simple or forked veinlets; compital in two rows between the primary veins and in about two rows between the costa and margin. *Bl. Pl. Jav. Fil. 172, t. 80.* *Sp. Fil. v. 72.* Pleopeltis heterocarpa, *Hook. Syn. Fil. p. 360, (l.)* *Bedd. F. B. I. 319, (venation not correct, as not showing the veinlets.)*

Himalayas, Dalhousie (Dyas), Sikkim and Bhotan, 2,000-4,000 feet elevation, Khasya, 2,000-4,000 feet elevation, (not in Java as stated by Clarke).

Also in Java and Philippines.)

8. PLEOPELTIS PUNCTATA. (*Linn. under Acrostichum.*) Rhizome creeping, stout, clothed with brownish ovate or lanceolate scales; fronds 1-3 feet long, 1-3 inches broad, lanceolate, sessile,

gradually narrowed at the base, acute or blunt at the apex, coriaceous, very glabrous; venation immersed (visible in a dried state), main veins fine but distinct to the margin, areoles copious, including smaller areoles, in which are free simple or forked veinlets, with clavate apices; sori compital, very small and scattered very irregularly, generally only on the upper half of the frond. *Linn. Sp. Pl.* 1524. *Sw. in Schr. Jour.* 1800, ii. 21. *Pleopeltis irioides*, *Hook. Syn. Fil.* 360. *Bedd. F. S. I. t.* 178. *Wall. Cat.* 281, glabrum, and 273, polycephalum.



Nº 201.

PLEOPELTIS PUNCTATA.
(*Lin.*)

Bedd. F. B. I. t. 317.

Malacca.

(Also in the Malay Islands.)

South India, Western mountains up to about 3,000 feet elevation; Ceylon; North India, in the Bengal plains, and up to 3,000 feet elevation; Malay Peninsula.

(Also in China, Malay Islands, North Australia, Polynesia, Southern and Central Africa and the islands.

19. PLEOPELTIS MUSÆFOLIA. (*Bl.*)

In every way like punctata, so that the same description will do for both, except that the texture in this is, like that of nigrescens and hemionitidea, thin and papery, with the veins very prominent, whilst punctata has fleshy leaves, in texture like *Thamnopteris Nidus*. *Bl. Fil Jav. p.* 171, t. 79 *Hook. Syn. Fil.* 360

20. PLEOPELTIS HEMIONITIDEA. (*Wall.*) Rhizome creeping, scaly, furnished with wiry scaly fibrous roots; stipes scaly, winged upwards; fronds 6 inches to 1½ foot long, 1–2 inches broad, membranaceous subchartaceous, subtransparent very dark shining green, broad lanceolate, gradually acuminate into a fine point at the apex.


and gradually attenuated at the base, being winged nearly all down the stipe, entire glabrous, costa slightly scaly beneath near the base; venation very prominent, main veins blackish, pinnate nearly horizontal, areoles in about five series, subquadrate, in which are free veinlets which are either simple or forked, but without clavate apices; sori compital, rather large, forming one series between the main veins very irregular in shape and size, subglobose or oblong, or even (by confluence) linear. *Wall. Cat. p. 284. Hook. Sp. Fil. v. 73. Bedd. F. B. I. t. 182.*

South India, on the Western mountains, rather common in Coorg, 3,000 feet elevation, rare elsewhere—Himalayas, Nepal, Bhotan, 2,000–7,000 feet elevation, Khasya, 1,000–5,000 feet elevation, Chittagong, 1,000 feet elevation.


(Also in South China and Malay Islands.)

21. *PLEOPELTIS PTEROPUS.*

(Bl.) Aquatic, rhizome creeping, branched, the young apices only paleaceous, with blackish lanceolate-subulate scales; stipes more or less apart, 1–3–4 inches to 1 foot long, winged upwards, and as well as the back of the costa and costules furfuraceous-squamose; fronds 2–3–9 inches long, $\frac{1}{4}$ –1 $\frac{1}{2}$ and 2 inches wide, firm membranaceous, very dark dirty green (when dry often black), lanceolate acuminate entire, or 3-lobate, or 3-partite, or 5-lobed, terminal lobes up to 8 inches long, lateral ones 3–5 inches, below long-tapering into a gradually decurrent wing upon the petiole, glabrous above, margin entire; venation very conspicuous, main veins prominent beneath, rather wide apart, extending about two-thirds of the way to the margin, then uniting and forming large costal



PLEOPELTIS HEMIONITIDEA. (Wall.)




areoles, within which the sori have their origin, a second series of smaller areoles is formed nearer the margin, and these and the rest of the frond are filled up with a net-work of smaller irregular areoles, including free simple or forked veinlets which have clavate apices; sori small not very numerous, 1-3 in each large areole, compital upon the secondary veins of the primary areole, often confluent into transverse, oblong or linear (grammitoid) sori. *Bl. Fl. Jav. Fil.* 168, t. 76. *Polypodium tri-dactylon*, *Wall. Cat. n.* 315. *Hook. Sp. Fil.* v. 75. *Hook. et Grav. Ic. Fil.* t. 209. *Bedd. F. B. I. t.* 11.

North India, Sikkim and Bhotan, 1,000-4,000 feet elevation; Khasya from no elevation up to 4,000 feet, in the plains at Mymensingh, Chittagong, plains up to 1,000 feet; Malay Peninsula.

(Also in the Philip-pines and South China.)

VAR. MINOR. Fronds always small and simple 3-4 inches long, by $\frac{1}{4}$ - $\frac{1}{2}$ inch broad, never lobed. *Bedd. F. S. I. t.* 179.

South India, Anamallays and Bolampatty Valley, 3,000-4,000 feet elevation, in rivers on rocks under water, fructifying when the water subsides after the rainy season; Ceylon. This may not be entitled to rank as a permanent variety, as Mr. Clarke says the North



Nº 204.

PLEOPELTIS PTEROPUS. VAR. MINOR.
(Bl.)

Indian plant is small and simple at the higher elevations; but I never saw 3-lobed examples in South India or Ceylon, or fronds more than about 4 inches long.

VAR. *ZOSTERÆFORMIS*. (*Wall. Cat.* 280.) Fronds very narrow and ribbon-like, 4-7 inches long, by 1-3 lines broad. *Hook. Syn. Fil.* 357. *Bedd. F. B. I. t.* 123. Baker is wrong in saying there are no free veinlets; it differs from the last in no way except in the narrowness of the fronds, and it is certainly only a local variety.

Tenasserim, beds of rivers at the foot of Mooleyit.

** *Fronds deeply pinnatifid.*



Nº 205.

PLEOPELTIS HASTATA. (*Thunb.*)

22. *PLEOPELTIS HASTATA*. (*Thunb.*) Rhizome creeping, stout, paleaceous with dense subulate or hair-pointed falcate ferruginous scales; stipes 2-5 inches long; fronds coriaceous, 2-12 inches and more long, deltoid-ovate acuminate, trifid or deeply pinnatifid to within half an inch of the rachis, with 5-11 segments, which are 3-6 inches long, $\frac{1}{2}$ to 1 inch wide, very patent from a broad base oblong-lanceolate, very finely acuminate entire or repand or denticulate, thickened at the margin, the lowest ones cuneate-decurrent, terminal segment often the longest and most narrowly

acuminate; venation conspicuous, main veins often prominent beneath, distant, united by transverse veins forming 3-4 series of primary areoles, which are filled up by lesser ones including free veinlets; sori large not sunk, arranged in a single series nearer the costa than the margin. *Thunb. Fl. Jap.* 335. *Polypodium oxylobum*. *Wall. Cat.* 294. *Bedd. F. S. I. t.* 175. *Pleopeltis trifida*, *Des.*

Prod. Fl. Nep. 3. *Hook. Syn. Fil.* 363. *Pleopeltis malacodon*, *Bedd. Sup. t.* 387.

Young rhizomes often produce only very small fronds, which are simple or trilobate only; (var. *Thunbergii*, *Clarke*). This fern can at once be known from *malacodon* by the base of the lowest pair of pinnæ (*i.e.*, base of the frond) being always a little decurrent at the apex of the stipe, (not scooped out and subcordate) and in being subentire or only a little denticulate instead of acutely serrulate; Mr. Clarke, however, is wrong in saying that it is always quite entire, my specimens (looked at under a lens) are always more or less toothed.

Throughout the Indian region on the mountains; in South India between 5,000–8,600 feet elevation; in the North 2,000 to 5,000 feet.

23. *PLEOPELTIS MALACODON*. (*Hook.*) In all respects like *hastata*, only that the base of the frond is scooped out and subcordate instead of being a little decurrent), and the margins are acutely serrulate, the teeth mucronate or spinescent. *Hook. Sp. Fil.* v. 87.

Stewartii, *Clarke, F. N. I. p.* 563, *a variety less serrated*. Mr. Baker and Mr. Clarke both protest against this being considered a variety only of *hastata*, otherwise I should have so placed it, the differences being those of a variety rather than of a species.

Himalayas, Nepal to Bhotan, 10,000–13,000 feet elevation.

VAR. β MAJUS. (*Hook. Sp. Fil.* v. 88.) Base of the frond cordate or sometimes decurrent; pinnæ (or rather segments) narrower and longer, and much more erect (*i.e.* pointing upwards); margins less prominently serrated, scales of rhizome blacker. *Pleopeltis Stewartii*, *Bedd. F. B. I. t.* 204, and *Baker, Syn. Fil.* 513 (*not Clarke*). *Pol. propinquum*, var. *Wall.* *P. cryptolobum*, *J. Smith, MS.* *Clarke, F. N. I. t.* 83.

Nepal to Bhotan, 9,000–12,000 feet elevation (pendant from trees). I think that Sir W. Hooker may be safely followed in considering this only a variety of *malacodon*.

24. *PLEOPELTIS EBENIPES*. (*Hook.*) Rhizome stout creeping,

lowest pair generally deflexed, and entire
 stipe, or scooped out and subcordate at
 patent as are the other pinnæ, all a little



№206.

PLEOPELTIS INCURVATA. (Bl.)

though closely allied to *Pleopeltis*

gl
 w
 lo
 so
 ve
 are
 let
 the
 v.
 I
 var
 typ
 a I
 mei
 dor
 a s
 with
 ebe

of the sterile frond, 2 feet of the fertile, glossy brown ; fronds firm-coriaceous glossy dimorphous, sterile ones 6-10 inches long, 8 inches to 1 foot wide, hastate, tripartite, simply ovate or pinnatifid with 4-6 lateral lobes, lobes ovate-acuminate entire horizontal, terminal one very large, all thickened at the margin ; main veins distinct, but immersed, other veins internal and very indistinct ; areoles with free included simple or forked veinlets ; fertile fronds very long-stipitate, 10 inches to a foot long, nearly as much wide, very broad-ovate deeply pinnatifid nearly to the rachis, below subpinnate, segments 5-15, 4-8 inches long, very remote, linear and acuminate, decurrent at the base ; sori large, copious, oval, sunk into a very deep cavity which forms a circular elevated truncated tubercle on the opposite side, occupying the space between the costa and the margin in a single series. *Bl. Fil. Jav. p. 151, t. 65. Hook. Sp. Fil. v. 77. Bedd. F. B. I. 124.*

Malacca.

(Also in the Malay Islands.)



Nº 207.

PLEOPELTIS INSIGNIS. (Bl.)

26. *PLEOPELTIS INSIGNIS*.
(Bl.) Rhizome short-creeping, paleaceous, with deciduous ferruginous scales ; stipes 1-2 inches or more long (according to the length of the decurrent wing of the frond) ; fronds subtriangular-ovate, membranaceous, glabrous, 4-9 inches long (independent of the decurrent wing), and nearly as much broad, subdeltoid, deeply pinnatifid, with 5-9 lanceolate spreading acuminate segments ; the base of the fronds subcuneate and decurrently attenuated so that the stipes is winged for the

Malacca.

(Also in the Philippines and J

27. PLEOPELTIS PHYMATODES.

ing, woody, the scales dark brown f
firm, erect, glossy; fronds varying
6 inches long, to deeply pinnatifid
entire, acuminate, lanceolate oblon
broad, texture coriaceous, both sides
areoles fine with copious free vein
immersed, 1-2 serial or scattered.

Syn. Fil. 364. *Bedd. F. S. I.* 17
Cat. 289, *type sheet.*

Ceylon; Malabar and Travance
Peninsula.

(Also throughout the tropics of
India excepted.)

28. PLEOPELTIS LONGISSIMA. (

the scales ovate adpressed; stipes 3-4 f
fronds 1-4 feet long, 6-12 inches broa
rachis (often leaving only the narrow
erect ...)

Assam, Gowhatty, Sylhet, Furidpore (floating in jheels.)
(Also in the Philippines, Formosa, and Malay Islands.)

29. *PLEOPELTIS NIGRESCENS*. (Bl.) Very like *longissima*, but wing to the rachis broader and the pinnæ broader ($1-1\frac{1}{2}$ inches), ~~er~~ more crisp in texture and the veins more prominent; it is ~~aps~~ only a variety, but I have never seen *longissima* in a wild

~~2.~~ *Bl. En. Fil. Jav.* 127.

Longissima, *Bedd. F. S. I.*


Wall. Cat. 289, 2nd sheet
(*rnifolium*).

South India, on rocks, ~~ut~~ at the foot of the Western ~~ntains~~ mountains, (foot of Sispara Carcoor ghats), not ascend- the mountains to any ele- ~~on~~ on, and never growing in ~~er~~ er; Ceylon. An exceed- y handsome fern.

30. *PLEOPELTIS DILATATA*.

(all.) Rhizome stout creep-
- paleaceous with ovate
- culated scales; stipes $1\frac{1}{2}$
- tand more long, but strongly
- aged the whole length by

- decurrent base of the frond; fronds ample $1\frac{1}{2}-2$ and 3 feet long,
- ot and more wide, membranaceous, light-green, glabrous, oblong-
- ate, pinnatifid to within $\frac{1}{2}-1$ inch of the rachis (less towards the base),
- ments 6-11, 5-8 inches long (shorter towards the apex), $\frac{5}{4}-1\frac{1}{2}$ inch
- le, oblong-lanceolate acuminate, subentire, suddenly decurrent;
- in veins wavy, forming large primary areoles which extend two-
- rds of the way to the margin and include the sori, these and the rest
- the segments are occupied by lesser irregular areoles including
- ny free veinlets, their branches moderately divaricating; sori nu-
- rous, very small, often oval or more or less elongated, compital on



PLEOPELTIS NIGRESCENS. (Bl.)

the secondary veins and veinlets. *Wall. Cat. n. 295. Hook. Sp. Fil. v. 85. Bedd. F. B. I. t. 122.*

North India, from Nepal to Bhotan, 3,000–8,000 feet elevation, Khasya, 2,000–6,000 feet; Ceylon, forests above Telgamma, 4,000 feet elevation; Tenasserim and Malay Peninsula.

(Also in Samoa.)

*** *Fronde pinnate.*

31. *PLEOPELTIS PALMATA.* (*Bl.*)



N°209.

PLEOPELTIS DILATATA. (*Wall.*)


including free incurved clavate veinlets; sori rather large in a single row, midway between the costa and margin. *Bl. Fil. Jav. p. 150, t. 64. Hook. Syn. Fil. 368. Bedd. F. B. I. t. 156.*

Malay Peninsula, Singapore, Penang and Malacca.

(Also in the Malay Islands and Philippines.)

32. *PLEOPELTIS JUGLANDIFOLIA.* (*Don*). Rhizome stout creeping, with many spreading lanceolate subulate bright-ferruginous scales; stipes 1 foot or more long, firm, erect, glossy; fronds 1½–2 feet long, 1 foot or more broad; pinnae 8–10 on a side in pairs, which are 1–2

Rhizome stout, the scales large lanceolate; stipes 6–12 inches long, firm, erect, glossy; fronds 6–18 inches long, 8–12 inches broad, with a linear or linear-oblong entire or slightly toothed or repand terminal lobe, and 1–6 similar ones on each side, which are ½–1½ inch broad, narrowed or dilated, always slightly adnate at the base, long caudate at the apex, those of the barren frond the broadest, texture subcoriaceous, both sides glabrous; main veins distinct to the edge, areoles fine, hidden,



PLEOPELTIS JUGLANDIFOLIA. (Don.)

inches apart, not quite opposite, articulated with the rachis, 1-1½ inch broad, the apex caudate; the margin thickened and wavy, rarely lobed, sessile from a rounded base, or attenuated and petioled, texture subcoriaceous, both sides glabrous; main veins distinct to the margin; areoles copious, hidden, including free veinlets; sori large, one between each main vein, forming a single row, much nearer the midrib than the margin (rarely in two rows). *Don. Prod. Fl. Nep. 3. Hook. Syn. Fil. 368. P. capitellata, Wall. Cat. 306. Bedd. F. B. I. t. 12.*

VAR. TENUICAUDA. (*Hook.*) Pinnæ gradually narrowed and acute at the base and petiolate, quite runs into the type and cannot be considered a variety. *P. leiorhizon, Wall. Cat. 303, 3rd sheet.*


Himalayas, from Gurwhal to Bhotan, 2,000-9,000 feet elevation, very common, Khasya, 2,000-5,000 feet.

33. PLEOPELTIS LEHMANNI. (*Mett.*) Rhizome creeping, clothed with ovate or orbicular glabrous scales, each ending in a very long acumination; stipes 8 inches or more long, glabrous as well as the rachis and frond; fronds 1½-2 feet long, submembranaceous pinnate; pinnæ opposite or subopposite, sessile, articulated at the base (terminal one long petioled), linear oblong entire, with a caudate acumination 4-6 inches long, about ¾ inch broad, superior base obliquely excised, inferior base amplexicaule, the margin entire or slightly repand; main veins evident not extending quite to the margin, connected by the transverse veins which form about 4 series of areoles in which are free simple or forked veinlets with clavate apices; sori irregular, generally 4-6 in two rows between the main veins. *Mett. Poly. p. 229. Bedd. F. B. I. t. 260. Hook. Syn. Fil. p. 369.*

Sikkim, 4,000-8,000 feet elevation; Birma.

(*Pleopeltis moulmeinensis, Bedd. F. B. I. t. 205*, drawn from a single specimen in Mr. Parish's possession, is probably this, imperfectly seeding; it does not agree in habit with *juglandifolia*.)

34. PLEOPELTIS HIMALAYENSIS. (*Hook.*) Rhizome glaucous



10° 211.

PLEOPELTIS LEHMANNI. (Metz.)

creeping, clothed with bright ferruginous hair-like scales; stipes a span to 1 foot and more long; fronds 1-1½ foot long, submembranaceous, sometimes pubescent or even subtomentose, glabrous subcoriaceous in age, pinnated, pinnæ almost invariably opposite, sessile in distant pairs 5-8-9 inches long, by 1½-2 inches broad from an obtuse base, elliptical-oblong, finely and long-caudately acuminate entire, but with a very distinct membranous hyaline margin; main veins manifest, connected by transverse ones forming areoles, of which one or all are soriferous, these areoles are filled up with a net-



N°212

PLEOPELTIS HIMALAYENSIS. (Hook.)

work of lesser areoles which have free included veinlets; sori often large, 1-2 in the primary areoles, hence they are 1-2 serial between the main veins, and 3-4 serial in a direction parallel with the costa. *Hook. Sp. Fil.* v. 91. *Bedd. F. B. I. t.* 318. *Polypodium venustum*, *Wall. Cat.* 305, not of *Desr.*

Himalayas, 6,000-10,000 feet elevation, from Nepal to Bhotan, Khasya Hills (Jerdon), young plants often have simple or 3-lobed frond. Mettenius considers it on a variety of *Lehmanni*.

Clarke calls the tomentose variety, var. *niphoboloides*, but it graduates into the type.

35. *PLEOPELTIS LEIORHIZA.* (*Wall.*) Rhizome very thick fleshy, wide-creeping, covered with ovate adpressed peltate, slaty-brown scales; stipes 1-2 feet long, erect; fronds 2-4 feet long, 1-2 feet broad; pinnæ narrow lanceolate-linear coriaceous acuminate at the apex, margins entire, sessile and rather decurrent at the base, or

venuated and petioled, both sides glabrous; main veins scarcely more prominent than the rest, areoles with copious free veinlets; sori large, in a single row, a little nearer the midrib than the margin. Polypodium, *Wall. Cat.* 303, *type-sheet.* *Hook. Syn. Fil.* 369. *Bedd. S. I.* 174.

South India, common on all the Western mountains, 2,000–4,000 feet, on rocks; North India, from no elevation up to 4,000 feet.

TRIBE II.—GRAMMITIDÆ.


Sori on the back of the lobes, more than twice as long as broad, usually linear.

GENUS LXVI.—NOTHOLÆNA. (*R. Br.*)

Nothos, spurious; *lænos*, wool—the scales on back of frond pseudo-woolly.)

Sori marginal, at first oblong or roundish, soon confluent into a continuous marginal line, without a distinct involucre, but with the edge of the frond frequently inflexed; veins free; fronds adherent to the caudex, pinnate or bipinnate.

1. NOTHOLÆNA MARANTÆ. (*L. under Acrostichum.*) Rhizome stout, horizontal, densely paleaceous with soft silky ferruginous finely pointed subulate scales; stipes generally stout purplish-black, 3–10 inches long, aggregated, setosely and densely hirsute, as well as the fronds; fronds 4–10 inches long, 2–2½ inches wide, oblong-lanceolate, paleaceous, glabrous above, beneath densely clothed with ferruginous long-lanceolate imbricated scales, bipinnate, primary pinnae petiole or sessile from a broadish obtuse base, oblong-acuminate, sori numerous, approximate, sessile, 2–3 lines long, very obtuse entire, upper ones confluent at the base (as are the terminal primary pinnae), the edge scarcely reflexed; sori forming a broad border extending some way from the margin towards the costule.



A. S. CATTELL & C^O ENG^S.

NOTHOLAENA MARANTAE. (Linn.)

much concealed by the paleaceous covering. *Linn. Sp. Pl.* 1527. *Notholæna Marantæ*, *R. Br.* *Nothochlæna*, *Hook. Sp. Fil.* v. 120. *Bedd. F. B. I. t.* 1.

Alpine Himalayas, from Kashmir to Kumaon, rare ; Sikkim, 9,000–15,000 feet elevation, Lachen Valley.

(Also in South Europe and the Mediterranean region, from Macaronesia to the Caucasus and Abyssinia.)

2. *NOTHOLÆNA VELLEA*. (*R. Br.*) Stipes densely tufted, woolly, wiry, short ; fronds 8–9 inches long, 1–1½ inch broad, oblong-lanceolate, bipinnate, pinnæ close lanceolate, the central ones the largest, with close roundish or oblong entire or 3-lobed pinnules, texture herbaceous but thick, both sides, especially the lower, coated with whitish or subferruginous tomentum, rachis bright chesnut-brown, more or less woolly. *R. Br. Prod. p.* 146. *N. lanuginosa*, *Desv. Hook. Syn. Fil. p.* 370.

Pangi and Lahul in Chumba, Cashmire.

(Also in Afghanistan, South Europe, Madeira, Cape Verd Isles, Algiers.)

GENUS LXVII.—MONOGRAMME. (*Schk.*)


Mono, one ; *gramme*, a line—sori in a single line on each frond.)

Sori subimmersed, linear elongated close to the midrib on one or both sides, the receptacles formed of a portion of the costa ; veins consisting only of a costa ; fronds small, grass or rush-like, simple or forked, rhizome creeping.

1. *MONOGRAMME PARADOXA*. (*Fée.*) Rhizome creeping, hairy ; fronds linear filiform, grass-like, 2–12 inches long, ¼–½ line broad ; sori within a vaginiform expansion of the costa, one side of which is larger than the other. *Fée. Vitt. p.* 38. *M. Junghuhnii*, *Hook. Sp. Fil.* v. 123. *Bedd. F. S. I. t.* 210.

Ceylon, 3,000–5,000 feet elevation, not very common.

(Also in Java, Philippines, Queensland and Polynesian Islands.)



A. & C. G. TEL. & C. P. G. RAY

MONOGRAMME PARADOXA. (Fie.).

GENUS LXVIII.—LEPTOGRAMME. (*J. Sm.*)

(*Leptos*, slender ; *gramme*, a line.)

Sori arising from the veins over the under surface of the frond, linear or linear oblong, simple ; veins free ; fronds bi-tripinnatifid, adherent to the caudex, habit and mode of growth of *Phegopteris* and *Lastrea*, and only differing from the former in its elongated sori.

1. LEPTOGRAMME TOTTA. (*Schl.*) Rhizome scarcely creeping, stipes subtufted, 6–12 inches long, pilose, below scaly ; fronds 12–18 inches long, 6–8 inches broad, pinnate, both sides pilose, pinnæ lanceolate, $\frac{1}{4}$ –1 inch broad, pinnatifid half-way to the midrib, lobes entire, entire or subentire, $1\frac{1}{2}$ –2 lines broad, herbaceous in texture ; veinlets simple 5–7 on each side with the linear oblong sori medial. *Schlecht, Adumbr.* 15, t. 6. *Hook. Syn. Fil.* 376. *Grammitis Totta*, *Bedd. F. S. I. t.* 49.

South India, very common on the mountains on the west side to the highest elevations ; Ceylon at the highest elevations ; North India, North-west Himalayas, Kashmir to Bhotan, 6,000 feet elevation, rare, Khasya, 3,000–5,000 feet elevation, common.

(Also in Java, China, Japan, and Africa and its islands.)

2. LEPTOGRAMME AURITA. (*Hook.*) Rhizome extensively creeping ; stipe 1 foot long, naked, glossy, the base curved and furnished with a few deflexed ovate acute scales ; fronds up to 2 feet long, and 1 foot broad, pinnate, pinnæ in distant pairs pinnatifid rarely or quite to the rachis into lanceolate entire or crenate lobes, lowest on one or both sides longer than the others and pinnatifid, others sometimes unequal, texture subcoriaceous, rachis glossy, both surfaces generally glabrous ; veinlets mostly forked ; sori oblong. *Hook. Syn. Fil.* 377. *Gymnogramme*, *Hook. Sp. Fil.* v. 141. *Grammitis aurita*, *Bedd. F. B. I. t.* 152.

Sikkim and Bhotan, 3,000–6,000 feet elevation, Assam, Khasya, 2,000–5,000 feet elevation.

VAR. LEVINGII. (*Clarke.*) Fronds weak and flaccid, pinnæ $1\frac{1}{2}$ inch



H. B. K. & F. H. C. H. C.

LEPTOGRAMME TOTTA. (Schl.)

y, and $\frac{1}{2}$ inch broad, not auriculate, lower ones very distant, rugly covered on both sides with long needle-like weak hairs. *F. N. I. p.* 568.


hmir, Jhelum and Chittapani valleys, 4,000-7,000 feet (Levinge), Sutlej, 9,000 feet elevation (Stewart).

LEPTOGRAMME OPACA. (*Spr.*) Stipes $1\frac{1}{2}$ and more feet long, hairy near the base ; fronds 2 feet and more long 12-15 inches



Nº 216

LEPTOGRAMME AURITA. (*Hook.*)



Nº 217.

LEPTOGRAMME OPACA. (*Spr.*)

base, subdeltoid-ovate acute membranaceous, pubescent on sides and costæ, bipinnate, lower primary pinnæ petiolate as long, 3-4 inches wide, oblong-ovate, their pinnules an inch long sessile and adnate, and slightly decurrent at the base, or acute, pinnatifid, the lobes retuse or emarginate and veins pinnate in each lobe of the pinnules or large, veinlets forked or simple, soriferous towards the base of the branches ; sori oblong. *Hook. Syn. Fil.* 378. Gymno-

gramma, *Hook. Sp. Fil.* v. p. 143. *Spr. Syst. Veg.* iv. 39. *Bedd. F. B. I. t.* 238. *Gynnogramma obtusata*, *Bl. Fil. Jar.* p. 97, t. 43. and *Hook. Sp. Fil.* v. 143.

Nepal to Bhotan, 4,000–7000 feet elevation; Khasya, 3,000–5,000 feet elevation.

(Also in Java.)

GENUS LXIX.—STEGNOGRAMME. (*Bl.*)

(*Stegnos*, a cover; *gramme*, a line.)

As in *Leptogramme*, only the veinlets of contiguous groups uniting as in *Nephrodium*, fronds pinnate; habit and mode of growth of *Nephrodium* and *Goniopteris*, only differing from the latter in its linear sori.

I. STEGNOGRAMME ASPIDIODES. (*Hook.*) Fronds hairy on both sides, especially the veins, ovate-lanceolate acuminate, pinnate, firm-membranaceous, pinnæ opposite or alternate, sessile subtruncate or adnate at the base, 3–4 inches long, rarely an inch wide, oblong-lanceolate, pinnatifid $\frac{1}{4}$ – $\frac{1}{3}$ the way to the midrib; veins 5–6 pair, 2–5 lower united, with an excurrent spurious vein reaching to the sinus, those in the lobes free, all soriferous; sori linear-oblong, nearly the length of the veins. *Hook. Sp. Fil.* v. 150. *Bl. Fil. Jar.* p. 172. *Bedd. F. B. I. t.* 149.

Khasya, 4,000–6,000 feet elevation; Ceylon (?)


(Also in Java.)

GENUS LXX.—GYMNOGRAMME. (*Desv.*)

(*Gymnos*, naked; *gramme*, a line.)

Sori arising from the veins over the under surface of the frond; linear or linear-oblong, forked or sometimes simple; veins forked, veinlets free; fronds various, adherent to the caudex, habit and mode of growth of *Cheilanthes*.

19215.



STEGOGRAMME ASPIDIOIDES. (*Hook.*)

1. *GYMNORHAMME ANDERSONI*. (*Bedd.*) Caudex furnished with numerous black wiry roots; stipes tufted, 1-2 inches long, shaggy, with long soft golden hairs; fronds (including stipes) 1-4 inches long, $\frac{1}{2}$ inch broad, membranaceous pinnated, copiously hairy, especially beneath, pinnae sessile, cordate-ovate or oblong-ovate, pinnatifid or crenated; costa not very prominent, veins pinnate



N°218.


GYMNORHAMME ANDERSONI. (*Bedd.*)

in the lower segments of the pinnae, forked or simple in the others; all free and not reaching the margin; sori forked copious. *Bedd. F. B. I. t.* 190. *Hook. Syn. Fil.* 380.

Kumaon, Soondadunga Valley, on dry rocks, 13,000 feet elevation.

2. *GYMNORHAMME LEPTOPHYLLA*. (*Desv.*) Root a small annual

mass of fibres; stipes tufted, glossy brown, slender, filiform, 1-2 inches long; fronds small 2-4 inches long, 1-1½ inch broad, membranaceous very transparent, ovate or deltoid, all varying from reniform crenated to variously pinnate or pinnate, those on the shorter stipes being less compound, pinnate late to obovate, decurrent, deeply crenated, lobes entire or crenate; veins simple or forked, not reaching the margin; sori



GYMNOGRAMME LEPTOPHYLLA. (Desv.)

; simple or forked. *Desv. Journ. Bot. i. p. 26. Hook. Sp. Fil. . Bedd. F. S. I. t. 270. Anogramma, Link.*
 Western ghats of the Peninsula of India, Ootacamund, Marwar, Suttara Fort walls.
 Also in Europe, the Azores, Madeira, Canaries, Africa, Persia, India, New Zealand, South America.)

3. *GYMNOGRAMME MICROPHYLLA*. (*Hook.*) Stipes densely tufted, 2-4 inches long, slender, fragile, glossy brown; fronds $1\frac{1}{4}$ -3 inches long, deltoid quadripinnatifid, pinnæ close, the lower ones deltoid, pinnules deltoid, ultimate segments $\frac{1}{8}$ - $\frac{1}{6}$ of an inch, obovate-oblong, texture pellucid-herbaceous; veins and sori one to each ultimate segment. *Hook. Syn. Fil.* 383. *Bedd. F. B. I.* t. 148, (habit of *leptophylla*, but not annual.)




GYMNOGRAMME MICROPHYLLA. (*Hook.*)

Khasya, Surareen, 5,000 feet elevation, Sikkin, Tonglo, 7, 10,000 feet elevation, Sinchal, 8,000 feet elevation, Dikee 7,500 feet elevation.

GENUS LXXI.—SYNGRAMME. (*J. Sm.*)

(*Syn* to unite; *gramme*, a line—sori often in united lines.)

Sori long linear, arising from the veins and veinlets on the under surface, veins forked close to the midrib or higher up, veinlets parallel



F. & C. 1863.

SYNGAMME PRAXINEA. (Don.)

creeping; sapes naked, glabrous,
simply pinnate with the pinnæ at
the base, stalked, and 2 inches
pinnæ 3 inches long by $\frac{5}{8}$ inch
broad rounded base, in both fo
herbaceous, glabrous, or a little
or finely toothed; veins very clo
midrib, or much higher up, one
forked, rarely joining with the
along all the branches, but stoppi
zium fraxineum, *Don. Fl. Nep.*
(*Bl.*), *Bedd. F. S. I. t. 232, the*
bipinnate form. *G. serrulata, Bl. Pl*

Himalayas, from Chumba to
very common; Khasya, 1,000-5,0
6,000 feet elevation; Malay Penin
(Also in the Malay Islands;
Japan.)

This Fern should certainly be
"alismaefolia," with which it quite
of the veins is rare, but I detect it

2. SYNGRAMME VESTITA. (H

rked or pinnate, where pinnate the lower branches generally
astomose; sori universal
the veins. *Bedd. F. B.*
t. 154. *Grammitis vestita*,
Vall. Cat. 12. *Gymno-*
ramme vestita, *Hook. Syn.*
Fil. 379

Himalayas from Chum-
a to Nepal, 6,000-9,000
et elevation.

(Also in China.)


3. *SYNGRAMME WAL-*
CHII. (*Hook.*) Stipes
fted up to 20 inches long,
n, glossy brown, fibrillose
ar the base; fronds up to
e foot long by 2½ inches
ad, elliptic-lanceolate,
ple, acuminate, entire at
margin, the base narrow-
very gradually, texture
coriaceous, both sides
brous; veins fine parallel,
ple, or forked from the
e or much higher up,
ely joining with the con-
uous group about the
tre, but all anastomosing
ar the margin and forming
3 small oblong hexagonal
oles; sori on all the
traight veins, but scarcely
extending to anastomosing
nes near the margin. *Bedd.*

F. B. I. t. 153, *Gymnogramme*, *Hook. Sp. Fil.* v. p. 155, t. 302



№ 229.

SYNGRAMME VESTITA. (*Wall.*)



Singapore.

Also in Borneo.)

1. SYNGRAMME ALISMÆFOLIA. (*Hook.*) Caudex creeping ; a span to a foot and more long, blackish-purple ; fronds subcoriaceous, simple 5–8 inches long, $2\frac{1}{2}$ inches wide near the base, from rounded base, ovate or ovate-lanceolate, finely acuminate ; veins pinnate, approximate, simple or forked, anastomosing near the apex into 1–2 series of oblong hexagonal areoles ; sori narrow-linear on all the straight veins, but scarcely extending to the anastomosing ones. *Bedd. F. B. I. t.* 240. *Gymnogramme*, *Hook. Sp.* 155.

Singapore.

GENUS LXXII.—SELLIGUEA. (*Bory.*)

(After Selligie, a Frenchman.)

Veins prominent, compound, anastomosing, with free included veins in the areoles ; sori long linear ; fronds simple, pinnatifid or bipinnate, articulate with the caudex ; habit and mode of growth of *Pleopeltis*, from which genus it only differs in the sori long linear.


* *Fronds simple.*

1. SELLIGUEA FEEI. (*Hook.*) Rhizome firm, wide-creeping, scales bright brown, fibrillose ; fronds dimorphous, the sterile ones 1–2 inches long, $1\frac{1}{2}$ – $2\frac{1}{4}$ inches broad, on stems $3\frac{1}{2}$ –6 inches long, the fertile ones narrower and more contracted at the base on stems 6–10 inches long, both acute, entire, very coriaceous and glabrous ; main veins very prominent to the margin, $\frac{1}{4}$ inch apart, veinlets obscure, forming copious areoles, with free included clavate venules ; sori linear, immersed in single rows, rarely interrupted, between the main veins, not quite reaching the margin. *Hook. Syn. Fil.* 389. *Sp.* 158. *Bedd. F. B. I. t.* 151.

Malay Peninsula, Penang, Singapore.

(Also in the Malay Islands.)

the broad, fertile ones 4-8 inches
more than 1 foot long, texture ra-
spicuous; main veins very distinct.



Nº 226.

SELLIGUEA HAMILTONIANA.

(Wall.)


both sides not...

It
vi
re
at
H
no
Be

Ne
the

(H
the
late
erec
inch
the
1-2 i

Nº 227.



SCIENCE

SELLIQUEA ELLIPTICA. (Thunb.)

subpalmately divided into 3-5 l-
quite pinnate, lobes or pinnæ l-
broad, texture subcoriaceous, quite
distinct to the edge, areoles un-
clavate veinlets; sori linear obliqu-
margin, sometimes interrupted
ellipticum, *Thunb. Fl. Jap.* 335.
Fil. 389. *Bedd. F. B. I. t.* 150.

Himalayas, Nepal to Bhotan
feet; Malay Peninsula, Tenasserim
(Also in the Philippines, Quee-

5. *SELLIGUEA MAINGAYI*. (*B.*
naked; fronds deltoid, under 1 foot
lanceolate pinnæ, the end one largest
broad, slightly repand, narrowed to
pair distant, texture membranaceous
to the edge, $\frac{1}{4}$ inch apart, areoles cor-
veinlets; sori very copious, minute in
lines or curves. *Baker in Hook.*
Malacca.

GENUS LXXIII.—LOXI

Fronds slender, wide-creeping; scales small, linear, brown; fronds 6–10 inches long by $\frac{1}{8}$ – $\frac{3}{4}$ inch broad, linear lanceolate, the apex acute, the margin entire, the lower third narrowed gradually into a very short stipe, texture thick, coriaceous but flaccid, both sides glabrous; venation obscure, costal areoles numerous, small, without, or more rarely with, free included veinlets, the midrib much more slender than in *involuta*; the sori shorter and often nearly parallel, or quite parallel, to the midrib. *Grammitis lanceolata*, Sw. Syn. Fil. 22, 212, t. fig. 4. *Selliguea*, Hook. Syn. Fil. 387. Sp. Fl. v. 156. *Loxogramma lanceolata*, Bedd. F. S. I. t. 51.

Southern India, Nilgiris and the higher Western mountains, 6000–8,000 feet elevation, (rare compared with *involuta*). Ceylon, above Newera Elya, rare; Khasya, 4,000–5,000 feet elevation.

(Also in China, Japan, Fiji, Samoa, and Africa, with its eastern islands.)



2. *LOXOGRAMME INVOLUTA*. (*Don. under Grammitis*.) Rhizome short, creeping, the scales lanceolate, brown; fronds 8–18 inches long, 1–2½ inches broad, lanceolate, the apex acuminate, the margin entire, the lower part narrowed very gradually into a short broad compressed stipe; texture thick coriaceous, but flaccid, both sides glabrous; venation obscure, areoles copious with free included veinlets; sori in long parallel very oblique lines reaching from the midrib nearly to the margin. *Don. Fl. Nep.* 14. *Selliguea*, Hook. Sp. Fil. v. 155. Syn. Fil. p. 387. *Loxogramma involuta*, Bedd. F. S. I. t. 50. Wall. Cat. 6, 7 and 10.

Very common on the mountains throughout the Indian region; Himalayas, 2,000–7,000 feet elevation; Southern India and Ceylon, 5,000–8,000 feet.

Perhaps only a large form of *lanceolata*. Baker says there are no free included veinlets, but I find them always present in this plant, (Sir J. Hooker says sometimes), generally absent though sometimes present in *lanceolata*.

(Also in the Malay Islands and Polynesia.)


3. *LOXOGRAMME AVENIA*. (*Baker*.) Rhizome creeping; stipe



1 inch long, fronds glabrous, coriaceous, 18-20 inches long, linear-g, spatulate, broader up-
 , but narrowed and acute
 : apex, about 3 inches
 n the widest part, glabrous
 oth sides, texture less
 :eous than in the last
 venation more evident,
 lated with free veinlets
 me of the areoles, no
 nent costules though
 imary veins are slightly
 r than the others; sori
 row linear more or less
 se, parallel lines in the
 portion of the frond
 ouching the costa nor
 ding to the margin.
 ; *Syn. Fil. p.* 388. *Bedd.*
I. t. 266.

The Malay Peninsula,
 ng.

Also in the Malay Islands.)





LOXOGRAMME AVENIA. (*Baker.*)

GENUS LXXIV.—BRAINEA. (*J. Sm.*)

(After C. Braine who first introduced the fern.)

Sori linear, simple or branched, usually on the veins which form
 costal areoles and on the bases of the free transverse ones, often
 extended and confluent; veins united so as to form one series
 long or triangular costal areoles, the rest free, simple or forked,
 enerally all free towards the apex of the pinnæ; caudex erect,
 ascent; fronds pinnated continuous with the caudex.

. BRAINEA INSIGNIS. (*Hook.*) Caudex as thick as a man's



m, clothed with shaggy dark-brown ferruginous subulate-lanceolate scales, $\frac{3}{4}$ –1 inch long; stipes stout, firm, 3–4 inches long, scaly only the base; fronds coriaceous bright-green (Lomaria-like), pinnate, or occasionally below partially bipinnate; pinnæ numerous, close, horizontal, 5–6 inches long, 4 lines broad, base cordate, linear-oblong, acuminate, finely serrate. *Hook. Syn. Fil.* 390. *Bedd. F. B. I. t.* 139.

Khasya Hills, 3,000–4,000 feet elevation, Pomrang and Jainka; Malay Peninsula, in Tenasserim, the fir forests (Parish), on Mooleyit, 6,000 feet (Beddome.)

(Also in Hong Kong.)

GENUS LXXV.—MENISCIUM. (*Schreb.*)




(*Meniskos*, crescent; shape of fructification,)

Sori naked, oblong or linear, occupying the transverse connivent veinlets; veins pinnate, veinlets numerous, the opposite ones uniting in an arc or angle, and sending out from the angle a free or continuous venule; fronds simple or pinnate, (only differs from *Goniophlebium* in the shape of the sori.)

1. *MENISCIUM TRIPHYLLUM*. (*Sw.*) Rhizome firm, wide-creeping, clothed with chesnut lanceolate-linear scales at the extremities; stipes slender, slightly pubescent, of the fertile frond 1 foot long or more, of the sterile often shorter; fronds 3-foliate or with 5 or more pinnæ, the terminal one much the largest, all oblong-lanceolate with a broad base and acute apex, about 4–6 inches long by $\frac{3}{4}$ –1 $\frac{1}{2}$ inch broad, sessile or subsessile, margin entire or subrepand, the fertile ones often lower, texture herbaceous, slightly pubescent beneath; areoles between midrib and margin. *Sw. Syn. Fil.* 19, 206. *Hook. Syn.* 391. *Bedd. F. S. I. t.* 56. *Wall. Cat.* 61.

South India, Bolampatty Valley, Anamallays, Travancore, 2,000–3,000 feet elevation, gregarious and forming very large beds, nearly always trifoliate, rarely 5-foliate; Ceylon, Matale and Saffragam, 2,000–3,000 feet; North India, Eastern Bengal, Cachar, Chittagong, Sikkim, at no great elevation; Malay Peninsula.

(Also in China and Philippines.)



. β PARISHII, pinnæ 9-13, Tenasserim and Mishmee. *B. I. t.* 184; this probably runs into the type, but there is ke it in South India.

MENISCIUM THWAITESII. (*Hook.*) Rhizome wide-creeping, a crowquill, stipes 9-12 inches long, angled, naked except base, fronds 8-10 inches long, 4-6 inches broad at base, subdeltoid, the apex acuminate and more or less pinnatifid, which are several linear oblong rather deeply crenated




MENISCIUM THWAITESII. (*Hook.*) MENISCIUM SALICIFOLIUM. (*Wall.*)

the lowest stalked, the others sessile or subsessile, $\frac{1}{2}$ -1 inch long, suddenly narrowed at the base, texture herbaceous, little pubescent, above glabrous or nearly so, except the veins generally 5 series between the costa and margin. *t. Fil. p.* 391. *Bedd. F. S. I. t.* 223.

India, Nilgiris, west slopes below Sispara, 3,000 feet; Ceylon, Matale, 3,000 feet elevation.

MENISCIUM SALICIFOLIUM. (*Wall.*) Stipe 1 foot or more brownish or pale-brown, glossy; fronds 12-18 inches long,

longitudinal ven
 formed of two confluent rounded o
Syn. Fil. 391. *Bedd. F. B. I. t. 2*
Malay Peninsula, Penang and :



MENISCIMUM CUSPIDATUM. (Bl.)

Bl. En. Pl. Jav. Fil. 114. *Hooker, S.*
 309. *M. longifrons* " " -

Himalayas and Khasya, from 500-4,000 feet, and Mr. Clarke makes it a separate variety, under the name of *longifrons*; it, however, graduates into the type, I believe it is not separated at Kew, nor are there any characters to distinguish it; they both have the costa very red sometimes, Mr. Clarke also gathered small specimens with simple fronds (not in fruit, however).

(Mr. Clarke is quite wrong in referring *Gymnopteris costata*, var. *deltigera*, to this genus, it has quite different venation.)

GENUS LXXVI.—ANTROPHYUM. (*Kaulf.*)


(*Antron*, a cave, hollow; *phyo*, I grow.)

Sori reticulated or interrupted, carried along the veins in line, the receptacles immersed and forming grooves, or superficial; veins uniform reticulated; fronds adherent to the caudex, simple, with or without a defined midrib, fleshy-coriaceous in texture, all the species closely allied.

I. ANTROPHYUM RETICULATUM.

(*Kaulf.*) Stipe none or very short, the frond being decurrent down to the base; fronds 6-15 inches long, by 1-1½ inch broad, linear-lanceolate or acuminate, very gradually narrowed downwards; midrib none or sometimes present towards the base of the frond, the areoles very long and narrow and distinctly raised on the upper surface; sori immersed, sometimes confluent. *Kaulf, Féc, 3rd Mém. Foug. 14.* *Bedd. F. S. I. t. 231.* *Ant. coriaceum, Wall. Cat. 43.* *Ant. semicostatum (Bl.), Hook. Syn. Fil. 393.*

All the South Indian, Ceylon, and North Indian specimens seem to me to belong to one species; "coriaceum" is said to differ



N°235
ANTROPHYUM RETICULATUM. (*Kaulf.*)



by being plicate on the upper surface, but this peculiarity occurs also in the South Indian plant and is not a specific character; reticulatum can always be distinguished from "plantagineum" by its longer and narrower fronds, and generally also by the presence of numerous barren sporangiastra.

South India, rare, on the Tinnevely and Travancore Mountains, 3,000 feet elevation; Ceylon, southern and central provinces, up to 4,000 feet; Himalayas and Khasya, up to 5,000 feet; Malay Peninsula.

(Also in Polynesia and Queensland.)

VAR. β PARVULUM.

(Bl.) Fronds very small, 1-4 inches long, by $\frac{1}{4}$ - $\frac{1}{2}$ inch broad. Antr. parvulum (Bl.), Fil. Jav. 78, t. 34. Hook. Sp. Fil. v. 170. Bedd. F. B. I. t. 267.


Sikkim, Yoksum, 4,500 feet, Khasya; Penang.

(Also in Java.)

2. ANTROPHYUM PLAN-
TAGINEUM. (Kaulf.) Stipe distinct, 1-4 inches long; fronds oblong, broadest towards the apex, then suddenly narrowed into an acute point, 4-10 inches long, and up to 2 inches broad, no midrib or

an inconspicuous one towards the base; sori deeply immersed, sometimes distinctly raised on the upper surface making the frond plicate above. Kaulf, Bory. in Voy. de la Coq. Bot. Cryp. t. 28. Bedd. F. S. I. t. 52 (reticulatum).

South India, on the Western mountains, 2,000-5,500 feet



ANTROPHYUM LATIFOLIUM. (Bl.).

elevation, not very common ; Ceylon, central provinces ; Himalayas and Khasya up to 5,000 feet ; Malay Peninsula.

(Also in the Malay Islands, Philippines and Polynesian Islands.)

3. *ANTROPHYUM LATIFOLIUM*. (*Bl.*) Stipe long up to 7 inches ; fronds large, broadly obovate or round, up to $4\frac{3}{4}$ inches wide, acuminate or acutely lobed at the apex, no midrib ; sori superficial or immersed, with the frond plicate above. *Bl. Fl. Jav. Fil.* 75. *Bedd. F. B. I. t.* 176.

Sikkim and Bhotan, 2,000–6,000 feet ; Assam and Khasya 1,000–4,000 feet elevation.

(Also in Java.)


Supposed to differ from the last by the sori being superficial ; they however are deeply immersed in some of my specimens, so it only differs in its much broader frond and longer stipes ; all the three so-called species are probably only varieties of one plant.

GENUS LXXVII.—VITTARIA. (*Sm.*)

(*Vitta*, a riband, the riband-like frond.)

Veins simple, forming an acute angle with the midrib, their apices prolonged into a transverse marginal vein, which becomes the receptacle, or veins forked without the transverse marginal vein ; sori seated in an extrorse groove of the margin, or in a slightly intramarginal line with the unaltered edge of the frond produced beyond and often rolled over it ; fronds adherent to the caudex, linear, grass- or tape-like.

1. *VITTARIA ELONGATA*. (*Sw.*) Rhizome creeping, scales many, with black hair-like points ; fronds up to $2\frac{1}{2}$ feet long, generally only $\frac{1}{4}$ inch broad, acuminate, grass-like, but rather firm in texture, midrib generally more or less distinct beneath ; veins simple, oblique, immersed, parallel, connected by an intramarginal veinlet ; sori quite sunk in an extrorse marginal groove. *Sw. Syn. Fil.* 109, 302. *Hall. Cat.* 144. *Bedd. F. S. I. t.* 21.



A. D. S. 1904

VICTARIA ELONGATA. (Sw.)

South India, on the Western mountains, 2,000–5,000 feet elevation; Ceylon, central provinces; North India, from the plains up to about 4,000 feet elevation; Malay Peninsula, Birma, &c.

(Also in the Malay Islands; Queensland; Polynesia; Tropical Africa, and Mauritius.)


2. *VITTARIA SIKKIMENSIS*. (*Kuhn.*) Rhizome very shortly creeping, with slaty-rufous hair-pointed scales; stipes densely tufted; fronds up to 4 inches long, but often very much shorter, $\frac{1}{2}$ inch broad, subobtuse; midrib beneath obscure or slightly depressed; sori sunk

in a large extrorse marginal furrow. *Kuhn in Linnæa* xxxvi. 66. *Clarke, F. N. I.* p. 574. *V. minor* var. *minima*, *Hook. Sp. Fil.* v. 183. *Bedd. F. B. I.* t. 56 (not minor of *Fée*).

Sikkim, 2,000–6,000 feet elevation, common; Khasya, Mowlong, 2,500 feet elevation; Tenasserim.

The Tenasserim specimens are certainly the same as the Sikkim, and when Mr. Clarke stated that the Moulmein and Malay fern was distinct, he had in his eye only the Malacca plant (*i.e. falcata* or the next species). I have never seen the Tenasserim plant more than 2 inches long, and the Sikkim plant

also common in this small state, though other specimens are 4 inches long, the soral groove is extrorse as in *Vittaria elongata* (not intramarginal as in the section *Tæniopsis*) and this plant can hardly be said to differ from *elongata* except in its very small size, and probably only a variety of it.



Nº 239.

VITTARIA SIKKIMENSIS.
(*Kuhn.*)

3. *VITTARIA FALCATA*. (*Kunze.*) Fronds 4–5 inches long, 1 inch broad, the apex blunt, the lower part narrowed gradually to the base, texture leathery and very thick; a distinct raised midrib

attaining the apex in the barren fronds, but lost in the fertile ones ; veins short, oblique, parallel, immersed ; sori quite sunk, in deep intramarginal grooves. *Kunze. Hook. Syn. Fil.* 395. *V. falcata and minor Fée 3rd. Mém. Foug.*

Malacca.

(Also in the Philippines.)

4. *VITTARIA AMBOINENSIS.* (*Fée.*) Fronds 4–5 inches long, by 3–4 lines broad, smooth, submembranaceous, falcate acuminate, tapering below into a petiole ; costa slender, disappearing below the apex ; veins curved equal approximate ; sori closely marginal, cuticle of the margin resembling a false involucre ; caudex flexuose, contorted scaly, scales cancellate, rigidly toothed at the margin. *Fée, Vittar. p.* 44, *t.* 1, *f.* 1. (*not Mett.*) *Hook. Sp. Fil.* v. 177. *Bedd. F. B. I. t.* 117.

Martaban. This seems as distinct a species as most of this group, though it is possible it may be a small form of scolopendrina, the veins are combined where fertile by an intramarginal vein which runs along the centre of the receptacle.

(Also in Amboyna).

5. *VITTARIA LINEATA.* (*Sw.*) Rhizome very short creeping ; **stipes** tufted, scales acuminate with caudate points ; fronds up to 8 inches long, $\frac{1}{8}$ – $\frac{1}{3}$ inch broad, narrowed gradually downwards to the **stout** compressed stipe, the margin often reflexed, texture thick, a **distinct** raised midrib from the base to the apex ; veins simple, **immersed**, parallel, very oblique ; sori in a broad intramarginal shallow **furrow**, the edge of the frond distinctly beyond the furrow and at first **wrapped** over it. *Sw. Syn. Fil.* p. 109. *Hook. Syn. Fil.* 396. *Tæniopsis lineata*, *Bedd. F. S. I. t.* 54. *Fée* separated the Indian plant **from** the American under the name of *flexuosa*, and Mr. Clarke has **adopted** that name, but the two plants seem identical.

South India, on the Western mountains, 2,000–6,000 feet elevation ; Ceylon, ~~central~~ provinces, common ; Himalayas, 2,000–12,000 feet elevation, from Gurwhal to Bhotan, Khasya, 1,000–6,000 feet elevation ; Malay Peninsula.

(Also in Tropical America, and the West Indies, Africa and its Eastern Islands.)



N° 240.

VITTARIA SULCATA. (Kuhn.)


scolopendrina, *Pr. Tent. Pter. p.* 141. *Hook. Syn. Fil. p.* 396.

Tæniopsis, Bedd. F. S. I. t. 212.

Vitt. Zeylanica, Fée.

Ceylon, southern and central provinces, on rocks up to 3,000 feet elevation; Himalayas, Sikkim, Assam, Bhotan.

(Also in New Guinea, Philippines, Malay Islands, Seychelles, and Mozambique.)



N° 241.


VITTARIA SCOLOPENDRINA. (*Presl.*)

deep furrow confined to the very much thickened upper half of the frond, where the midrib and veins

7. VITTARIA SULCATA. (Kuhn.)

Rhizome short creeping, scales dense lanceolate, pale-brown; fronds crowded, ligulate, obtuse, sessile, narrowed to the base, 2-4 inches long, $\frac{1}{8}$ inch broad; sori in an intramarginal

NP 242



UTER & CO. STERIL

TÆNITIS BLECHNOIDES. (Sw.)

are quite lost ; midrib visible in the barren lower half, veins forked or rarely with two branches, veinlets slightly clavate at the apex not reaching the margin, margin beyond the soral groove very thick. *Kuhn, Linnæa*, 36, p. 68. *Tæniopsis falcata*, *Bedd. F. B. I. t.* 175.

Ceylon, 4,000–5,000 feet elevation, the forked venation is abnormal in the genus.

GENUS LXXVIII.—TÆNITIS. (*Willd.*)

(From *tainia*, a fillet or ribbon.)

Veins reticulated, forming oblong hexagonal oblique areoles; sori linear, but the line sometimes interrupted, forming a transverse band between the midrib and margin ; fronds adherent to the caudex.

1. TÆNITIS BLECHNOIDES. (*Sw.*) Rhizome creeping, setose; stipes 8–12 inches long, firm, naked, glossy; fronds 1–2 feet long, 8–12 inches broad, pinnate; pinnæ of barren frond 2–3 on each side, 1–2 inches broad, oblong-lanceolate, the point acuminate, the edge thickened and wavy, the base cuneate, the lower ones stalked, fertile pinnæ more numerous and narrower, texture coriaceous, areoles copious, oblique, without free veinlets; sori in a continuous (rarely interrupted) line, about midway between the edge and midrib. *Swartz, Syn. Fil.* 24 and 220. *Hook. Syn. Fil.* 397. *Bedd. F. B. I. t.* 54.

Ceylon, in the forests about Galle; Malay Peninsula, in Tenasserim and further south; Sylhet (?).

(Also in the Philippines.)

GENUS LXXIX.—DRYMOGLOSSUM. (*Presl.*)

(*Drymos*, wood; *glossa*, tongue).


Veins obscure, compoundly anastomosing in the sterile fronds, forming 3–4 series of areoles between the midrib and the margin, each including simple or forked free veinlets, with clavate apices; fronds articulate with the caudex, dimorphous, the sterile broad and

ort, the fertile long and narrow; sori linear, scarcely immersed, ntral or submarginal often at length confluent and covering the hole underneath surface of the frond; caudex wide-creeping on es.

1. *DRYMOGLOSSUM CARNOSUM*. (*Hook.*) Rhizome long, filiform, ry, clothed with peltate lanceolate-linear toothed scales, which are en hair-pointed; stipes 2 lines to 1 inch long, distant slender; nds simple, of two kinds, sterile one from $\frac{1}{2}$ an inch, (and en generally orbicular or sub- rdate) to 2 and even 3 inches ng, and then elliptical or obovate spathulate, or even lanceolate d acuminate, thick and fleshy, riaceous when dry, faintly cos- e on the under side; veins an- tomosing, the areoles including e veinlets, fertile fronds 1-2 $\frac{1}{2}$ ches long, linear-spathulate ob- se; sori linear continuous form- g a line intermediate between e costa and the margin, soon nfluent, and representing one ad band, nearly as broad as e frond, when young covered numerous peltate pedicellated ales.


Hook. Sp. Fil. v. 189. *Nothochlæna?* (*Tænitis?*) *carnosa*, *Wall. Cat. n.* 138. *Bedd. F. B. I. t.* 55.

Nepal, Sikkim, Bhotan, 2,000-5,000 feet elevation, common. (Also in China and Japan.)



DRYMOGLOSSUM CARNOSUM. (*Hook.*)

2. *DRYMOGLOSSUM PILOSELLOIDES*. (*Presl.*) Rhizome long iform, wiry, clothed with adpressed, diamond shaped peltate laci- ated scales, which are sometimes hair-pointed; stipes about 2 lines ng in the sterile, often about 1 inch long in the fertile fronds, nds dimorphous, the barren ones roundish or obovate, $\frac{1}{2}$ -2 inches



A. P. CASTYLLA & C. V. SORIANO

DRYMOGLOSSUM VILOSELLOIDES. (Presl.)

long, $\frac{3}{4}$ inch broad, very thick and fleshy, and when young more or less covered with stellate hairs, the fertile ones, 2–4 inches long $\frac{1}{8}$ – $\frac{1}{4}$ inch broad; veins immersed, areoles with copious free veinlets; sori in broad continuous marginal lines often at length confluent and covering the whole under surface, capsules mixed with a few stellate paraphyses. *Presl. Tent. Pterid.* 227, t. 10. *Bedd. F. S. I. t.* 55 and *F. S. I. t.* 186 (*Niphobolus nummularifolius*).

Bengal Plains; Birma; Ceylon; South India, common in the Malabar plains, (Calicut, &c. on trees), also on the mountains, up to about 2,000 feet elevation, (Anamallays, Wynad, Coorg, &c.)

(Also in Java Philippines and Japan.)

VAR. β BEDDOMEI. Mr. Clarke proposes the name *Beddomei* for a specimen from the Anamallays figured by me (*tab.* 186, *F. S. I.*) because the fronds have stellate hairs and the scales of the rhizome are more pointed, but I find this stellate pubescence is always more or less present on the young fronds of Himalayan examples, and the rhizome scales are quite as hair-pointed in some specimens from Java, Philippines, and Japan, the fertile fronds are shorter and narrower in this South Indian form, and the broader usually sterile fronds sometimes fructify towards the apex, but I am not inclined to consider it even a permanent variety, as I believe it runs into the type with longer fertile fronds, which is also found in Southern India.



(*Niphobolus nummularifolius*, though I have included it in that genus, rather belongs here, it has exactly similar venation.)

GENUS LXXX.—HEMIONITIS. (L.)

(*Hemionos*, a mule—the mule fern.)

Sori continuous along the veins and copiously reticulated; veins copiously anastomosing, forming numerous areoles which have rarely a free veinlet in them, which when present is also soriferous; fronds adherent to the caudex, simple, pinnatifid, or pinnate.

1. HEMIONITIS ARIFOLIA. (*Burm. under Asplenium.*) Caudex




act, short; stipe of the barren frond, 2-4 inches long, of the fertile frond 1 foot long, dark chesnut-brown, glossy densely fibrillose towards the base; fronds 2-3 inches each way, cordate-hastate, the barren ones bluntish at the apex, with generally rounded (rarely indented) basal lobes and a deep sinus, the fertile ones with the lobes more produced and pointed, texture more or less coriaceous, glaucous above, somewhat villous beneath; areoles oblique, numerous, the veinlets none or very rare. *Burm. Fl. Ind.* 231. *H. cordifolia* (Lam.) *Bedd. F. S. I. t.* 53.

South India, common in the plains and on the mountains up to nearly 3,000 feet elevation; East Bengal plains; Ceylon; Birma.

(Also in the Philippines.)

2. *HEMIONITIS GRIFFITHII*. (*Hook. Fil. et Thoms.*) Rhizome short creeping; stipes coriaceous and coarsely hirsute, 8-12 inches long; fronds 1-6 inches long, 6-10 inches broad, subdeltoid, pinnatifid, bipinnate with 2-4 pinnae on each side, which are 1-1½ inches broad, entire or creased, acuminate, texture coriaceous, rachis and both faces hairy on the venation; main veins prominent, generally venose, the veinlets reticulated into copious areoles all soriferous, rarely there are free included veinlets, which are also soriferous. *Hook. Sp. Fil.* v. 192. *Dictyocline Griffithii* (*Moore*), *Bedd. F. B. I.* 155.

Khasya, Cherra, 4,000-5000 feet elevation.
(Also in Formosa.)



Nº 296

HEMIONITIS GRIFFITHII.
(*Hook. et Thoms.*)

TRIBE XII.—ACROSTICHEÆ.

Sori spread in a stratum over the under surface, or rarely over both surfaces of the frond, not confined to the veins only.

GENUS LXXXI.—ELAPHOGLOSSUM. (*Schott.*)

(*Elaphos*, a stag ; *glossa*, tongue.)

Veins free, simple or forked, their apices sometimes clavate, fronds simple, entire, sessile or stipitate, the fertile somewhat contracted and generally sporangiferous over the whole under surface ; stipes adherent to the rhizome, but generally pseudo-articulate a little above the base.

1. ELAPHOGLOSSUM CONFORME. (*Sw.*) Rhizome woody, wide creeping, scales blackish, ovate, jagged, not hair-pointed ; stipes firm erect, 2–3 inches long in the sterile, and often much more in the fertile, black at the base up to the pseudo-articulation, where it breaks off in age, generally clothed with sheathing scales ; sterile fronds 2–9 inches long, seldom more than 1 inch broad, narrow-lanceolate acuminate, furnished with deciduous scales on both sides, quite glabrous in age, margin slightly revolute in age ; veins hidden, generally once-forked, just reaching the margin ; fertile fronds somewhat contracted. *Sw. Syn. Fil.* 10, 192, *t.* 1. *Bedd. F. S. I. t.* 198. *Hook. Syn. Fil.* 401.

South India, Western mountains, at the higher elevations, very common ; Ceylon ; Malay Peninsula ; Sikkim and Nepal, 6,000–9,000 feet elevation, Khasya, 4,500–6,000 feet elevation.


(Also in Australia, Central and South Africa, Queensland, Polynesia, and the Malay Islands.)

2. ELAPHOGLOSSUM LATIFOLIUM. (*Sw.*) Rhizome woody, wide-creeping, scales bright chesnut or golden, lanceolate and more pointed than in *conforme* ; stipes longer than in *conforme* and the deciduous scales not sheathing ; fronds much larger than in *conforme*, and generally over 2 inches broad, margin diaphanous and much




J. C. Pringle

ELAPHOGLOSSUM CONFORME. (Sw.)



3. ELAPHOGLOSSUM NORRIS



ELAPHOGLOSSUM LATIFOLIUM. (Ste.)

sheathing, very dense towards the l




N° 249.

ELAPHOGLOSSUM STIGMATOLEPIS. (Fée.)



N° 250

ELAPHOGLOSSUM VISCOSUM. (Sw.)



N° 251

ELAPHOGLOSSUM SQUAMOSUM. (Sw.)



N° 252

ELAPHOGLOSSUM SPATHULATUM. (Sw.)

South India, Nilgiris, and Palghat Hills.

5. *ELAPHOGLOSSUM VISCOSUM*. (Sw.) Rhizome woody, creeping, the scales bright-brown, narrow, linear, hair-pointed; sterile fronds linear-lanceolate, 6-9 inches long, by $\frac{1}{2}$ - $\frac{3}{4}$ inch broad, narrowed gradually at both ends, densely covered on the under surface with stellate pubescence, glabrous above, at least in age, its stipe 2-3 inches, scaly and with stellate pubescence; veins apparent, their clavate apices well within the margin, generally once forked, rarely the forks again forked, fertile fronds contracted and on a longer stipe. Sw. *Syn. Fil.* 10, 193. *Bedd. F. S. I. t.* 196. *E. stelligerum*, Wall. *Cat.* 2167.

South India, Western mountains, Anamallays 4,000 feet, on rocks up the Toracadu River, Coorg, Travancore; North India, Sikkim and Nepal, 6,000-8,000 feet elevation, Khasya, 4,000-6,000 feet. (It has been proposed to separate the South Indian plant from the Himalayan under the name of *stelligerum*, but I cannot distinguish them.)

(Also in Tropical America; Tropical Africa, and its eastern islands; and the Malay Islands.)

6. *ELAPHOGLOSSUM SQUAMOSUM*. (Sw.) Rhizome short creeping, scaly; the scales black margined and ciliate; stipes 1-2 inches long, very scaly, fronds linear-lanceolate obtuse, gradually attenuated at the base, 4-12 inches long, $\frac{1}{2}$ - $\frac{3}{4}$ inch broad, densely covered on both sides with velvety ciliated scales, fertile fronds often not contracted, texture flaccid; veins hidden, simple or forked. Sw. in *Schrad. Journ.* 1800, ii. p. 11. *Bedd. F. S. I. t.* 197.

South India, Nilgiris, Anamallays, Travancore hills, 3,000-4,000 feet elevation, not common; Ceylon, central provinces.

(Also in Tropical America and West Indies; Sumatra; Sandwich Islands; Mascareen Islands; Guinea Coast; Madeira and Azores.)

7. *ELAPHOGLOSSUM SPATHULATUM*. (Sw.) Rhizome short, creeping, densely scaly, furnished with numerous wiry roots; stipes 2-4

ches long, densely covered with reddish setaceous scales, fronds near-lanceolate to rhomboid-lanceolate 1-2 inches long, covered on both sides with numerous hair-like scales, fertile fronds broad ovate, smaller than the sterile ones. *Sw. Syn. Fil. p. 10. Bedd. F. S. I. 213. Acrostichum piloseloides, var. ♂ spathulatum. Hook. Sp. Fil. 228.*

Ceylon, about Newera Elya, and in the southern provinces.


(Also in Tropical America and West Indies ; Natal and Mascaren Islands ; and Tristan d'Acunha.)

GENUS LXXXII.—STENOCHLÆNA. (*ℱ. Sm.*)

(*Stenos*, narrow ; *chlæna*, cloak ; the narrow involute margin.)

Fronds simply pinnate, the fertile contracted and very narrow, the sterile with the habit of *Lomaria* ; veins simple or forked, fine and close, generally quite free to the margin, or rarely the two forks or even two separate veins anastomose ; stipes adherent to the rhizome ; pinnæ pinnately pinnate with the rachis. (In *palustre*, the rachis or costa of the sterile pinnæ is winged, particularly towards the apex, which wing has been called an obscure transverse vein, anastomosing in loops ; the same occurs in *Pteris patens*, and some *Athyriums* (very apparent in *Athy. fimbriatum, var. sphæropteroides*) and it can scarcely be called a true vein.

1. *STENOCHLÆNA PALUSTRE. (Linn. under Polypodium.)* Rhizome scandent, (often reaching the tops of the highest trees), fronds pubescent, shining, of hard texture, pinnate, 1-4 feet long, pinnæ pinnately pinnate numerous, alternate or opposite, lanceolate acuminate, apically serrate towards the apex, oblique at the base, and furnished with a marginal gland on the upper edge, 5-10 inches long, 1-1½ inch broad, fertile fronds very much contracted ; veins simple or forked, generally free to the thickened margin, rarely the forks or two separate veins anastomose in the middle of, or towards the margin of the pinnæ ; rachis of sterile pinnæ winged, particularly towards the margin, and forming a pseudo vein parallel with it




STENOCHLOA PALUSTRE. (Linn.)

Burm. Fl. Zey. 234. *S. scandens*, *J. Smith*, in *Hook. Journ. of Bot.* iii. 401. *Bedd. F. S. I. t.* 201. *Lomaria scandens*, *Willd. Sp. Pl.* 293.

South India, in the plains on the West Coast and up the mountains to about 3,000 feet elevation; Ceylon. North India in the plains of Bengal and at low elevations on the hills; Malay Peninsula. (*Davallia achilleifolia*, *Wall.* *Teratophyllum aculeatum*, *Mett. Ann. Mus. Lug. Bat.* 4, 296. *Bedd. F. B. I. t.* 209, is an abnormal form of this plant, showing clearly a winged partial rachis.)

(Also in South China; Queensland; and Fiji.)

2. *STENOCHLÆNA SORBIFOLIA*. (*L.*) Rhizome thick, woody, often 40 feet long, clasping trees like a cable, sometimes prickly, scales lanceolate-subulate, large; fronds up to 18 inches long, simply pinnate, barren pinnae 3-8 inches long, about 1 inch broad, bluntly pointed, margin entire or toothed, 3-20 on each side, articulated at the base, texture subcoriaceous, glabrous, or nearly so on both sides, rachis often winged, fertile pinnae smaller, much contracted, about $\frac{1}{4}$ inch



STENOCHLÆNA SORBIFOLIA. (*L.*)

broad. *Acrostichum sorbifolium*, *Linn. Sp. Pl.* p. 1526. *Lomariopsis*, *Hook. Syn. Fil.* p. 412. *Bedd. F. B. I. t.* 192.

The Malay Peninsula, Tenasserim, Malacca.

Bedd. F. B. I. t. 210, is an abnormal bipinnate form of this plant in which the rachis of the pinnae is very broadly winged with small pinnules resembling the leaves of *Feronia elephantum*. *Lomaria limonifolia*, *Wall. Cat.* 35, is the same form.

(Also in Tropical America and West Indies; Fiji; Samoa;

New Caledonia ; Philippines ; Cochin China ; Mascareen Islands, and Tropical Africa.)

GENUS LXXXIII.—POLYBOTRYA. (*H. B. K.*)

(*Poly*, many ; *botrys*, bunch—in allusion to the fructification.)

Fronds pinnate, bipinnatifid or subbipinnate, the sterile not lomarioid in habit, generally viviparous, fertile much contracted; veins pinnate, all free; stipes adherent to the rhizome.

1. POLYBOTRYA APPENDICULATA. (*Willd.*) Rhizome thick, short-creeping, stipes and rachis scaly, scales linear, not adpressed; fronds pinnate, glabrous, the sterile ones viviparous at the apex; pinnae 25 to 50 pair, subopposite or alternate, oblong-lanceolate, obtuse, 2–3 inches long, $\frac{1}{2}$ an inch broad, rather deeply crenated with a setaceous bristle between each crenature, superior basal crenature the largest, inferior base cuneate and slightly unequal; veins not prominent, pinnate free; fertile fronds much contracted, pinnae much shorter than the sterile ones. *Willd. Sp. Pl.* 114. *Bedd. F. S. I.* t. 194. *Wall. Cat.* 28 and 2685.

Common throughout the Indian region.


(Also in Philippines and Hong Kong.)

The above description only relates to the type, but there are several varieties more or less permanent.

VAR. β MAJOR. Stipes and rachis very thick, $\frac{1}{2}$ inch or rather more in diameter, rough with dense adpressed scurfy scales; pinnae 1 inch in breadth, not auricled at the superior base or cuneate and unequal at the inferior; main veins very prominent and straight and costa-like veinlets more numerous and very prominent.

Sikkim; a very large fern, unlike any forms in Southern India or Ceylon.

VAR. γ ASPLENIIFOLIA. (*Bory.*) Rachis with copious linear patent scales, fronds seldom proliferous at the apex; pinnae very



LECTOTYPE

D. YBOTRYA APPENDICULATA. (Willd.) VAR. Y ASPLENIIFOLIA.

unequal sided, the inferior basal portion being much cut away, the upper side deeper cut than in the type. *P. aspleniifolia*, *Bory*, in *Belang. Voy. Bot.* 23, t. 3. *Bedd. F. S. I. t.* 195. *Acrost. Wightianum*, *Wall. Cat.* 2163.



N° 256
POLYBOTRYA APPENDICULATA.
(*Willd.*) VAR. δ *HAMILTONIANA.*

acters in cultivation.

South India, in moist evergreen forests on the western side, up to nearly 4,000 feet elevation. (Not found in North India.)

This can only be considered a variety of this variable species, it looks, however, very different when growing, being a much brighter green and a far prettier species than the type, it is constant to its char-


VAR. δ *HAMILTONIANA.* (*Wall.*) Fertile pinnæ interrupted,

the sori in bead-like clusters.

Wall. Cat. 29. *Bedd. F. B. I. t.* 111, the lower 3 figures.

Polyb. Helferiana (*Kze*), *Schott. Supp.* 2 p. 47. t. 114.

Chittagong, Assam; Tenasserim, and Malay Peninsula generally.



N° 257
POLYBOTRYA APPENDICULATA. (*Willd.*)
VAR. ϵ *COSTULATA.*

pinnules up to $3\frac{1}{2}$ inches long, and again deeply pinnatifid.

VAR. ϵ *COSTULATA.* (*Hook.*) Sterile pinnæ pinnatifid $\frac{1}{2}$ - $\frac{3}{4}$ down to the rachis; main veins costulate and prominent, the lowest pair of pinnæ often more or less bipinnate both in the sterile and fertile, sometimes half deltoid and deeply bipinnate at the base, the

Hook. Sp. Fil. p. 252. Bedd. F. B. I. t. 110, and 111 right hand top fig.

Tenasserim; Khasya; Jeypore Hills, West of Vizagapatam. This variety can at once be distinguished by the much more deeply pinnatifid pinnæ, and by its costulate main veins, but the bipinnate character is not so constant; some specimens from Birma (*vide Bedd. F. B. I. t. 110*) have the lowest pinnæ deeply bipinnate in both sterile and fertile; and some specimens from Khasya and Jeypore are only slightly bipinnate in the sterile only, whilst other specimens from Birma have the lowest pinnæ quite unchanged (*vide Bedd. F. B. I. t. 111, right-hand top figure.*)

VAR. ζ SUBINTEGRA. Pinnæ almost entire, of a very dull colour and more coriaceous than in the type, not at all or very slightly auricled at the superior base, and the inferior base rounded and not at all unequal. *Bedd. F. B. I. t. 111, left-hand top fig.*


Birma only.

GENUS LXXXIV.—ACONIOPTERIS. (*Presl.*)

(*Akon*, a point, and *pterus*; the veins forming pointed angles near the margin.)

As in *Elaphoglossum*, only the veins are combined near the margin by a straight or zigzag vein.

1. ACONIOPTERIS GORGONEA. (*Kaulf.*) Rhizome very short, with ovate-lanceolate obtuse, brown-red scales; stipes tufted, scarcely an inch long, being usually winged near the base by the decurrent frond; barren fronds simple entire narrowly elliptic, tapering at both ends, 3–5 inches long, by $\frac{3}{4}$ – $1\frac{1}{4}$ inch broad; glabrous, but with peltate flat scales, sparingly scattered on the surface beneath; midrib strong, texture firm, but diaphanous; veins parallel, above simple or furcate, the marginal vein more distinct, and further from the margin than in examples from Polynesia. *Kaulf. En. Fil. 63. Clarke, F. N. I. p. 578.*



J. D. COLEMAN & C. E. DORRIS

GYMNOPTERIS VARIABILIS. (Hook.)

Khasya, above Shaila, 2,500 feet elevation, lately discovered by Mr. Clarke, but only in a barren state, so it is doubtful whether it is a new species, though it must be closely allied.

(Also in the Sandwich and Society Islands.)

GENUS LXXXV.—GYMNOPTERIS. (*Bernh.*)

(*Gymnos*, naked ; *pteris* ; seeding naked.)

Veins copiously anastomosing with free veinlets in the areoles ; the primary veins costæform or obsolete ; fronds simple or pinnate, generally dimorphous the fertile much contracted, or rarely simple and uniform and bearing the sori on the contracted apex ; stipes adherent to the rhizome.

1. GYMNOPTERIS VARIABILIS. (*Hook.*) Rhizome more or less creeping, squamose, with ovate or lanceolate scales ; sterile fronds membranaceous to subcoriaceous, generally about 1 foot or a little more long and 1–2½ inches wide, ovate-lanceolate acuminate, often being decurrent on the stipe nearly to its base ; costules generally more or less distinct, sometimes very prominent, zigzag or quite straight ; areoles copious with free included veinlets ; fertile fronds contracted and narrow, and normally entirely covered with sori, but in some cases the broader fronds are dotted all over with large polypodioid sori. *Hook. Sp. Fil.* v. 277. *Bedd. F. B. I. t.* 272.

Sikkim, Bhotan, Assam, Khasya, Cachar, up to 4,000 feet elevation. South India (rare, compared with the variety *lanceolata*) in the inner mountainous regions, fronds very thin and membranaceous and decurrent nearly to the base of the stipe, main veins wavy, but very prominent, Jeypore Hills (Vizagapatam), texture thicker and colour deeper green, very decurrent, main veins very prominent and nearly straight, polypodioid sori over many of the broad fronds ; *Varma*.

VAR. β LANCEOLATA. (*Hook.*) Main veins none, or more or less indistinct. *Gymnopteris lanceolata*, *Hook. Sp. Fil.* v. 276. *G. Féei* (*Moore*), *Bedd. F. S. I. t.* 48.

Very common throughout the Western forests of the Madras Presidency and Bombay; Ceylon; Birma. Also in Chota Nagpore, and on Parasnath: in typical forms there are no main veins, and all the venation is indistinct, but other forms run too closely into *variabilis* for it to be considered a distinct species; the fronds are often quite rounded at the base, but at other times nearly as decurrent as in *variabilis*, the seeding is normally over the whole of the under surface of the fertile frond, but sometimes it is in a broad line on each side of the costa, leaving a considerable margin of the laminae without sori, or at other times the seeding is punctiform or grammitoid (*Bedd. F. B. I. t. 274*), or the upper half



GYMNOPTERIS MINUS. (*Mett.*)

of the frond is contracted and sori-ferous, as in *Gymnopteris spicata*, (*Hymenolepis* of authors). In South Canara and Coorg there are forms with both sterile and fertile fronds 3-lobed (*Bedd. F. B. I. t. 273*), and in Ceylon, pinnatifid forms (*Bedd. F. S. I. t. 211*) with often as many as five distinct pinnae on each side the rachis, with only a narrow wing, but as the ordinary form is sometimes mixed with these even on the same root they can only be considered abnormal forms, not distinct varieties.

VAR. γ AXILLARIS. (*Car.*) This is a name given to a variety with a long slender tortuous rhizome, which creeps up trees, but it scarcely differs otherwise, the main veins are less prominent than in *variabilis*, but more so than in *lanceolata*. *Car. Praelect. 1801, n. 582. Hook. Sp. Fil. v. 276. Bedd. F. B. I. t. 271.*

South India, in all the western forests; Plains of Bengal and Assam; Birma.

2. **GYMNOPTERIS MINUS.** (*Mett.*) Small, rhizome creeping.

ck as a sparrow's quill, scaly at the apex, scales lanceolate ; stipes
note, slender, 1-3 or in those of the fertile frond 4 inches long ;
rile fronds membranaceous, $1\frac{1}{2}$ -2 inches long, oblong or broad-
iceolate, costate, tapering into a stipe $2\frac{1}{2}$ inches long ; costules in-
stinct, areoles with a free included clavate veinlet, smaller towards
e margin ; fertile fronds 2 inches long, linear obtuse. *Metten. Fil.*
ort. Lips. p. 20. Hook. Sp. Fil. v. 277. Bedd. F. B. I. t. 116.


Khasya Hills, 2,000-4,500 feet, in river sands, perhaps only
small form of *variabilis* var. *lanceolata*.

3. GYMNOPTERIS WALLII. (*Baker.*) Rhizome slender, wide-



Nº 260.

GYMNOPTERIS METALLICA. (*Bedd.*)



Nº 261

GYMNOPTERIS SPICATA. (*L.f.*)

eeeping, scales minute, grey, lanceolate acuminate ; sterile fronds
near, subsessile, 8-9 inches long, under $\frac{1}{4}$ inch broad, narrowed to
oth ends, obscurely repand, texture thin but rigid, surfaces naked,
ight green ; midrib stramineous, veins slender, immersed, forming
single costal row of long areoles, usually with a free decurved vein-
t springing from the outer border ; fertile frond filiform, 6-8 inches

long, long-stalked, fructification often interrupted. *Baker in Hook. Syn. Fil. p. 523. Bedd. F. B. I. Sup. t. 389.*

Ceylon, southern provinces, Moorooa, rocky ravines.

4. GYMNOPTERIS METALLICA. (*Bedd.*) Fronds quite sessile, 3-7 inches long, up to nearly 1 inch broad, of a deep shining metallic blue; main veins not prominent and often not distinguishable from the others; fertile fronds only soriferous towards the apex. *Bedd. F. Sup. t. 390.*

Ceylon, Haycock Mountain near Galle, in dense forests on rocks, 2,000 feet elevation.

5. GYMNOPTERIS SPICATA. (*L. f.*) Rhizome woody, short, creeping; stipes 1-2 inches long, firm; fronds 6-18 inches long, $\frac{1}{2}$ -1 inch broad, the upper part for 4-6 inches, suddenly or gradually contracted and fertile, the edge entire, the lower part narrowed very gradually, texture subcoriaceous, surfaces naked; no distinct main veins, areoles uniform, hexagonal, with copious free, simple or forked veinlets; sori mixed with peltate sporangiasters. *Linn. Fil. Sup. 444, under Acrostichum. Gymnopteris, Presl. Tent. Pter. 244, t. 11. Acrostichum, Hook. Syn. Fil. 424. Hymenolepis spicata, Bedd. F. S. I. t. 46.*

North India, Sikkim and Bhotan, 4,000-7,000 feet elevation, Khasya 3,000-7,000 feet; South India, on the Western mountains; Ceylon.

(*Hymenolepis* has been proposed as a genus or subgenus for this species, as the fructification is on the contracted apex of the fronds, instead of being on a separate frond, the same peculiarity occurs constantly in other species, and I have similarly contracted fronds of *G. variabilis* var. *lanceolata*, which if separated from their rhizomes and other fronds could not be distinguished from this species, so I do not think the genus can hold good.)

6. GYMNOPTERIS QUERCIFOLIA. (*Retz.*) Rhizome stout, creeping, scales linear, hair-pointed; stipe of the barren frond 1-2 inches long, clothed with soft spreading ferruginous hairs; barren frond 3-4 inches

long, $1\frac{1}{2}$ –2 inches broad, the terminal pinna with blunt rounded lobes reaching sometimes $\frac{1}{4}$ of the way to the rachis, below this usually a single pair of small sessile ovate blunt auriculate lateral ones, texture thin, herbaceous, veins beneath hairy; main veins sometimes distinct to the edge, at other times hardly distinguishable from the other veins, with copious areoles, including free veinlets; fertile fronds with a terminal pinna, 1–2 inches long, $\frac{1}{2}$ inch broad, and a pair of smaller lateral ones, the stipe slender, 6–9 inches long, naked except at the base. *Acrostichum quercifolium*. *Retz. Obs. Bot.* 6, p. 39. *Hook. Syn. Fil.* p. 418. *Bedd. F. S. I. t.* 47.

Tinnevely and Travancore Hills, up to 2,000 feet elevation, Anamallays; Ceylon, about Colombo.

7. *GYMNOPTERIS FLAGELLIFERA*. (*Wall.*) Rhizome short, terrestrial, or scandent several feet up trees, strong, densely scaly with brown lanceolate scales at the extremities, and throwing out rootlets encircling the boughs; stipes often densely scaly, fronds simple or 3-foliate or pinnate with 5–11 pinnae, the terminal pinna often much prolonged and rooting at the apex; pinnae stalked entire or repand, or even lobed, 4–5 inches long by 2 inches broad; main veins very prominent to the margin, areoles copious with or without free veinlets; fertile pinnae more or less contracted, generally without any free veinlets in the areoles, wholly soriferous or partially covered with meniscioid sori; texture herbaceous, drying a dull brown colour. *Wall. Cat.* 25. *Clarke, F. N. I.* p. 579. *Pacilopteris flagellifera*, *Bedd. F. B. I. t.* 112.

North India, Sikkim and Assam, Cachar and Chittagong up to 3,000 feet elevation; Birma. Some of my Birma specimens have all the fronds simple, and are then hardly distinguishable from “*variabilis*” (type) except by the peculiar colour and scaly stipe, others have



GYMNOPTERIS QUERCIFOLIA. (*Retz.*)

the fronds all 3-foliate and the pinnæ often deeply lobed, particularly the two lower ones, these simple and 3-foliate fronds grow on the same rhizome, the broad sterile fronds are often partially seeded, particularly on the long terminal cauda. I have only seen pinnate fronds on North Indian examples.

Sir W. Hooker in his *Sp. Fil.* gives Malabar for a locality and quotes Rheede, Hort. Malab. xii. 11, which plate is *Drynaria quercifolia*; Pl. 19 is probably intended, which however is *Gymnopteris subcrenata*. On one of Wallich's sheets of this fern (25, No. 3) there is written, Rheede, Malab. xii. t. 19.

8. *GYMNOPTERIS SUBREPANDA*. (*Hook.*) Rhizome woody, wide-creeping; stipes of barren fronds stout erect, nearly naked; barren fronds varying from simple, 1 foot long, $1\frac{1}{2}$ –2 inches broad, to 2 feet long, 1 foot broad, copiously pinnate, with several linear-oblong entire or subrepand pinnæ on each side, which are sometimes 6–9 inches long, 2 inches broad, texture subcoriaceous, surfaces and rachis naked; main veins distinct nearly to the edge, with copious areoles, with free veinlets between them; fertile fronds like the others, but smaller. *Syn. Fil. p.* 419. *Hook. Sp. Fil. v. p.* 275. *Bedd. F. B. I. t.* 339.

Penang.

This is nearly allied to the more pinnate varieties of *flagellifera*. Dodgson's specimen from the Himalayas referred here in the Kew Herb. certainly belongs to *flagellifera*.

9. *GYMNOPTERIS TRICUSPIS*. (*Hook.*) Rhizome creeping, clothed with subulate setaceous ferruginous scales; stipes 1 foot long, $1\frac{1}{2}$ of the fertile frond; sterile frond subcoriaceous, subtriangular-caudate, 8–10 inches long, 10–12 inches wide at the base, including the spread of the lateral lobes, deeply trilobate or 4-lobed, lateral lobes 5 inches or more long, $1\frac{1}{2}$ inch wide, horizontally patent, intermediate one 6 inches and more long, 2 inches wide, all of them oblong-lanceolate acuminate, entire unicostate; venation manifest, costules distant, indistinct, connected by transverse irregular curved veins forming large areoles enclosing lesser ones which have copious

free branched and divaricating veinlets; fertile fronds much elongated but contracted, tripartite nearly to the base, segments scarcely $\frac{1}{2}$ an inch wide, linear strap-shaped acuminate, lateral ones 9-10 inches long, erect-subpatent, intermediate one a foot and more long, sori generally universal except on the costa, or sometimes the terminal lobe only is soriferous, or rarely the apex only of one or more lobes is contracted and soriferous as in *Gymnopteris spicata*. *Hook. Sp. Fil.* v. 272. *Bedd. F. B. I. t.* 53.


Sikkim, at Goke below Darjeeling, 1,500 feet elevation.

10. *GYMNOPTERIS CONTAMINANS*. (*Wall*) Rhizome thick, short creeping; stipes scaly, fronds glabrous, pinnate, 1-2 feet long, of which the stipe is about $\frac{1}{3}$; rachis generally more or less winged sometimes prominently, more or less scaly, pinnæ 11-30 alternate sessile or very shortly petiolate, lanceolate from nearly entire to slightly crenated or pinnatifid one-third down, with the segments serrated, terminal pinnæ often more or less elongated, generally viviparous (the lateral



N° 283. *GYMNOPTERIS TRICUSPIS*. (*Hook.*)

ones rarely so); primary veins rather distant more or less conspicuous two-thirds of the way to the margin, areoles broad, in 3-4 series, the costal ones without any veinlets, the others with generally several veinlets irregularly disposed, mostly joining with the areole above, but some free with clavate apices; fertile pinnæ much contracted, entire or sinuate pinnatifid according to the form of the sterile, generally wholly soriferous, rarely only partially contracted into seed. *Acrostichum contaminans*, *Wall. Cat.* 22, and *crispatum*, *Cat.* 24. *Pæcilopteris repanda*, *Bedd. F. S. I. t.* 202. *Semicordata*, *Baker, Syn. Fil.* 422. *Bedd. F. B. I. t.* 270. *Blumeana*, *Hook. Syn.*




GYMNOTERIS CONTAMINANS. (Wall.)

Fil. p. 423 (at least as far as the North Indian specimens are concerned).
Acros. terminans, Wall. Cat. 2168.

South India, most abundant in all the Western coast moist forests from the plains up to about 4,000 feet; East Bengal, the lower hills up to 3,000 feet elevation; Ceylon (repandum of *Thw. En.*); Birma, common.

11. *GYMNOPTERIS SUBCRENATA. (Hook. and Grev. under Acrostichum.)* Rhizome thick, fronds glabrous pinnate, 1-4 feet, of which the stipe is sometimes nearly half; stipes and rachis furnished with a few scales; sterile fronds, pinnæ 4-12 alternate or sub-opposite petiolate, broad lanceolate sinuate or waved, with a longish serrated acumination, terminal pinnæ much the longest (sometimes 2 feet long), proliferous at the apex; primary veins close, costate and conspicuous nearly to the margin, veinlets anastomosing pretty regularly at right angles, from which proceed one or two generally free veinlets with clavate apices; fertile fronds conform to sterile, but much contracted. *Hook and Grev. Ic. Fil. 110. Pæcilopteris terminans, Bedd. F. S. I. t. 203 (not, Wall.). P. Hookerianum, Moore, Thw. En. p. 380, prolifera. Hook. Ic. Pl. t. 681-2. Rheede, Hort. Malab. xii 19.*



GYMNOPTERIS SUBCRENATA.
 (Hook. and Grev.)

South India, not uncommon in the Western moist forests of the Madras and Bombay Presidencies, up to about 4,000 feet elevation; Ceylon, central provinces.

This fern is not known from North India (Mr. Clarke's Chittagong specimens being *costata*), it has generally fewer and larger pinnæ than *contaminans*, sometimes up to 2 inches broad, and they are never quite sessile, the venation differs considerably, the primary veins are much more conspicuous or closer together, the areoles narrow and generally with an acute apex, the lowest ones empty, the others with often only one free excurrent veinlet (and then the venation is quite that of *Meniscium*), there are, however, often two or sometimes three excurrent veinlets, which are very irregular, free or anastomosing amongst themselves, or with the superior areole.



GYMNOPTERIS COSTATA. (Wall.)

12. GYMNOPTERIS COSTATA (Wall.) Rhizome creeping, furnished with subulate scales; stipes up to $1\frac{1}{2}$ feet long; fronds up to 2 feet long or more, pinnate, pinnæ up to 14 inches long by 3 inches broad, petiolate, acuminate, margin entire, sinuate, or crenate; primary veins very prominent and generally much raised, close together, areoles numerous, but varying with the breadth of the pinnæ, costal ones small (sometimes obsolete) empty, several excurrent veinlets from all the other areoles which


are irregular, sometimes free and sometimes anastomosing. Wall. Cat. 26. Bedd. F. B. I. 113. Hook. Sp. Fil. v. 262, *deltigerum* and *undulatum*, Wall. Cat. 59 and 140. Bedd. F. B. I. 114 and 115, are only abnormal forms, such as occur more or less in nearly all the other species of *Gymnopteris* and cannot be recorded as varieties (Mr. Clarke has again transferred the former, which was *Meniscium* of Wallich, to that genus, but he now acknowledges that he was wrong, and that the venation is different to *Meniscium*, the same.

meniscioid sori being very common in flagellifera). Sir W. Hooker in his *Sp. Fil.* quite correctly described this species and its varieties; but he lumped subcrenata with contaminans, though he had formerly distinguished it, first under the name of subcrenata, and afterwards as flagellifera. Mr. Baker, not understanding the three species, lumped them all in his edition of the Synopsis; and Mr. Clarke, not knowing subcrenata, has puzzled pteridologists with a lot of varieties which do not exist; I have had subcrenata and contaminans (very common South Indian plants) for many years in cultivation, and subcrenata (not South Indian) I have seen in abundance in Birma, and know them to be three distinct species without any permanent varieties, though all three variable, as are nearly all other Gymnopteris.

North India, Nepal and Chittagong, up to 6000 feet elevation; Birma. (Some of the larger fronds turn very red in drying.) The meniscioid form (flagellifera) is common in Sikkim and Assam, the undulate form (formerly described as a genus, Jenkinsia) is, I believe, only from Birma.

Acros. virens. Wall. Cat. 1033, from Tavoy, is a blank sheet, so it is impossible to say whether the fern intended to be represented as this species or contaminans; it could not be subcrenata, as it does not grow in Birma.

13. *GYMNOPTERIS PRESLIANA.* (Hook.) Rhizome stout creeping, paleaceous with narrow attenuated lanceolate scales; stipes 6-10



GYMNOPTERIS PRESLIANA. (Hook.)

inches long, more or less scaly, as is the rachis and petioles, fronds 6–12 inches long, pinnated glabrous, pinnæ submembranaceous, remote, subopposite, petiolate, 2–4 inches long, $\frac{1}{2}$ an inch wide, narrow lanceolate acuminate at both ends, rachis more or less winged, specially towards the apex of the frond; fertile fronds generally longer stipitate, pinnæ much contracted, wholly or sometimes only partially fertile, costules indistinct, connected by generally 2 pairs of opposite veinlets, which meet at an acute angle, from which proceed a veinlet, which is either free or joined to the veins above, marginal veins free, terminating in a dot within the margin. *Hook. Sp. Fil.* v. 265. *Syn. Fil.* 420. *Bedd. F. B. I. t.* 269 (under *Pæcilopteris*.)

South India, Coorg, at the foot of the Bhagamandal ghat at no elevation, Concan (Law). I have only found this species in the single locality mentioned, but it is probably more common in the Bombay Presidency; it is of much thinner texture than *contaminans*, the pinnæ differently shaped, and the venation more that of *subcrenata*.


(Also in the Philippines.)

GENUS LXXXVI.—ACROSTICHUM. (L.)

(*Akros*, highest; *stichos*, order—fructification at the top of the frond.)

Veins uniform, copiously anastomosing, forming numerous subquadrangular areoles without free included veinlets, no main veins present; fronds pinnate, the upper pinnæ smaller and wholly sterile on the lower surface; stipes adherent to the caudex.

1. *ACROSTICHUM AUREUM*. (L.) Caudex erect; stipe 1–2 feet long, tufted, strong, erect, glossy; fronds 2–6 feet long, 1–2 feet broad, pinnate, the upper pinnæ fertile and slightly smaller than the barren ones which are usually stalked, ligulate-oblong, $\frac{1}{2}$ –3 inches broad, the apex acute or blunt, sometimes retuse with a mucro, the edge quite entire, the base subcuneate, texture coriaceous, rachis and surfaces glabrous; areoles very small and copious without free veinlets. *Linn. Sp. Pl.* 1525. *Bedd. F. S. I. t.* 204.



WILLD. C. 1800

ACROSTICHUM AUREUM. (Linn.)

Tidal backwaters throughout the Indian region.

(Also throughout the world in warm countries near the sea.)

.GENUS LXXXVII.—PHOTINOPTERIS. (*J. Sm.*)

(*Photeinos*, shining ; *pteris*, a fern.)

Veins copiously anastomosing, forming numerous areoles with free included veinlets, the primary veins costæform ; fronds pinnate, the upper pinnæ much contracted and wholly soriferous on the under side ; stipe adherent to the rhizome, pinnæ articulate with the rachis. (Very like *Drynaria* in its pinnæ and venation, but the vernation is adherent not articulate.)

1. PHOTINOPTERIS RIGIDA. (*Wall.*) Rhizome repent or scandent, paleaceous, with elongated brown subulate fringed scales mixed with hair-like ones in age, white glaucescent ; stipes short, nearly as thick as a writing pen ; fronds $1\frac{1}{2}$ –3 feet long, broad-lanceolate, very hard-coriaceous glossy, pinnate ; sterile pinnæ long-petiolate, distant 3–6 inches and more long, broad ovate, acuminate, the apex generally caudate and curved upwards ; venation manifest, elevated on both sides, costules subflexuose, united by transverse veins into four-angled areoles, which are again divided into lesser areoles including branched free clavate veinlets ; petioles articulated on the rachis and dilated into a large orbicular scutiform base ; sometimes these sterile pinnæ occupy the whole frond, more frequently the rachis is suddenly elongated at the apex, and bears 1–13 distant very contracted narrow-linear and fertile pinnæ, 5–12 inches long, soriferous except on the costa and on the slightly revolute margin. *Wall. Cat. n. 27. Hook. Sp. Fil. v. 281. Bedd. F. B. I. t. 211.*

Malay Peninsula ; Singapore.



(Also in the Malay Islands, and Philippines.)

2 PHOTINOPTERIS DRYNARIOIDES. (*Hook.*) Fronds sessile, several feet long, 1 foot or more broad, the upper part with numerous distant Lomarioid pinnæ, sometimes 1 foot long, 1 line broad, below



LICHTEN & CO. 1884

PHOTINOPTERIS RIGIDA. (Wall.)

N^o 270PHOTINOPTERIS DRYNARIOIDES. (*Hook.*)N^o 271PLATYCERIUM GRANDE. (*J. Sm.*)

this pinnatifid, with numerous close entire lanceolate acuminate lobes 4-6 inches long, $1\frac{1}{4}$ - $1\frac{1}{2}$ inch broad which reach down nearly to the rachis, all articulated, the lower ones passing down gradually into a mere undulated wing to the rachis, which is continued to its very base; texture very coriaceous; main veins raised, with transverse veinlets, and between them copious small areoles with free veinlets. *Hook. Syn. Fil. p. 425. Sp. Fil. v. 282. Bedd. F. B. I. t. 325.*

Malay Peninsula, Penang.

(Also in the Solomon Islands.)

GENUS LXXXVIII. — PLATYCERIUM. (*Desf.*)

(*Platys*, broad; *keros*, a horn —stag's-horn fern.)

Sori occupying a portion only of the under side of the disk of the fertile fronds and forming large, often reniform patches, frequently at the sinuses of the primary lobes, main veins costæform.

prominent, furcate, free or anastomosing, veinlets copiously anastomosing, forming large oblong areoles, usually with free included simple or forked veinlets; fronds dimorphous, the sterile sessile, oblique, reniform, round or elongated, the fertile dichotomously forked with stag-horn-like divisions, in texture and pubescence resembling *Niphobolus*; articulate with the caudex.

1. *PLATYCERIUM GRANDE*. (*J. Sm.*) Barren fronds very large, suborbicular, convex or the upper ones erect, deeply laciniated with spreading or inflexed divisions; fertile fronds 4–6 feet long, pendent in pairs, the disk broad-cuneate, with the sorus against the upper edge, occupying the disk only, with an elongated dichotomously forked division beyond it at each corner; both surfaces green and smooth. *J. Sm. in Hook. Journ. Bot.* iii. p. 402. *Hook. Syn. Fil.* p. 425. *Bedd. F. B. I. t.* 326. There is only one soriferous patch to each frond.


Singapore.

• (Also in Philippines and North Australia.)

2. *PLATYCERIUM WALLICHII*. (*Hook.*) Fronds ample, bifarious, at length glabrous; sterile ones imbricated below, irregularly sinuate-lobate, above elongated, dilated deeply and much dichotomously pinnatifid, segments patent-inflexed; fertile fronds geminate, very broadly flabellate-cuneate, twice dichotomous suddenly narrowed at the base, each primary division or segment bears a sorus on the disk (hence there are two sori on each frond), which is semicircular and very much produced in the sinus itself, the terminal segments beyond the soriferous disk are several times dichotomous and pendent, primary veins dichotomous, parallel elevated here and there, anastomosing into broad elongated areoles which are occupied by lesser ones with numerous free-branched included veinlets. *Hook. Sp. Fil.* v. 284. *Acrostichum alcicorne*, *Wall. Cat. n.* 19, (*not Linn.*) *Bedd. F. B. I. t.* 108.

Malay Peninsula, Tenasserim.

3. *PLATYCERIUM BIFORME*. (*Bl.*) Fronds ample, bifarious, at



PLATYCERIUM WALLICHII. (Hook.)

length glabrous; sterile ones imbricated, very thick and corky towards the base, subrotund but very varied in circumscription, obed and sinuated at the margin, coarsely-reticulate-venose; fertile fronds rather long-stipitate of great size, 5-15 feet long, from a subcuneate base, repeatedly dichotomous, the segments loriform, flaccid and pendent, fertile segments quite different from the rest, forming a large reniform, stipitate, shield-like receptacle, 6-8 inches or more in diameter, wholly soriferous beneath, except at the margin.

Bl. Fil. Jav. p. 14, t. 18. Hook. Sp. Fil. v. 285. Bedd. F. B. I. t. 109 and 224. A. fuciforme, Wall. Cat. n. 20.

Malay Peninsula, Tenasserim, Mergui, Singapore.
(Also in the Malay Islands and Philippines.)



N° 273

PLATYCERIUM BIFORME. (B. I.)

SUB-ORDER III.—OSMUNDACEÆ.


Capsule 2-valved, opening across the apex, furnished with a short horizontal wing; vernation circinate.

GENUS LXXXIX.—OSMUNDA. (Linn.)

(*Osmunder*, one of the names of Thor, a Celtic divinity.)

Fertile frond wholly, on the upper or middle portion, contracted, forming simple or compound sporangiferous panicles; veins forked, free; fronds pinnate or bipinnate, articulated with the rachis.

1. *OSMUNDA JAVANICA*. (B. I.) Stipes tufted, 6-12 inches long, firm erect, naked; fronds up to 3 feet long, 8-12 inches broad, simply pinnate, the lower or central pinnæ fertile; barren pinnæ 4-8 inches



6.3. 1971. 1. 1. 1971.

OSMUNDA JAVANICA. (Bl.)

ing, $\frac{3}{8}$ – $\frac{1}{2}$ inch broad, cuneate at the base, and often slightly stalked, the edge entire or sharply toothed; texture coriaceous, rachis and both sides glabrous; fertile pinnæ shorter, made up of numerous close but distinct oblong sessile clusters. *Bl. Enum. p. 252. Bedd. S. I. t. 77.*

Ceylon, about Newera Elya; South India, cultivated only (?); Malay Peninsula; Tenasserim, banks of rivers in the plains. This commonly in cultivation in Southern India, but I very much doubt it is wild; I have never found it myself, though I have specimens said to have been collected in the Western ghats.

(Also in Java.)



Nº 275.

OSMUNDA CLAYTONIANA. (L.)



Nº 276.

OSMUNDA REGALIS. (L.)

2. OSMUNDA CLAYTONIANA. (Lin.) Stipes tufted, 1 foot or more long, clothed with loose woolly ferruginous tomentum when young, at length glabrous; fronds 1–2 feet long, 8–12 inches broad, simply pinnate; pinnæ deeply pinnatifid, the uppermost and lowest barren, some of the intermediate fertile, barren ones lanceolate, 4–6 inches long, 1 inch or more broad, the lobes oblong, entire, texture

herbaceous, fertile pinnæ shorter, pinnules dense cylindrical. *Lin. Sp. Pl.* 1521. *Bedd. F. B. I. t.* 187.

Himalayas from Kashmir to Bhotan, 6,000–10,000 feet elevation. Khasya, 4,500–6,000 feet.

(Also in North America.)

3. OSMUNDA REGALIS. (L.)

Stipes tufted, 12–18 inches long, firm, erect, naked; fronds 2–4 feet long, 1 foot or more broad, bipinnate, the barren and fertile separate, or the frond barren below and fertile above, barren pinnæ 6–12 inches long, 2–4 inches broad, pinnules sessile or slightly stalked, 1–2 inches long, $\frac{1}{4}$ – $\frac{1}{2}$ inch broad, oblong, blunt, often unequal at the base, the edge finely serrulate, texture subcoriaceous, rachis and both sides naked, fertile pinnules cylindrical, forming a copious panicle. *L. Sp. Pl.* 152. *Bedd. F. S. I. t.* 76.

South India, common on the Western mountains at the higher elevations; North India, Kumaon, Bhotan, Khasya, 4,000–6,000 feet, (small, the fertile and barren fronds generally separate.)




№ 277.

SCHIZÆA MALACCANA. (Baker)

SUB-ORDER IV.—SCHIZÆACEÆ.

Capsule 2-valved, opening down the side, crowned by a complete operculiform ring; venation circinate.




SCHIZAEA DICHOTOMA. (Sw.)

GENUS XC.—SCHIZÆA. (*Smith.*)(*Schizo*, I cut ; the cut fertile segments.)

Capsules sessile in 2-4 rows which cover one side of close distichous spikes, which form separate fertile segments at the apex of the fronds ; veins free.

1. *SCHIZÆA MALACCANA*. (*Baker.*) Stipes dense, not distinguishable from the frond, which is 4-8 inches long, weak, flexuose, subterete, channelled in front not more than $\frac{1}{2}$ line thick, the barren

*SCHIZÆA DIGITATA*. (*Sw.*)

and fertile ones similiar ; fertile segment erect, often bilateral, $\frac{1}{2}$ inch deep with 3-6 slender spreading spikes on each side, the lowest 2-2 $\frac{1}{2}$ lines long. *Baker, Syn. Fil. p. 428. Bedd. F. B. I. t. 255.*

Malay Peninsula.

(Also in the Malay Islands and Philippines.)

2. *SCHIZÆA DICHOTOMA*. (*Sw.*) Stipes 6-18 inches long, firm, erect, glossy, channelled on the face above ; fronds fan-like in general outline, 6-9 inches each way, many times dichotomously forked, the ultimate divisions $\frac{1}{2}$ -1 inch broad, with 1 fertile segment to each, $\frac{1}{2}$ - $\frac{1}{3}$ inch long, the

rachis often curved with 4-10 close spreading spikes on each side. *Hook. Syn. Fil. p. 430. Bedd. F. S. I. t. 65.*

South India, on the Western mountains ; Malay Peninsula.

(Also in Tropical America and West Indies ; Australia, New Zealand, Polynesia, Philippines, and Mascareen Islands.)

3. *SCHIZÆA DIGITATA*. (*Sw.*) Fronds long, linear, grass-like,

10–15 inches long by 2–3 lines broad, bearing a digitate fertile crest at the apex ; crest 8–14, parted to the base, segments 1 inch long by a line broad ; sori in 4 series (*i.e.*, 2 series each side the costa).

Sw. Syn. Fil. 150, 380, *t.* 4. *Bedd. F. S. I.* 268.

North India, Khasya and Chittagong ; Ceylon ; Malay Peninsula.
(Also in the Malay Islands ; Fiji and Philippines.)

GENUS XCI.—ANEMIA. (*Sw.*)

(From *aneimon*, naked—the naked spikes.)

Capsules small, very abundant, forming a copiously branched panicle quite distinct from the leafy part of the frond ; fronds pinnate or bi-tripinnatifid ; veins free.

1. ANEMIA TOMENTOSA. (*Sw.*) Stipes 6–12 inches long strong, erect, clothed with deciduous ferruginous hairs ; fronds ternately divided, the two lateral branches fertile, the terminal one spreading, sterile bipinnate, with the pinnules variously lobed or pinnatifid, texture herbaceous, rachis and both surfaces pilose ; veins fine, flabellate. *Hook. Syn. Fil.* 433. *Anemia Wightiana* (*Gard.*), *Bedd. F. S. I. t.* 66.


South India, Nilgiris (Sispara ghat), Anamallays, Pulney Hills, Travancore Hills, 3,000–4,000 feet elevation.

(Also in Mexico, Peru and West Indies.)

GENUS XCII.—LYGODIUM. (*Sw.*)

(*Lygodes*, flexible—the climbing habit.)

Capsules solitary (or casually in pairs), in the axils of large imbricated clasping involucre, which form spikes either in separate pinnæ or in lax rows along the edge of the leafy ones ; fronds scandent, pinnæ conjugate palmate-lobed, pinnatifid or pinnate ; veins forked, free.



ANEMIA TOMENTOSA. (Str.)


1. *LYGODIUM CIRCINATUM*. (Sw.) Fully developed barren frond, bipartite into 2 palmate lobes or simply palmate, primary petiole so much reduced that the fork seems almost to spring from the main rachis, secondary petiole 1-2 inches long, firm, naked, pinnules digitate, with 5-6 long lanceolate lobes, reaching nearly down to the base, or once or even twice-forked, ultimate barren divisions 4-12 inches long, $\frac{1}{4}$ - $\frac{1}{2}$ inch broad, the fertile ones contracted sometimes so much so, that the lamina is nearly lost, the spikes 1-2 lines long, in close marginal rows, texture subcoriaceous, surfaces naked. Sw. *Syn. Fil.* 153. *Thw. En. Pl. Zey. p.* 379. *pedatum*, Sw. 154. *L. dichotomum*, Bedd. *F. S. I.* 62. *Wall. Cat.* 176. *Hook. Syn. Fil.* 437.

North India, Chittagong Hills; Ceylon, western, central, and southern provinces, up to 2,000 feet elevation; Malay Peninsula.

(Also in the Malay Islands; Philippines; Hong Kong and Chusan.)

2. *LYGODIUM MICROPHYLLUM*. (R. Br.) Fronds simply pinnate, pinnules petioled, 3-4 on each side of the zigzag rachis, with a terminal one which is more or less lobed, barren pinnæ ovate-oblong, blunt, the margin subentire or rarely somewhat lobed, the base rounded or cordate, fertile ones short, deltoid, with generally a very rounded apex, and a square base and lobed round the margin. *R. Br. Prod.* 162. *Wall. Cat.* 174. *L. scandens*, Bedd. *F. S. I. t.* 61.

South India, Malabar and West Coast generally, very common in the plains, and also in the Wynad, up to about 3,000 feet eleva-



N° 281
LYGODIUM CIRCINATUM. (Sw.)



LYGODIUM MICROPHYLLUM. (R. Br.)


tion; Ceylon, abundant; North India, Bengal plains, rare; Malay Peninsula.

(Also in the Malay Islands.)

3. *LYGODIUM FLEXUOSUM*. (Sw.) Fronds glabrous or slightly hairy, pair of fronds stipitate-pinnate with the pinnules again pinnate or variously lobed, or subpalmate, all serrulate; sori protruding from the margin; texture subcoriaceous. Sw. Syn. Fil. 153. Bedd. F.



LYGODIUM FLEXUOSUM. (Sw.)



LYGODIUM POLYSTACHYUM. (Wall.)

S. I. t. 63. *L. pinnatifidum*, Sw. 153. *L. longifolium*, Wall. Cat. 175.

South India, on both sides of the Madras Presidency, common up to about 4,000 feet elevation; North India, plains, and up to 5,000 feet on the Himalayas; Ceylon; Malay Peninsula.

(Also in the Malay Islands; Philippines; North Australia, and Tropical Africa.)

4. *LYGODIUM JAPONICUM*. (Sw.) As in *flexuosum*, only that

the pinnæ are much smaller, with the pinnules smaller and finely cut, the fertile ones often so contracted that there is little or no lamina present. *Sw. Syn. Fil.* 154. *Bedd. F. S. I. t.* 64. *Wall. Cat.* 2201.

North India, abundant, extending west to Kashmir, up to 5,000 feet elevation; South India, Western mountains, rare. It is probably only a form of *flexuosum*.

(Also in Japan, China, Australia, Malay Islands and Philippines.)

5. *LYGODIUM POLYSTACHYUM*. (*Wall.*) Stem creeping, slightly pilose; fronds conjugate pinnate, membranaceous, pinnæ petiolate, furnished with a tufted gland at the apex of the petiole, deltoid-ovate to lanceolate, glandular-pilose on the rachis of the pinnæ costa and veins, pinnatifid more than half-down to the costa, segments with a rounded apex, entire or slightly crenate; costa of the pinnæ and central vein (or costule) of segments flexuose, veinlets simple or forked; fertile segments contracted. *Hook. Syn. Fil.* 438. *Bedd. F. B. I. t.* 300.

Malay Peninsula, Tenasserim.


SUB-ORDER V.--MARATTIACEÆ.

Capsule opening by a slit down one side or a pore at the apex, without a wing, usually joined together in concrete masses (*synangia*); vernation circinate.

GENUS XCIII.—ANGIOPTERIS. (*Hoffm.*)

(*Angio*, open; *pteris*, fern—the open sporangia.)

Capsules opening by a slit down the side, sessile, very close to one another, but not concrete, arranged in a linear-oblong or boat-shaped band of sporangia near the edge of the frond; veins simple or forked, free; fronds very large, bipinnate, springing from between two fleshy stipulæform appendages; the base of the stipes clavate, pseudo-articulate with the axis; pinnæ and pinnules articulate with the rachis.



1. *ANGIOPTERIS ERECTA*. (*Hoffm*) Caudex erect, often 2 feet thick, and as much or more in height ; fronds 6–20 feet long, pinnae 1–3 feet long, spreading, the lowest the largest, rachis swollen at the base, pinnules 4–12 inches long, $\frac{1}{2}$ –1 $\frac{1}{2}$ inch broad, linear oblong, sessile or shortly stalked, the apex acuminate, the edge entire or toothed, particularly towards the apex, texture herbaceous to subcoriaceous, glabrous, shining ; veins subparallel ; sori of 8–15 capsules. *Hoffm. Schk. Krypt. Gew. t. 151. Bedd. F. S. I. t. 78.*

Throughout the Indian region up to 7,000 feet elevation.

(Also in Japan, Tropical Australia, New Caledonia, Madagascar, and Polynesia.)


GENUS XCIV.—MARATTIA. (*Sm.*)

(After Maratti of Tuscany, a writer on Ferns.)

Capsules sessile or stalked, 4–12 concrete in boat-shaped synangia, which consist of two opposite rows of capsules and open by slits down their inner faces, with or without an inferior involucre ; veins simple or forked, free ; fronds bi-tripinnate, large, springing from between two fleshy stipulæform appendages (which sometimes assume the character of abnormal fronds) ; pinnules articulate with the rachis.

1. *MARATTIA FRAXINEA*. (*Smith.*) Stipes 1–2 feet long, 1–1 $\frac{1}{2}$ in. thick, smooth deciduously scaly or swollen in the lower part ; fronds up to 15 feet long, bipinnate, or sometimes tripinnate ; pinnae 1–2 feet long, pinnules oblong-lanceolate, 4–6 inches long $\frac{1}{2}$ –1 $\frac{1}{2}$ inches broad, the apex acuminate, the edge generally serrate, more rarely entire, the base cuneate, or slightly rounded, texture rather coriaceous, glabrous, the rachis of pinnae sometimes slightly winged ; synangia submarginal, in close rows, the receptacle linear with 6–12 capsules on each side, an obscure fimbriated inferior involucre often present. *Hook. Syn. Fil. 440. Bedd. F. S. I. t. 79.*

South India, Western forests of the Madras Presidency, 4,000–6,000 feet elevation (not nearly so common as *Angiopteris* which it



M. J. B. P. 1971.

MARATTIA FRAXINEA. (Sm.)

much resembles in general habit) ; Ceylon, central provinces, above 5,000 feet elevation ; Malay Peninsula.

(Also all round the world in the tropics and a little beyond in the southern zone.)

GENUS XCV.—KAULFUSSIA. (*Blume.*)

(After Kaulfuss, of Halle, a writer on Ferns.)

Capsules sessile, 10–15, quite concrete, in raised circular masses, which are hollow in the centre, with the oblong apertures on the inner face ; veins costæform, parallel veinlets copiously anastomosing with free venules in the areoles ; fronds 2–3 feet high, long stipate palmately lobed, and springing from two short fleshy stipulæform appendages ; lobes of the fronds oblong elliptical ; a genus of a single species.

1. KAULFUSSIA ÆSCULIFOLIA. (*Bl.*) Stipes 12–18 inches long, herbaceous, auricled at the base ; fronds digitate (like a chesnut leaf), or ternate ; the central pinnæ the largest, oblong, spathulate, 6–12 inches long, 3–4 inches broad, the others smaller, margins entire or lobed, texture fleshy-herbaceous ; sori copious scattered. *Bl. En. Pl. Jav. Fil.* 260. *Bedd. F. B. I. t.* 185, (free included veinlets not shown). *K. assamica*, *Griff. Not.* l. ii. 628.

North India, Assam, Cachar, Chittagong Hills, 250 feet elevation.

(Also in the Malay Islands and Philippines.)


SUB-ORDER VI.—OPHIOGLOSSACEÆ.

Capsule deeply 2-valved, opening down the side nearly to the base, without a ring ; vernation erect ; terrestrial or epiphytic.

GENUS XCVI.—OPHIOGLOSSUM. (*L.*)

(*Ophis*, a snake ; *glossa*, a tongue.)

Capsules sessile, arranged in two rows, forming a narrow close spike, which arises from the base or centre of the barren segment ;




Б. С. ГИТЕНКО, 1957, 1958, 1959

KAULFUSSIA AESCULIFOLIA. (Bl.)

rarely distinct, rising direct from the corm ; veins reticulated ; fronds simple entire, rarely palmate.

1. *OPHIOGLOSSUM VULGATUM* (L.) Rhizome not tuberous, short, or elongated, producing annually 1-2 fronds ; fronds 6-9 inches long, the sterile division generally placed about the middle 2-4 inch long, $\frac{1}{4}$ -2 inches broad, ovate or ovate-oblong, without a



Nº 288.

distinct haft, texture stouter than in the others, the midrib usually indistinct ; fertile spike 1 inch long or rather more, on a peduncle 2-4 inches long, and considerably overtopping the sterile division when fully mature. *Lin. Sp. Pl.* 1518.

Sikkim, Goke, 4,000 feet elevation, Rungait camp, below Darjeeling, 2,000 feet elevation.

(Widely spread in Europe ; Africa, and its eastern islands ; America ; Japan ; Australia ; New Zealand, and Sandwich Islands.)

2. *OPHIOGLOSSUM NUDICAULE* (L. f.) Rhizome small, slightly tuberous ; fronds 1 inch or more long, the sterile division placed not far from the base, $\frac{1}{4}$ -1 inch long, 2-5 lines broad, linear to ovate, without a haft, or with only a slight one, the texture thin but with no evident costa and veins not distinct ; fertile spike $\frac{1}{2}$ inch long, the peduncle often 2 inches long, very slender. *Sw. Syn. Fil.* t. 4. *O. parvisolium*, *Hook. and Grev. Bedd. F. S. I.* t. 71.

South India, Anamallay Forests, 2,500 feet elevation, and elsewhere on the Western mountains ; Malay Peninsula.

(Also in America from United States southward to Brazil, New Caledonia, and Tropical Africa.)

3. *OPHIOGLOSSUM FIBROSUM*. (Schum.) Rhizome a large round white bulb with numerous fibrous rootlets; stipe very short with the sterile division close to its base, the latter $1\frac{1}{2}$ –2 inches long, by $\frac{1}{2}$ inch broad, lanceolate, acute or obtuse, the texture thick and opaque; midrib prominent, venation indistinct (except when dried), fertile spike $1\frac{1}{2}$ –2 inches long on a spike 4–5 inches long. Schum. *K. Dansk. Vid. Afh.* iv. 226. *O. Wightii*, Hook and Grev. *Bot. Mis.* iii. 218. *O. brevipes*, Bedd. *F. S. I. t.* 72.

South India, Anamallay teak forests, 2,000–3,000 feet elevation.

(Also in Tropical Africa, Guinea Coast; Ascension Island.)

4. *OPHIOGLOSSUM RETICULATUM*. (L.) Rhizome cylindrical, elongated with many fibrous rootlets; fronds 6–12 inches long, the sterile division placed a little below the middle, $1\frac{1}{2}$ –3 inches long, by 1–2 inches broad, with a distinct haft, and distinctly cordate at the base, the apex blunt or acute, texture thin; veins prominent, but usually no midrib; fertile spike 1 inch or more long on a slender peduncle, 2–4 inches long. Linn. *Sp. Pl.*

1518. Hook. *Syn. Fil. p.* 446. Bedd. *F. S. I. t.* 70. *O. cordifolium* (Roxb.), Wall. *Cat.* 47.


South India, Nilgiris and Anamallays, 2,000 feet elevation and upwards; Ceylon, Newera Elya; Himalayas; Malay Peninsula.

(Also in Malay Islands; Polynesia; Tropical America; Africa, Mascareen Islands; Philippines.)



OPHIOGLOSSUM FIBROSUM. (Schum.)

5. *OPHIOGLOSSUM PENDULUM*. (L.) Epiphytic on trees;




18. PAVILLON & P. PAVILLON

OPHIOGLOSSUM RETICULATUM. (L.)

fronds pendulous, ribbon-like, without a distinct stem, 1-3 feet long, 1-3 inches broad, simple or forked, texture fleshy; no midrib and the veining not distinct; fertile spike single, arising low down on, but above the base of, the sterile segment, 2-6 inches long, on a peduncle shorter than itself. *Linn. Sp. Pl.* 1518. *Bedd. F. S. I.* t. 269.

Ceylon, at no great elevation; Malay Peninsula; Assam (?)

(Also in Malay and Polynesian Islands; Mauritius; Australia; Philippines.)



OPHIOGLOSSUM PENDULUM. (L.)


GENUS XCVII.—HELMINTHOSTACHYS. (*Kaulf.*)

(*Helminthos*, a worm; *stachys*, a spike.)

Capsules in long crested clusters which form a long loose spike; veins forked, free; fertile spike rising from the base of the leafy segment; fronds stipate, sterile segments foliaceous, digitate. A genus of a single species.

1. HELMINTHOSTACHYS ZEYLANICA. (*Linn. under Osmunda.*)

Rhizome thick, fleshy, creeping; stipes often 1 foot long, barren segment palmately pinnate, often in three principal divisions which are stalked, and again forked or pinnate, the ultimate divisions linear-oblong, 3-4 inches long, $\frac{1}{4}$ -1 inch broad, the apex acuminate, the edge slightly toothed or entire, texture herbaceous; fertile spike solitary, arising from the base of the barren segment, 3-4 inches long, $\frac{1}{4}$ inch broad, the firm peduncle about as long as the fructification. *Linn. Sp. Pl.* 1519. *Hook. and Bauer, Gen. Fil.* t. 48B. *Bedd. F. S. I.* t. 69.



A. B. PIERCE & CO. DRAUGHTSMEN.

HELMINTHOSTACHYS ZEYLANICA. (Linn.)

South India, Western forests in swampy places up to 3,000 feet elevation ; Ceylon, about Colombo and other parts of Western and Southern provinces ; North India, Bengal plains to Assam and Cachar ; Malay Peninsula.

(Also in Tropical Australia, Malay Islands, Philippines and New Caledonia.)

GENUS XCVIII.—BOTRYCHIUM. (*Sw.*)


(*Botrys*, a bunch ; fructification like a bunch of grapes.)

Capsules sessile, arranged in two rows, on the face of spikes which form a compound panicle ; veins forked, free ; fronds erect, the sterile segments foliaceous, deltoid, bi-tripinnatifidly compound, rarely pinnate ; fertile segments rachiform, compound paniculate.

1. BOTRYCHIUM LUNARIA. (*L. under Os-munda.*) Rhizome small, scarcely thickened, enclosed by brown sheaths furnished with stoutish fleshy brittle branched roots ; stipes erect, smooth, cylindrical, hollow, succulent, vernation plicate or folded straight, the fertile branch clasped by the sterile before unfolding, fronds solitary, 3-10 inches high, firm, stout fleshy, sterile branch oblong, pinnate smooth, pinnæ 4-7 pairs flabellate or lunate, the margins crenate (rarely partially fertile) fertile branch pinnate or bipinnate ; venation (barren pinnæ) flabellately-furcate, *i.e.* the vein enters at the base and is repeatedly forked, veins not quite extending to the margin. *Linn. Sp. Pl.* 1519. *Sw. Syn. Fil.* 171. *Bedd. F. B. I. t.* 208.

North India, Sikkim, Lachen, 11,000-13,000 feet elevation, Kumaon, 12,000 feet.

(Also the arctic and cold temperate zone, extending to South Europe ; Patagonia ; Australia.)



Nº 293.

BOTRYCHIUM
LUNARIA. (*L.*)

2. BOTRYCHIUM DAUCIFOLIUM. (*Wall.*) Stipes stout, 6-12



A. K. G. & C. G. G. G.

BOTRYCHUM DAUCIFOLIUM. (Wall.)

inches long, the sterile portion with a stalk, 1-6 inches long, *i.e.* prolonged beyond the fertile spike, 6-12 inches each way; deltoid, tripinnatifid or tripinnate; the lower pinnæ much the largest; segments lanceolate-oblong, $\frac{1}{4}$ - $\frac{3}{8}$ inch broad, finely toothed; fertile segments about equalling the sterile segment when mature; panicle 2-4 inches long tripinnate, not very close. *Wall. Hook. and Grev. Ic. Fil. t. 161.* *B. subcarnosum*, *Wall. Cat. 49.* *Bedd. F. S. I. t. 69.*

Throughout the Indian region, up to 8,000 feet elevation.

(Also in Java and Polynesia.)

3. *BOTRYCHIUM VIRGINIANUM*. (*L. under Osmunda.*)

Stipes 3-18 inches long, sterile portion not prolonged beyond the fertile spike 4-12 inches each way; deltoid, quadripinnatifid, lower pinnæ much the largest, pinnules ovate-oblong, close cut down to a broadly winged rachis into finely cut linear-oblong segments, both sides naked or slightly hairy; fertile branch of the rachis springing from the base of the sterile portion (*i.e.* sterile portion, sessile), or from the middle of it. *Linn. Sp. Pl. 1519.* *Sw. Syn. Fil. 171.*



N° 295.

BOTRYCHIUM VIRGINIANUM. (*Sw.*)
VAR. LANUGINOSUM.

VAR. β LANUGINOSUM. (*Sp. Wall. Cat. 48.*) Fertile branch arising from the middle of the sterile portion. *Bedd. F. S. I. t. 67.* *Moore, Ind. Fil. 213.* *Hook. Gard. Ferns, t. 29.* In the typical American plant, the fertile branch arises from the base of the sterile portion, and the latter is quite glabrous; in the Indian plant the fertile branch always springs from well above the base, and the sterile

portion is more or less hairy, they are, however, only considered varieties.


South India, at the higher elevations on the Western mountains (only appearing in the rains) ; Ceylon, about Newera Elya ; North India, on the Himalayas, Kumaon to Bhotan, 5,000–8,000 feet elevation, Khasya, 4,000–6,000 feet elevation.

(The typical plant is widespread in Europe, America and Japan.)





LEUCOSTEGIA MULTIDENTATA. (Wall.)



H. C. GATTELL & CO. PHOTODUPLICATORS

MICROLEPIA MAJUSCULA. (Lowe.)

ADDITIONS AND CORRECTIONS.

At page 16, after No. 10, insert :—

ALSOPHILA KINGI (Clarke MS. in Kew Herb.), a tree fern, 10 feet high, main rachis purplish shining glabrous, rachis of pinnæ purplish glabrous and somewhat glaucous, pinnæ 20-24 inches long, pinnules about 3 inches long by half inch broad, coriaceous, of a blueish glaucous tint, the rachis very scaly beneath and hairy above, cut down nearly to the base into oblong crenated segments, the midrib much raised and very prominent (particularly in the fertile pinnules) and scaly ; veins once forked from a little above the base, or more rarely simple, receptacles much raised and very prominent.

Malay Peninsula at Laroot, in Perak, at 5,000 feet elevation, lately discovered by Dr. King's collectors.

At page 20, line 15, for 900—1,200, *read* 9,000—12,000.

At page 28, for plate 14, *read* plate 15.

At page 29, for plate 15, *read* plate 14.

At page 292, 9th line from bottom, for *This variety*, *read The variety*.

At page 295, 11th line from bottom, for *rugulosum Labill*, *read rugosulum Labill*.

At page 439, 4th line from bottom, for *as it read*, *as that*.



INDEX TO THE PLATES.

PLATE		PAGE	PLATE		PAGE
GLEICHENIACEÆ.			19	Trichomanes digitatum ...	39
1	Gleichenia linearis ...	3	20	„ Kurzii ...	40
296	„ glauca ...	xv	21	„ javanicum ...	45
POLYPODIACEÆ.			<i>Davallieæ.</i>		
<i>Cyatheaæ.</i>			22	Humata parallela ...	47
2	Cyathea sinuata ...	5	23	„ pedata ...	49
3	„ spinulosa ...	7	24	Leucostegia assamica ...	51
297	„ Hookeri... ...	xvi	25	„ pulchra ...	53
4	Amphicosmia Walkeræ ...	9	25	„ nodosa ...	55
5	Alsophila Oldhami ...	13	298	„ multidentata ...	473
6	„ crinita ...	15	27	Prosaptia Emersoni ...	56
7	Diacalpe aspidioides ...	17	28	„ contigua ...	57
8	Matonia pectinata ...	19	29	Davallia triphylla ...	58
<i>Dicksonieæ.</i>			30	„ solida ...	59
9	Struthiopteris orientalis ...	21	31	„ bullata ...	63
10	Woodsia elongata ...	23	32	Microlepia Hookeriana ...	62
11	Peranema cyatheoides ...	25	33	„ platyphylla ...	65
12	Dennstædtia scabra ...	26	299	„ majuscula ...	474
15	„ appendiculata ...	29	34	Stenoloma chinensis ...	69
13	Cibotium Barometz ...	27	35	Cystopteris setosa ...	71
<i>Hymenophylleæ.</i>			<i>Lindsayeæ.</i>		
14	Hymenophyllum parvifo- lium ...	28	36	Lindsaya cultrata... ...	73
16	„ exsertum ...	31	37	„ scandens ...	74
17	„ Neesii... ...	35	38	„ divergens ...	76
18	Trichomanes parvulum ...	38	39	Schizoloma lobata, var. malabarica ...	78
			40	„ cordata ...	79
			41	„ ensifolia ...	81

PLATE		PAGE	PLATE		PAGE
	<i>[Pterideæ.</i>				
42	<i>Adiantum Parishii</i>	82	73	<i>Asplenium Wightianum</i>	146
43	„ <i>lunulatum</i> , var.		74	„ <i>tenerum</i>	147
	„ <i>Mettenii</i>	83	75	„ <i>Zenkerianum</i>	148
44	„ <i>caudatum</i>	85	76	„ <i>paradoxum</i>	152
45	„ <i>venustum</i>	87	77	„ <i>heterocarpum</i>	155
300	„ <i>hispidulum</i>	ii	78	„ <i>tenuifolium</i>	160
46	<i>Cheilanthes mysorensis</i>	90	79	„ <i>rutæfolium</i>	162
47	„ <i>varians</i>	91	80	<i>Athyrium falcatum</i>	164
48	„ <i>argentea</i>	95	81	„ <i>selenopteris</i>	167
49	<i>Onychium auratum</i>	97	82	„ <i>brevisorum</i>	171
50	<i>Cryptogramme crispa</i>	99	83	„ <i>fimbriatum</i>	173
51	<i>Pellæa Stelleri</i>	100	84	<i>Diplazium lanceum</i>	175
52	„ <i>concolor</i>	100	85	„ <i>porrectum</i>	176
53	„ <i>Boivini</i>	103	86	„ <i>bantamense</i>	177
54	„ <i>falcata</i>	104	87	„ <i>longifolium</i>	179
55	<i>Pteris longifolia</i>	105	88	„ <i>asperum</i>	183
56	„ <i>dactylina</i>	108	89	„ <i>polypodioides</i>	185
57	„ <i>Griffithii</i>	109	90	„ <i>umbrosum</i> , var.	
58	„ <i>semipinnata</i>	110		„ <i>australe</i>	189
59	„ <i>patens</i>	113	91	„ <i>umbrosum</i> , var.	
60	<i>Campteria Kleiniana</i>	117		„ <i>multicaudatum</i>	190
61	<i>Doryopteris ludens</i>	119	92	<i>Anisogonium cordifolium</i>	191
62	<i>Litobrochia incisa</i>	121	93	„ <i>heterophlebium</i>	192
63	<i>Ceratopteris thalictroides</i>	121	94	„ <i>esculentum</i>	193
64	<i>Lomaria Patersoni</i>	126	95	<i>Hemidictyum Ceterach</i>	194
65	<i>Plagiogyria adnata</i>	128	96	„ <i>Finlaysonia-</i>	
	<i>Blechnææ.</i>			„ <i>num</i>	195
66	<i>Blechnum orientale</i>	131	97	<i>Allantodia javanica</i>	196
67	<i>Blechnidium melanopus</i>	134	98	<i>Actiniopteris dichotoma</i>	198
68	<i>Woodwardia radicans</i>	136		<i>Aspidicæ.</i>	
69	<i>Doodia dives</i>	138	99	<i>Didymochlæna lunulata</i>	200
	<i>Asplenicææ.</i>		100	<i>Mesochlæna polycarpa</i>	202
70	<i>Thamnopteris Grevillei</i>	140	101	<i>Polystichum semicordatum</i>	201
71	<i>Asplenium ensiforme</i>	142	102	„ <i>auriculatum</i>	205
72	„ <i>alternans</i>	143	103	„ <i>ilicifolium</i>	206
			104	„ <i>aculeatum</i> , var.	
				„ <i>rufo-barbatum</i>	208

INDEX TO THE PLATES.

479

PLATE	PAGE
105 Polystichum Prescottianum	210
106 Cyrtomium falcatum, var.	
caryotideum	213
107 Aspidium singaporianum	214
108 " vastum	... 214
109 " subconfluens	215
110 " subtriphyllum	216
111 " variolosum	... 217
112 " heterocarpum	216
113 " decurrens, var.	
minor	... 221
114 Pleocnemia Trimeni	... 224
115 " membranifolia	226
116 " Clarkei	... 227
117 " leuzeana	... 228
118 Lastrea coniiifolia...	... 231
" aristata	... 231
119 " cuspidata	... 232
120 " hirtipes	... 233
121 " calcarata, var.	
ciliata	... 236
122 " Beddomei	... 239
123 " Thelypteris	... 242
124 " syrmatica	... 243
125 " flaccida	... 244
126 " Brunoniana	... 245
127 " barbigera	... 247
128 " odontoloma	... 247
129 " Filix-mas, var. nidus	249
130 " " var. cochleata	251
131 " sparsa, var. obtusissima	253
132 " spectabilis	... 256
133 " deparioides	... 258
134 " Parishii	... 259
135 " scabrosa	... 263
136 " ferruginea	... 265
137 Nephrodium otaria	... 268
138 " cucullatum	... 271
139 " aridum	... 273

PLATE	PAGE
140 Nephrodium urophyllum	274
141 " moulmeinense	275
142 " Arbuscula	... 276
143 " truncatum	... 281
144 Nephrolepis cordifolia	... 283
145 " ramosa	... 285
146 Oleandra neriiformis	... 286
147 " Wallichii	... 287

Polypodiæ.

148 Phegopteris Scottii	... 289
149 " auriculata	... 291
150 " rufescens	... 293
151 " Dryopteris	... 294
152 " ornata	... 294
153 Goniopteris prolifera	... 297
154 Dictyopteris Barberi	... 299
155 " chattagramica	300
156 " difformis	... 301
157 Polypodium parasiticum	.. 303
158 " Wallii	... 304
159 " hirtellum	... 306
160 " cornigerum	... 306
161 " cucullatum	... 307
162 " trichomanoides	308
163 " glandulosum	309
164 " Thwaitesii	... 310
165 " decorum	... 311
166 " obliquatum	... 312
167 " repandulum	... 313
168 " subfalcatum	... 314
169 " dareæforme	... 315
170 Goniophlebium amænum	316
171 " erythrocarpum	317
172 " molle	... 321
173 " subauriculatum	323
174 " argutum	... 323
175 " verrucosum	... 324
176 Niphobolus adnascens	... 326

PLATE		PAGE	PLATE		PAGE
177	<i>Niphobolus pannosus</i> ...	328		<i>Grammitideæ.</i>	
178	" <i>stigmatosus</i> ...	329	213	<i>Notholaena Marantæ</i> ...	374
179	" <i>fissus</i> ...	330	214	<i>Monogramme paradoxa</i> ...	376
180	" <i>flocculosus</i> ...	331	215	<i>Leptogramme Totta</i> ...	378
181	" <i>Gardneri</i> ...	332	216	" <i>aurita</i> ...	379
182	" <i>penangianus</i> ...	332	217	" <i>opaca</i> ...	379
183	" <i>mummulariæ-</i> <i>folius</i> ...	333	218	<i>Stegnogramme aspidioides</i>	381
184	<i>Dipteris Wallichii</i> ...	335	219	<i>Gymnogramme Andersoni</i>	382
185	" <i>Horsfieldii</i> ...	336	220	" <i>leptophylla</i> ...	383
186	" <i>Lobbiana</i> ...	337	221	" <i>microphylla</i> ...	384
187	<i>Dynaria coronans</i> ...	337	222	<i>Syngramme fraxinea</i> ...	385
188	" <i>splendens</i> ...	339	223	" <i>vestita</i> ...	387
189	" <i>propinqua</i> ...	340	224	" <i>alismæfolia</i> ...	388
190	" <i>mollis</i> ...	342	225	<i>Selliguea Féei</i> ...	388
191	" <i>quercifolia</i> ...	343	226	" <i>Hamiltoniana</i> ...	390
192	" <i>rigidula</i> ...	344	227	" <i>elliptica</i> ...	391
193	<i>Placopeltis rostrata</i> ...	346	228	<i>Loxogramme involuta</i> ...	394
194	" <i>linearis</i> ...	347	229	" <i>avenia</i> ...	395
195	" <i>sinuosa</i> ...	349	230	<i>Brainea insignis</i> ...	396
196	" <i>longifolia</i> ...	350	231	<i>Meniscium triphyllum</i> ...	398
197	" <i>lanceolata</i> ...	352	232	" <i>Thwaitesii</i> ...	399
198	" <i>rhyncophylla</i> ...	354	233	" <i>salicifolium</i> ...	399
199	" <i>Griffithiana</i> ...	355	234	" <i>cuspidatum</i> ...	400
200	" <i>ovata</i> ...	356	235	<i>Antrophyum reticulatum</i>	401
201	" <i>punctata</i> ...	358	236	" <i>plantagineum</i> ...	402
202	" <i>hemionitidea</i> ...	359	237	" <i>latifolium</i> ...	403
203	" <i>pteropus</i> ...	360	238	<i>Vittaria elongata</i> ...	405
204	" <i>pteropus</i> , var. <i>minor</i> ...	361	239	" <i>sikkimensis</i> ...	406
205	" <i>hastata</i> ...	362	240	" <i>sulcata</i> ...	408
206	" <i>incurvata</i> ...	364	241	" <i>scolopendrina</i> ...	408
207	" <i>insignis</i> ...	365	242	<i>Tænitis blechnoides</i> ...	409
208	" <i>nigrescens</i> ...	367	243	<i>Drymoglossum carnosum</i>	411
209	" <i>dilatata</i> ...	368	244	" <i>piloselloides</i> ...	412
210	" <i>juglandifolia</i> ...	369	245	<i>Hemionitis arifolia</i> ...	414
211	" <i>Lehmanni</i> ...	371	246	" <i>Griffithii</i> ...	415
212	" <i>himalayensis</i> ...	372		<i>Acrosticheæ.</i>	
			247	<i>Elaphoglossum conforme</i>	417
			248	" <i>latifolium</i> ...	418

INDEX TO THE PLATES. .

481

PLATE	PAGE	PLATE	PAGE
249 <i>Elaphoglossum</i>		271 <i>Platynerium grande</i> ...	444
<i>stigmatolepis</i>	419	272 <i>Platynerium Wallichii</i> ...	446
250 " <i>viscosum</i> ...	419	273 " <i>biforme</i> ...	447
251 " <i>squamosum</i>	419	274 <i>Osmunda javanica</i> ...	448
252 " <i>spathulatum</i>	419	275 " <i>Claytoniana</i> ...	449
253 <i>Stenochlaena palustre</i> ...	422	276 " <i>regalis</i> ...	449
254 " <i>sorbifolia</i> ...	423	277 <i>Schizaea malaccana</i> ...	450
255 <i>Polybotrya appendiculata</i> ,		278 " <i>dichotoma</i> ...	451
var. β <i>aspleniifolia</i>	425	279 " <i>digitata</i> ...	452
256 " <i>appendiculata</i> ,		280 <i>Anemia tomentosa</i> ...	454
var. δ <i>Hamiltoniana</i>	426	281 <i>Lygodium circinatum</i> ...	455
257 " <i>appendiculata</i> ,		282 " <i>microphyllum</i>	456
var. α <i>costulata</i> ...	426	283 " <i>flexuosum</i> ...	457
258 <i>Gymnopteris variabilis</i> ...	428	284 " <i>polystachyum</i>	457
259 " <i>minus</i> ...	430	285 <i>Angiopteris evecta</i> ...	459
260 " <i>metallica</i> ...	431	286 <i>Marattia fraxinea</i> ...	461
261 " <i>spicata</i> ...	431	287 <i>Kaulfussia aesculifolia</i> ...	463
262 " <i>quercifolia</i>	433	288 <i>Ophioglossum nudicaule</i> ...	464
263 " <i>tricuspis</i> ...	435	289 " <i>fibrosum</i> ...	465
264 " <i>contaminans</i>	436	290 " <i>reticulatum</i>	466
265 " <i>subcrenata</i>	437	291 " <i>pendulum</i>	467
266 " <i>costata</i> ...	438	292 <i>Helminthostachys zeylanica</i>	468
267 " <i>Presliana</i> ...	430	293 <i>Botrychium Lunaria</i> ...	469
268 <i>Acrostichum aureum</i> ...	441	294 " <i>daucifolium</i>	470
269 <i>Photinopteris rigida</i> ...	443	295 " <i>virginianum</i> ,	
270 " <i>drynarioides</i>	444	var. <i>lanuginosum</i>	471



INDEX TO THE SPECIES AND SYNONYMS, THE LATTER BEING IN ITALICS.

	PAGE		PAGE
Aconiopteris		<i>Edgeworthii</i> ...	84
gorgonea ...	427	<i>emarginatum</i> ...	86
<i>Acrophorus.</i> see <i>Leucostegia</i>		flabellulatum ...	88
<i>Acropteris</i> ...	145	hispidulum ...	86
Acrostichum		<i>Lanceum</i> ...	75
<i>alcicorne</i> ...	445	lunulatum ...	82
aureum ...	440	var. <i>Mettenii</i> ...	83
<i>contaminans</i> ...	435	<i>orbiculatum</i> ...	75
<i>costatum</i> ...	438	Parishii ...	82
<i>deltigerum</i> ...	438	pedatum ...	86
<i>dichotomum</i> ...	197	<i>pteropus</i> ...	83
<i>fuciforme</i> ...	447	<i>rhizophorum</i> ...	84
<i>quercifolium</i> ...	433	<i>soboliferum</i> ...	84
<i>septentrionale</i> ...	145	venustum ...	86
<i>sorbifolium</i> ...	443	Allantodia	
<i>subcrenatum</i> ...	437	<i>Brunoniana</i> ...	197
<i>terminans</i> ...	437	<i>javanica</i> ...	195
<i>undulatum</i> ...	438	<i>Allosorus.</i>	
<i>virens</i> ...	439	<i>crispus</i> ...	98
<i>Wightianum</i> ...	426	<i>gracilis</i> ...	100
Actiniopteris		Alsophila	
dichotoma ...	197	albo-setacea ...	16
<i>radiata</i> ...	197	<i>alternans</i> ...	10
Adiantum		Andersoni ...	12
<i>Æthiopicum</i> ...	84	<i>Brunoniana</i> ...	10
capillus-veneris ...	84	commutata ...	14
caudatum ...	83	comosa ...	13
var. <i>Edgeworthii</i> ...	84	<i>contaminans</i> ...	12
var. <i>soboliferum</i> ...	84	crinita ...	16

	PAGE		PAGE
glabra	14	<i>amabile</i>	229
glauca	12	<i>appendiculatum</i>	239
Kingi	addenda	<i>aristatum</i>	229
latebrosa	11	<i>var. affine</i>	230
<i>var. hemitelioides</i>	10	<i>var. assamica</i>	230
Oldhami	13	<i>var. Hamiltoniana</i>	230
ornata	12	<i>atratum</i>	252
<i>Schmidiana.</i> see latebrosa.		<i>Brunonianum</i>	246
<i>Scottiana.</i> see Oldhami.		<i>calcaratum</i>	235
<i>squamulata</i>	14	<i>canariense</i>	250
mphecosmia		<i>canum</i>	235
alternans	10	<i>cicutarium</i>	220
Brunoniana	10	<i>ciliatum</i>	235
decipiens	10	<i>coadunatum</i>	222
Walkeræ	9	<i>coniifolium</i>	230
nemia		<i>crassifolium</i>	238
tomentosa	453	<i>cucullatum</i>	272
<i>Wightiana</i>	453	<i>cuspidatum</i>	232
ngiopteris		<i>decurrens</i>	219
evecta	460	<i>var. minor</i>	221
nisogonium		<i>densum</i>	255
cordifolium	191	<i>eberneum</i>	239, 252
esculentum	192	<i>exaltatum</i>	284
heterophlebium	191	<i>fuscipes</i>	227
lineolatum	191	<i>glandulosum</i>	273
Smithianum	192	<i>gracilescens</i>	234
lnogramma	383	<i>Hænkianum</i>	282
ntrophyum		<i>heterocarpum</i>	219
<i>coriaceum</i>	401	<i>hirsutulum</i>	237
latifolium	404	<i>hirtipes</i>	232
<i>parvulum</i>	403	<i>immersum</i>	235
plantagineum	403	<i>lineatum</i>	274
reticulatum	401	<i>macrocarpum</i>	165
<i>var. β parvulum</i>	401	<i>marginatum</i>	252
<i>semicordatum</i>	401	<i>membranaceum</i>	225
Arthobotrys		<i>multicaudatum</i>	222
<i>macrocarpa</i>	251	<i>multidentatum</i>	50
Aspidium		<i>multijugum</i>	279
<i>alatum</i>	214	<i>multilineatum</i>	277, 280

		PAGE			PAGE
<i>obliquum</i>	204	<i>bullatum</i>	159
<i>obscurum</i>	273	<i>caudatum</i>	151
<i>ochthodes</i>	240	<i>Ceterach</i>	194
<i>oppositum</i>	255	<i>contiguum</i>	151
<i>pennigerum</i>	277	<i>crinicaule</i>	150
<i>filosulum</i>	258	<i>decurrens</i>	152
<i>polycarpum</i>	201	<i>drepanophyllum</i>	164
<i>polymorphum</i>	218	<i>eberneum</i>	171
<i>prionophyllum</i>	280	<i>elongatum</i>	147
<i>pteropus</i>	219	<i>ensiforme</i>	141
<i>puberulum</i>	277	<i>erectum</i>	148
<i>rhomboideum</i>	229	<i>exiguum</i>	158
<i>scabrosum</i>	264	<i>falcatum</i>	150
<i>semibipinnatum</i>	215	<i>fontanum</i>	158
<i>Simonsii</i>	218	<i>var. exiguum</i>	158
<i>singaporianum</i>	212	<i>formosum</i>	152
<i>solutum</i>	278	<i>furcatum</i>	157
<i>speciosum</i>	230	<i>Gardneri</i>	151
<i>subconfluens</i>	214	<i>Griffithianum</i>	142
<i>subdiaphanum</i>	258	<i>heterocarpum</i>	153
<i>subtriphyllum</i>	215	<i>hirtum</i>	149
<i>syrmaticum</i>	243	<i>Hookerianum</i>	195
<i>tectum</i>	278	<i>javanicum</i>	195
<i>variolosum</i>	216	<i>laciniatum</i>	154
<i>vastum</i>	212	<i>var. crinigerum</i>	154
<i>venulosum</i>	273	<i>„ depauperatum</i>	154
<i>Wallichii</i>	288	<i>lanceolatum</i>	158
<i>Zollingerianum</i>	218	<i>laserpitiifolium</i>	158
Asplenium			<i>lobulosum</i>	179
<i>acuminatum</i>	178	<i>longifolium</i>	179
<i>adiantum-nigrum</i>	156	<i>longissimum</i>	145
<i>affine</i>	157	<i>lunulatum</i>	147
<i>alternans</i>	142	<i>var. camptorachis</i>	148
<i>ariifolium</i>	413	<i>„ trapeziforme</i>	148
<i>auritum</i>	149	<i>macrophyllum</i>	150
<i>Beddomei</i>	150	<i>marginatum</i>	185
<i>Belangeri</i>	161	<i>monanthemum</i>	144
<i>bulbiferum</i>	159	<i>multijugum</i>	144
			<i>nitidum</i>	157

	PAGE		PAGE
normale	144	Atkinsonii	163
var. β	144	var. <i>Andersonii</i>	163, 174
paradoxum	151	<i>australe</i>	189
pekinense	156	brevisorum	171
<i>persicæfolium</i>	149	<i>Clarkei</i>	166
<i>pinnatifido-pinnatum</i>	178	<i>drepanophyllum</i>	164
<i>planicaule</i>	154	<i>erythrorachis</i>	168
<i>præmorsum</i> . see <i>furcatum</i>		falcatum	164
pumilum	154	Filix-foemina	168
var. <i>hymenophylloides</i>	156	var. <i>attenuata</i>	169
<i>resectum</i>	152	„ <i>dentigera</i>	169
Ruta-muraria	156	„ <i>flabellulata</i>	170
rutæfolium	159	„ <i>parasnathensis</i>	170
septentrionale	145	„ <i>pectinata</i>	169
<i>sikkimense</i>	184	„ <i>polyspora</i>	170
<i>spathulinum</i>	157	„ <i>retusa</i>	170
<i>spinulosum</i>	166	fimbriatum	172
<i>squarrosus</i>	165	var. <i>foliosa</i>	172
<i>stenophyllum</i>	141	„ <i>sphæropteroides</i>	174
subavenium	144	<i>Griffithii</i>	
tenerum	147	<i>gymnogrammoides</i>	168
tenuifolium	159	var. <i>erythrorachis</i>	168
<i>trapeziforme</i>	148	Hohenackerianum	163
Trichomanes	143	macrocarpum	165
unilaterale	152	var. <i>Atkinsoni</i>	165
var. <i>delicatulum</i>	153	„ <i>1-pinnata</i>	165
„ <i>rivale</i>	153	<i>multicaudatum</i>	190
„ <i>udum</i>	153	<i>nigripes</i>	166
varians	158	var. <i>Clarkei</i>	166
viride	143	„ <i>dissecta</i>	166
<i>vulcanicum</i>	146	„ <i>selenopteris</i>	166
Wightianum	146	<i>oxyphyllum</i>	170
„ var. <i>microphyllum</i>	146	var. <i>Kulhaitense</i>	171
<i>zamiioides</i>	151	<i>pectinatum</i>	168, 169
Zenkerianum	148	<i>selenopteris</i>	166
		<i>spinulosum</i>	161
Athyrium		<i>stramineum</i>	171
<i>allantodioides</i>	164	<i>subtriangulare</i>	163
<i>aspidioides</i>	166	<i>thelypteroides</i>	164

	PAGE		PAGE
Blechnidium		fragrans	88
melanopus	133	laxa	92
Blechnum		mysorensis	89
cartilagineum	130	<i>nitidula</i>	102
Findlaysonianum	132	rufa	94
<i>melanopus</i>	133	subvillosa	93
<i>nitidum</i>	130	Szovitzii	89
orientale	132	tenuifolia	92
serrulatum	132	<i>Thwaitesii</i>	92
Botrychium daucifolium	462	varians	91
<i>lanuginosum</i>	471	<i>Chnoophora</i>	
Lunaria	469	<i>glauca</i>	12
<i>subcarnosum</i>	471	<i>Chrysodium</i>	
virginianum	471	<i>aureum</i>	440
var <i>lanuginosum</i>	471	Cibotium	
Brainea insignis	395	Barometz	24
<i>Callipteris</i> . see <i>Anisogonium</i>		<i>glaucum</i>	24
<i>esculentum</i>	192	<i>Cælopteris</i>	313
Campteria		<i>Crepidomanes</i>	
<i>anamallayensis</i>	118	<i>nanum</i>	40
biaurita	116	Cryptogramme	
<i>geminata</i>	118	Brunoniana	98
Kleiniana	116	<i>crispa</i>	98
<i>patens</i>	114	<i>Cryptosorus</i>	313
Wallichiana	118	Cyathea	
Ceratopteris		Brunonis	5
thalictroides	123	Hookeri	6
<i>Ceterach</i>		sinuata	5
<i>officinarum</i>	194	spinulosa	6
Cheilanthes albo-marginata...	94	Cyrtomium	
argentea	95	caducum	211
„ var. <i>chrysophylla</i>	93	<i>caryotideum</i>	211
<i>bullosa</i>	93	falcatum	211
<i>Dalhousiæ</i>	93	var. <i>caryotideum</i>	211
farinosa	92	Cystopteris fragilis	70
var. <i>chrysophylla</i>	93	<i>retusa</i>	170
„ <i>Dalhousiæ</i>	93	setosa	70
„ <i>flaccida</i>	93	Davallia <i>achilleifolia</i>	423
fragilis	91	<i>affinis</i>	54

	PAGE		PAGE
<i>bullata</i> ...	61	<i>Dictyopteris</i>	
<i>calvescens</i> ...	64	<i>Barberi</i> ...	298
<i>chinensis</i> ...	70	<i>chattagrammica</i> ...	299
<i>Clarkei</i> ...	52	<i>difformis</i> ...	300
<i>divaricata</i> ...	60	<i>heterosora</i> ...	302
<i>elegans</i> ...	59	<i>irregularis</i> ...	300
<i>epiphylla</i> ...	60	<i>polycarpa</i> ...	350
<i>Griffithiana</i> ...	60	<i>tenerifrons</i> ...	298
<i>hymenophylloides</i> ...	54	<i>Didymochlæna</i>	
<i>Lorrainei</i> ...	61	<i>lunulata</i> ...	199
<i>ornata</i> ...	59	<i>polycarpa</i> ...	201
<i>parvula</i> ...	54	<i>Diplazium</i>	
<i>pilosula</i> ...	68	<i>asperum</i> ...	184
<i>polyantha</i> ...	60	<i>assimile</i> ...	190
<i>puberula</i> ...	68	<i>australe</i> ...	189
<i>pulchra</i> ...	52	<i>bantamense</i> ...	177
<i>pyramidata</i> ...	68	<i>bellum</i> ...	189
<i>Roxburghii</i> ...	68	<i>decurrens</i> ...	186
<i>schizophylla</i> ...	75	<i>decussatum</i> ...	180
<i>solida</i> ...	59	<i>dilatatum</i> ...	187
<i>speciosa</i> ...	61	<i>var. β minor</i> ...	186
<i>tenuifolia</i> ...	70	<i>fraxineum</i> ...	386
<i>trichomanoides</i> ...	75	<i>fraxinifolium</i> ...	177
<i>triphylla</i> ...	58	<i>Griffithii</i> ...	190
<i>Dennstædtia</i>		<i>japonicum</i> ...	180
<i>appendiculata</i> ...	26	<i>Jerdoni</i> ...	190
<i>deltoidea</i> ...	26	<i>lanceum</i> ...	174
<i>Elwesii</i> ...	26	<i>lasiopteris</i> ...	180
<i>scabra</i> ...	24	<i>latifolium</i> ...	187
<i>Diacalpe</i>		<i>Lobbianum</i> ...	187
<i>aspidioides</i> ...	18	<i>lobulosum</i> ...	179
<i>foeniculacea</i> ...	18	<i>longifolium</i> ...	179
<i>Dicksonia</i>		<i>maximum</i> ...	187
<i>scabra</i> ...	24	<i>multicaudatum</i> ...	189
<i>Diclisodon</i>		<i>pallidum</i> ...	175
<i>deparioides</i> ...	257	<i>pinnatifido-pinnatum</i> ...	178
<i>Dictyocline</i>		<i>polypodioides</i> ...	184
<i>Griffithii</i> ...	415	<i>var. decurrens</i> ...	186
		<i>var. β (Thw.)</i> ...	186

		PAGE			PAGE
<i>polyrhizon</i>	...	180	<i>mollis</i>	...	341
<i>porrectum</i>	...	176	<i>propinqua</i>	...	339
<i>Prescottianum</i>	...	178	<i>quercifolia</i>	...	341
<i>procerum</i>	...	189	<i>rigidula</i>	...	344
<i>Schkuhrii</i>	...	181	<i>splendens</i>	...	339
<i>sikkimense</i>	...	184	<i>Dryostachyum</i>	...	339
<i>sorzogonense</i>	...	181	<i>Elaphoglossum</i>		
<i>speciosum</i>	...	178	<i>conforme</i>	...	416
<i>Stoliczkæ</i>	...	182, 184	<i>latifolium</i>	...	416
<i>var. hirsutipes</i>	...	182	<i>laurifolium</i>	...	418
<i>subserratum</i>	...	174	<i>Norissii</i>	...	418
<i>sylvaticum</i>	...	177	<i>piloselloides</i>	...	421
<i>var. Prescottianum</i>		178	<i>spathulatum</i>	...	420
<i>thelypteroides</i>	...	184	<i>squamosum</i>	...	420
<i>Thwaitesii</i>	...	180	<i>stelligerum</i>	...	240
<i>tomentosum</i>	...	179	<i>stigmatolepis</i>	...	418
<i>travancoricum</i>	...	188	<i>viscosum</i>	...	420
<i>umbrosum</i>	...	180	<i>Galeoglossa</i>	...	394
<i>var. assimile</i>	...	190	<i>Gleichenia</i>		
<i>„ australe</i>	...	189	<i>circinata</i>	...	2
<i>„ multicaudatum</i>		190	<i>dichotoma</i>	...	4
<i>„ procerum</i>	...	189	<i>gigantea</i>	...	2
<i>Zeylanicum</i>	...	175	<i>glauca</i>	...	2
Dipteris			<i>linearis</i>	...	4
<i>Horsfieldii</i>	...	336	<i>longissima</i>	...	2
<i>Lobbiana</i>	...	336	<i>Norrisii</i>	...	2
<i>Wallichiana</i>	...	334	<i>semivestita</i>	...	2
Doodia			<i>Goniophlebium amænum</i>	...	317
<i>dives</i>	...	137	<i>argutum</i>	...	323
Drymoglossum			<i>erythrocarpum</i>	...	319
<i>Beddomei</i>	...	413	<i>Hendersoni</i>	...	320
<i>carnosum</i>	...	411	<i>lachnopous</i>	...	319
<i>piloselloides</i>	...	411	<i>microrhizoma</i>	...	322
<i>var. β Beddomei</i>	...	413	<i>molle</i>	...	322
Drynaria			<i>subamænum</i>	...	317
<i>conjugata</i>	...	339	<i>subauriculatum</i>	...	322
<i>coronans</i>	...	338	<i>verrucosum</i>	...	324
<i>diversifolia</i>	...	344	Goniopteris		
<i>Linnæi</i>	...	343	<i>lineata</i>	...	275, 276

INDEX TO THE SPECIES AND SYNONYMS.

489

	PAGE		PAGE
<i>multilineata</i> ...	275	Wallii ...	431
<i>penangiana</i> ...	276	<i>Haplopteris</i> ...	408
<i>prolifera</i> ...	296	<i>Helminthostachys</i>	
<i>urophylla</i> ...	274	<i>dulcis.</i> see <i>zeylanica</i>	
<i>Grammitis</i>		<i>zeylanica</i> ...	467
<i>aurita</i> ...	377	<i>Hemidictyum</i>	
<i>Hamiltoniana</i> ...	373	<i>Ceterach</i> ...	194
<i>lanceolata</i> ...	393	<i>Finlaysonianum</i> ...	195
<i>Totta</i> ...	377	<i>Hemionitis</i>	
<i>vestita</i> ...	387	<i>arifolia</i> ...	413
<i>Gymnogramme</i>		<i>cordifolia</i> ...	415
<i>alismæfolia</i> ...	389	<i>Griffithii</i> ...	415
<i>Andersoni</i> ...	382	<i>hastata.</i> see <i>arifolia</i>	
<i>Hamiltoniana</i> ...	390	<i>Hemitelea</i>	
<i>javanica</i> ...	386	<i>decipiens</i> ...	10
<i>leptophylla</i> ...	382	<i>Beddomei.</i> see <i>Cyathea spinulosa</i>	
<i>microphylla</i> ...	384	<i>Humata</i>	
<i>obtusata</i> ...	380	<i>alpina</i> ...	48
<i>serrulata</i> ...	386	<i>angustata</i> ...	47
<i>vestita</i> ...	387	<i>heterophylla</i> ...	46
<i>Wallichii</i> ...	387	<i>parallela</i> ...	47
<i>Gymnopteris</i>		<i>pedata</i> ...	48
<i>axillaris</i> ...	430	<i>vestita</i> ...	48
<i>contaminans</i> ...	438	<i>Hymenolepis</i>	
<i>decurrens.</i> see <i>variabilis</i>		<i>spicata</i> ...	432
<i>flagellifera</i> ...	483	<i>Hymenophyllum</i>	
<i>Feei.</i> see <i>variabilis</i>		<i>badium</i> ...	33
<i>lanceolata</i> ...	429	<i>Blumeianum</i> ...	32
<i>metallica</i> ...	432	<i>ciliatum</i> ...	33
<i>minus</i> ...	430	<i>crispatum</i> ...	33
<i>Presliana</i> ...	439	<i>denticulatum</i> ...	34
<i>quercifolia</i> ...	432	var. <i>flaccidum</i> ...	35
<i>spicata</i> ...	432	<i>emersum</i> ...	30
<i>subcrenata</i> ...	437	<i>exiguum</i> ...	37
<i>subrepanda</i> ...	434	<i>exsertum</i> ...	30
<i>tricuspis</i> ...	434	<i>flaccidum</i> ...	35
<i>variabilis</i> ...	429	<i>javanicum</i> ...	32
var. β <i>lanceolata</i> ...	429	var. <i>badium</i> ...	33
,, γ <i>axillaris</i> ...	430	<i>khasianum</i> ...	35

	PAGE		PAGE
Levingii ...	36	<i>concinna</i> ...	258
<i>microglossum</i> ...	32	<i>coniifolia</i> ...	238
Neesii ...	35	<i>crassifolia</i> ...	238
parvifolium ...	28	<i>crenata</i> ...	258
polyanthos ...	30	<i>cuspidata</i> ...	232
var. <i>Blumeum</i> ...	32	<i>deltoidea</i> ...	254
" <i>microglossum</i> ...	32	<i>deparioides</i> ...	257
" <i>minor</i> ...	32	<i>dissecta</i> ...	260
Simonsianum ...	34	var. <i>ingens</i> ...	261
Smithii ...	34	<i>divisa</i> ...	264
tenellum ...	30	<i>elongata</i> . see <i>Filix-mas</i> .	
<i>Tunbridgense</i> ...	35	<i>Elwesii</i> ...	239
<i>Hypolepis</i>		<i>eriocarpa</i> ...	258
<i>hostilis</i> ...	295	<i>Fairbankii</i> ...	240
<i>Kaulfussia</i>		<i>falciloba</i> ...	235
<i>æsculifolia</i> ...	462	<i>Falconeri</i> ...	248
<i>assamica</i> ...	462	<i>ferruginea</i> ...	264
<i>Lastrea</i>		<i>Filix-mas</i> ...	248
<i>affinis</i> ...	230	var. <i>apiciflora</i> ...	249
<i>alsophiloides</i> ...	8	" <i>cochleata</i> ...	250
<i>amabilis</i> ...	229	" <i>elongata</i> ...	250
<i>angustifrons</i> ...	262	" <i>fibrillosa</i> ...	250
<i>apiciflora</i> ...	249	" <i>marginata</i> ...	250
<i>aristata</i> ...	229	" <i>odontoloma</i> ...	248
var. <i>Hamiltonii</i> ...	230	" <i>panda</i> ...	251
<i>barbigera</i> ...	246	" <i>paralellogramma</i> ...	249
<i>Beddomei</i> ...	239	" <i>Schimperiana</i> ...	250
<i>Bergiana</i> ...	239	<i>flaccida</i> ...	244
<i>Blumei</i> ...	259	<i>fœniculacea</i> ...	18
<i>Boryana</i> ...	266	<i>fuscipes</i> ...	243
<i>Brunoniana</i> ...	246	<i>gracilescens</i> ...	239
<i>Buchanani</i> ...	255	<i>gracilescens</i> ...	234
<i>calcarata</i> ...	235	var. <i>decipiens</i> ...	234
var. <i>ciliata</i> ...	235	" <i>hirsutipes</i> ...	234
" <i>falciloba</i> ...	237	<i>gracilis</i> ...	207
" <i>sericea</i> ...	237	<i>grisea</i> ...	241
<i>cana</i> ...	238	<i>Hendersoni</i> ...	257
<i>Clarkei</i> ...	250	<i>hirtipes</i> ...	232
<i>cochleata</i> . see <i>Filix-mas</i> .		<i>immersa</i> ...	234

	PAGE
<i>ingens</i>	261
<i>intermedia</i>	250
<i>intermedia</i> (Baker)	260
<i>melanopus</i>	243
<i>membraniifolia</i>	261
<i>microstegia</i>	292
<i>nephrodioides</i>	238
<i>Nidus</i>	250
<i>obtusiloba</i>	264
<i>octhodes</i>	240
„ β <i>tylodes</i>	240
<i>odontoloma</i>	248
<i>Parishii</i>	259
<i>patentissima</i>	249
<i>platypus</i>	230
<i>pulvinulifera</i>	255
<i>recedens</i>	260
<i>rhodolepis</i>	260
<i>rigida</i>	251
<i>sagenioides</i>	243
<i>scabrosa</i>	263
<i>sericea</i>	235
<i>setigera</i>	267
<i>setosa</i> . see <i>Cystopteris</i>	
<i>sikkimensis</i>	259
<i>sparsa</i>	252
var. <i>deltoidea</i>	254
„ <i>gracilis</i>	207
„ <i>latesquama</i>	255
„ <i>minor</i>	254
„ <i>nitidula</i>	254
„ <i>obtussissima</i>	254
„ <i>squamulosa</i>	255
„ <i>undulata</i>	254
„ <i>zeylanica</i>	254
<i>spectabilis</i>	244
<i>spectabilis</i>	257
<i>spinulosa</i>	252
var. <i>remota</i>	252

	PAGE
<i>splendens</i>	262
<i>syrmatica</i>	243
<i>tenericaulis</i>	266
<i>Thelypteris</i>	241
<i>Thwaitesii</i>	258
<i>tylodes</i>	240
<i>undulata</i>	254
<i>viscosa</i>	238
<i>Walkeræ</i>	233
„ δ <i>bipinnata</i>	234
„ β <i>macrocarpa</i>	233
„ γ <i>pinnatifida</i>	233
<i>Leptochilus</i> . see <i>Gymnopteris</i> .	
<i>Leptogramme</i>	
„ <i>aurita</i>	377
var. <i>Levingii</i>	377
„ <i>obtusata</i>	380
„ <i>opaca</i>	379
„ <i>Totta</i>	377
<i>Leucostegia</i>	
„ <i>affinis</i>	54
„ <i>assamica</i>	51
„ <i>dareæformis</i>	52
„ <i>Hookeri</i>	52
„ <i>hymenophylla</i>	48
„ <i>hymenophylloides</i>	54
„ <i>immersa</i>	51
„ <i>membranulosa</i>	50
„ <i>multidentata</i>	30
„ <i>nodosa</i>	54
„ <i>parvula</i>	54
„ <i>pseudo-cystopteris</i>	54
„ <i>pulchra</i>	52
<i>Lindsaya</i>	
„ <i>caudata</i>	75
„ <i>cultrata</i>	72
„ <i>divergens</i>	76
„ <i>flabellulata</i>	75
„ <i>heterophylla</i>	77

			PAGE				PAGE
Lancea	75	<i>longifolium</i>	457
lanuginosa	77	microphyllum	455
<i>Lobbiana</i>	72	<i>pinnatifidum</i>	457
<i>nitens</i>	79	<i>pedatum</i>	455
orbiculata	75	polystachyum	457
var. tenera	75	<i>scandens</i>	455
<i>pectinata</i>	74	Marattia fraxinea	460
repens	74	Matonia pectinata	19
var. minor	74	Meniscium			
rigida	75	cuspidatum	400
scandens	74	<i>deltigerum</i>	438
<i>trapeziformis</i>	75	<i>longifrons</i>	400
Walkeræ	76	<i>Parishii</i>	399
Litobrochia				salicifolium	399
<i>aurita</i>	122	Thwaitesii	399
incisa	120	triphyllum	397
<i>ludens</i>	120	var. Parishii	399
marginata	122	<i>Mertensia.</i>	see	Gleichenia	
<i>pedata</i>	120	Mesochloena			
<i>tripartita</i>	122	polycarpa	199
Lomaria				Microlepia			
adnata	127	<i>flaccida</i>	68
<i>elongata</i>	125	<i>hirta</i>	68
<i>euphlebia</i>	129	Hookeriana	62
<i>glauca</i>	127	Kurzii	66
<i>limoniifolia</i>	423	majuscula	66
Patersoni	125	marginalis	64
<i>pycnophylla</i>	129	var. calvescens	64
scandens	423	pinnata	64
<i>Lomariopsis</i>	423	platyphylla	66
Loxogramme				<i>polypodioides</i>	68
avenia	393	<i>proxima</i>	67
involuta	393	<i>pteropus</i>	50
lanceolata	392	<i>rhomboidea</i>	68
Lygodium				<i>scabra</i>	64
circinnatum	455	speluncæ	67
<i>dichotomum</i>	455	var. hirta	68
flexuosum	457	strigosa	67
japonicum	457	urophylla	64

INDEX TO THE SPECIES AND SYNONYMS. 493

	PAGE		PAGE
Monogramme		otaria	267
<i>Junghuhnii</i>	375	<i>parasiticum</i>	278
<i>paradoxa</i>	375	<i>pennigerum</i>	277
Nephrodium		<i>var. multilineatum</i>	277
<i>abortivum</i>	277	<i>procurrens</i>	278
<i>abruptum</i>	277	<i>prolixum</i>	240
<i>amboinense</i>	278	<i>propinquum</i>	269
<i>Arbuscula</i>	276	<i>pteroides</i>	269
<i>aridum</i>	272	<i>punctatum</i>	270
<i>artinezum</i>	227	<i>simulans</i>	223
<i>brachyodon</i>	280	<i>singaporianum</i>	212
<i>costatum</i>	275	<i>terminans</i>	269
<i>crinipes</i>	279	<i>truncatum</i>	280
<i>cucullatum</i>	270	<i>unitum</i>	268, 272
<i>cuspidatum</i>	232	<i>urophyllum</i>	274
<i>didymosorum</i>	278	<i>variolosum</i>	218
<i>elatum</i>	272	<i>Wightii</i>	218
<i>eusorum</i>	280	<i>Zollingerianum</i>	218
<i>extensum</i>	269	Nephrolepis	
<i>var. microsorum</i>	270	<i>acuta</i>	284
" <i>minor</i>	278	<i>biserrata</i>	284
<i>falcilobum</i>		<i>cordifolia</i>	282
<i>var. pubera</i>	277	<i>exaltata</i>	282
<i>ferox</i>	279	<i>obliterata</i>	285
<i>glandulosum</i>	273	<i>ramosa</i>	284
<i>var. late-strigosa</i>	277	<i>tuberosa</i>	282
<i>Griffithii</i>	222	<i>volubilis</i>	284
<i>immersum</i>	235	Niphobolus	
<i>ingens.</i> see <i>Lastrea dissecta</i>		<i>acrostichoides</i>	327
<i>javanicum</i>	201	<i>adnascens</i>	325
<i>lineatum</i>	274	<i>angustatus</i>	353
<i>microsorium</i>	270	<i>Boothii</i>	333
<i>molle</i>	277	<i>costatus</i>	329
<i>var. amboinense</i>	278	<i>detergibilis</i>	331
" <i>aureum</i>	279	<i>fissus</i>	330
" <i>didymosorum</i>	279	<i>floccigerus</i>	331
" <i>multijugum</i>	279	<i>flocculosus</i>	331
" <i>procurrens</i>	278	<i>Gardneri</i>	331
<i>moulmeinense</i>	275	<i>Heteractis</i>	327

		PAGE			PAGE
<i>jaintensis</i>	...	327	<i>vulgatum</i>	...	464
<i>lævis</i>	...	325	<i>Wightii</i>	...	465
<i>Lingua</i>	...	327, 328	<i>Ornithopteris</i>	...	115
<i>nummulariæfolius</i>	...	324	<i>Osmunda</i>		
<i>pannosus</i>	...	328	<i>Claytoniana</i>	...	449
<i>penangianus</i>	...	332	<i>javanica</i>	...	447
<i>porosus</i>	...	330	<i>Leschenaultii.</i>	see <i>regalis</i>	
<i>Schmidianus</i>	...	331	<i>regalis</i>	...	450
<i>sphærocephalus</i>	...	353	<i>virginiana</i>	...	471
<i>stigmatosus</i>	...	328	<i>zeylanica</i>	...	467
<i>subfurfuraceus</i>	...	329	<i>Patania</i>		
<i>Niphopsis</i>	...	353	<i>appendiculata</i>	...	26
<i>Nothochlæna</i>	...	375	<i>Elwesii</i>	...	26
<i>Notholaena</i>			<i>Pellaea</i>		
<i>lanuginosa</i>	...	375	<i>Boivini</i>	...	102
<i>Marantæ</i>	...	373	<i>calomelanos</i>	...	104
<i>vellea</i>	...	375	<i>concolor</i>	...	100
<i>Odontoloma</i>			<i>falcata</i>	...	102
<i>repens</i>	...	74	<i>geraniifolia</i>	...	101
<i>Oleandra</i>			<i>gracilis</i>	...	100
<i>Cumingii</i>	...	288	<i>nitidula</i>	...	101
<i>musæfolia</i>	...	287	<i>Stelleri</i>	...	100
<i>neriiformis</i>	...	285	<i>Tamburii</i>	...	101
<i>Wallichii</i>	...	287	<i>Peranema</i>		
<i>Onoclea</i>			<i>cyatheoides</i>	...	22
<i>orientalis</i>	...	20	<i>Phegopteris</i>		
<i>Onychium</i>			<i>auriculata</i>	...	290
<i>auratum</i>	...	96	<i>davallioides</i>	...	296
<i>japonicum</i>	...	96	<i>distans</i>	...	292
var. <i>intermedia</i>	...	96	var. <i>glabrata</i>	...	292
„ <i>multidentata</i>	...	96	„ <i>minor</i>	...	292
<i>Ophioglossum</i>			<i>Dryopteris</i>	...	293
<i>brevipes</i>	...	465	<i>erubescens</i>	...	289
<i>cordifolium</i>	...	465	<i>ornata</i>	...	294
<i>fibrosum</i>	...	465	<i>pallida</i>	...	295
<i>nudicaule</i>	...	465	<i>polycarpa</i>	...	300
<i>parvifolium</i>	...	464	<i>punctata</i>	...	295
<i>pendulum</i>	...	465	<i>recedens</i>	...	260
<i>reticulatum</i>	...	464	<i>Robertiana</i>	...	293

	PAGE		PAGE
Phegopteris		ebenipes	363
<i>rufescens</i>	293	<i>var. Oakesii</i>	364
<i>rugulosa</i>	295	„ <i>Parishii</i>	364
<i>Scottii</i>	289	<i>Griffithiana</i>	354
<i>subdigitata</i>	295	<i>hastata</i>	362
<i>vulgaris</i>	290	<i>hemionitidea</i>	358
Photinopteris		<i>heterocarpa</i>	357
<i>drynarioides</i>	442	<i>himalayensis</i>	370
<i>rigida</i>	442	<i>incurvata</i>	364
<i>Phymatodes.</i> see <i>Pleopeltis.</i>		<i>insignis</i>	365
Plagiogyria		<i>irioides</i>	357
<i>adnata</i>	127	<i>juglandifolia</i>	368
<i>euphlebia</i>	129	<i>var. tenuicauda</i>	370
<i>glauca</i>	127	<i>lanceolata</i>	351
<i>pycnophylla</i>	129	<i>Lehmanni</i>	370
<i>triquetra.</i> see <i>euphlebia</i>		<i>leiorhiza</i>	372
Platycerium		<i>lepidota</i>	351
<i>biforme</i>	445	<i>linearis</i>	346
<i>grande</i>	445	<i>var. steniste</i>	347
<i>Wallichii</i>	445	<i>longifolia</i>	349
Platyloma		<i>longissima</i>	366
<i>falcata</i>	104	<i>macrosora</i>	353
Pleocnemia		<i>malacodon</i>	363
<i>aristata</i>	268	<i>var. β majus</i>	363
<i>Clarkei</i>	227	<i>membranacea</i>	355
<i>gigantea</i>	224	<i>musæfolia</i>	358
<i>leuzeana</i>	228	<i>nigrescens</i>	367
<i>membranacea</i>	225	<i>normalis</i>	353
<i>membranifolia</i>	225	<i>nuda</i>	347
<i>Thwaitesii</i>	223	<i>ovata</i>	354
<i>Trimeni</i>	224	<i>palmata</i>	368
Pleopeltis		<i>Parishii</i>	341
<i>accedens</i>	345	<i>phymatodes</i>	366
<i>angustata</i>	351	<i>pteropus</i>	359
<i>var. depauperata</i>	351	<i>var. minor</i>	361
<i>capitellata</i>	370	„ <i>zosteræformis</i> ..	362
<i>clathrata</i>	348	<i>punctata</i>	357
<i>cryptolobum</i>	363	<i>rhyncophylla</i>	353
<i>dilatata</i>	367	<i>rostrata</i>	345

			PAGE				PAGE
	simplex	...	347		<i>conjugatum</i>	...	336
	sinuosa	...	349		<i>contiguum</i>	...	350
	stenophylla	...	348		<i>cordifolium</i>	...	282
	<i>Stewartii</i>	...	363		cornigerum	...	307
	superficialis	...	351		<i>costatum</i>	...	275
	<i>trifida</i>	...	362		<i>crenatum</i>	...	258
	<i>Wightiana</i>	...	347		cucullatum	...	307
	Zippellii	...	357		dareæforme	...	316
	<i>zosteræformis</i>	...	362		decorum	...	310
<i>Pæcilopteris</i>					<i>dentigerum</i>	...	169
	<i>Blumeana</i>	...	435		<i>Dipteris</i>	...	336
	<i>costata</i>	...	438		<i>distans.</i>	see <i>Phegopteris</i>	
	<i>flagellifera</i>	...	433		<i>Dryopteris</i>	...	293
	<i>Hookeriana</i>	...	437		<i>elongatum</i>	...	232
	<i>Presliana</i>	...	440		<i>ellipticum</i>	...	392
	<i>prolifera</i>	...	437		<i>erubescens</i>	...	289
	<i>repanda</i>	...	435		<i>exaltatum</i>	...	282
	<i>semicordata</i>	...	435		<i>excavatum</i>	...	347
<i>Polybotrya</i>					<i>Filix-mas</i>	...	249
	appendiculata	...	424		<i>fragrans</i>	...	89
	var. <i>aspleniifolia</i>	...	424		<i>furfuraceum</i>	...	327
	„ <i>costulata</i>	...	426		fuscatum	...	311
	„ <i>Hamiltoniana</i>	...	424		<i>glabrum</i>	...	358
	„ <i>major</i>	...	424		<i>gladiatum</i>	...	346
	„ <i>subintegra</i>	...	427		<i>glandulosum</i>	...	309
	<i>aspleniifolia</i>	...	426		<i>glaucum</i>	...	2
	<i>Helferiana</i>	...	426		<i>grandifolium</i>	...	355
<i>Polypodium</i>					<i>Grevilleanum</i>	...	347
	<i>acutissimum</i>	...	346		<i>Griffithii</i>	...	291
	<i>adnascens</i>	...	325		<i>harpophyllum</i>	...	204
	<i>adnatum</i>	...	292		<i>hastatum</i>	...	364
	<i>alternifolium</i>	...	366 & 367		<i>heterocarpum</i>	...	355
	<i>appendiculatum</i>	...	290		<i>hirtellum</i>	...	305
	<i>argutum</i>	...	323		<i>hirtum</i>	...	305
	<i>auriculatum</i>	...	203 & 290		<i>jaintense</i>	...	327
	<i>Beddomei</i>	...	322		<i>khasyanum</i>	...	308
	<i>bifurcatum</i>	...	338		<i>lanceolatum</i>	...	351
	<i>brunneum</i>	...	292		<i>lasiosorum</i>	...	305
	<i>coniifolium</i>	...	296		<i>lepidota</i>	...	351

	PAGE		PAGE
<i>lineare</i> (Linn.) ...	5	<i>Russellianum</i> ...	267
<i>lineare</i> var. β <i>simplex</i> ...	347	<i>semibipinnatum</i> ...	215
<i>Linnæi</i> ...	343	<i>serra</i> ...	310
<i>Lonchitis</i> ...	203	<i>sesquipedale</i> ...	347
<i>longifrons</i> ...	353	<i>sessilifolium</i> ...	307
<i>longipes</i> ...	292	<i>simplex</i> ...	347
<i>loriforme</i> ...	346	<i>speluncæ</i> ...	57
<i>marginale</i> ...	64	<i>sphærocephalum</i> ...	353
<i>mediale</i> ...	303	<i>subconfluens</i> ...	300
<i>minutum</i> ...	313	<i>subdigitatum</i> ...	296
<i>multilineatum</i> ...	275	<i>subevenosum</i> ...	303
<i>mysurense</i> ...	331	<i>subfalcatum</i> ...	314
<i>nigrocarpum</i> ...	264	<i>subtripinnatum</i> ...	266
<i>normale</i> ...	353	<i>superficiale</i> ...	351
<i>nummulariæfolium</i> ...	334	<i>tenericaule</i> ...	266
<i>obliquatum</i> ...	311	<i>Thwaitesii</i> ...	309
<i>obscurum</i> ...	243	<i>trichomanoides</i> ...	308
<i>ornatum</i> ...	294	<i>tridactylon</i> ...	361
<i>ovatum</i> ...	354	<i>unitum</i> ...	269
<i>oxylobum</i> ...	362	<i>venustum</i> ...	370
<i>baludosum</i> ...	292	<i>verrucosum</i> ...	324, 325
<i>palustre</i> ...	421	<i>vittarioides</i> ...	325
<i>parasiticum</i> ...	278	<i>Walkera</i> ...	233
<i>parasiticum</i> ...	302	<i>Wallii</i> ...	305
var. <i>pilosiusculum</i> ...	305	<i>Wightianum</i> ...	346
<i>parvulum</i> ...	314	<i>Zeylanicum</i> ...	303
<i>penangianum</i> ...	276	<i>Polystichum aculeatum</i> ...	207
<i>pertusum</i> ...	325	var. <i>anomalum</i> ...	209
<i>Phegopteris</i> ...	292	,, <i>angulare</i> ...	207
<i>polycephalum</i> ...	358	,, <i>biaristatum</i> ...	209
<i>porosum</i> ...	331	,, <i>lobatum</i> ...	207
<i>pteroides</i> ...	269	,, <i>mucronifolium</i> ..	210
<i>quercifolium</i> ...	343	,, <i>rufo-barbatum</i> ...	207
<i>repandulum</i> ...	313	,, <i>simifertile</i> ...	209
<i>rivale</i> ...	31	,, <i>setosum</i> ...	209
<i>Robertianum</i> ...	293	,, <i>travancorium</i> ...	209
<i>rufescens</i> ...	293	<i>amabile</i> . see <i>Lastrea</i>	
<i>rugosulum</i> ...	295	<i>aristatum</i> . see <i>Lastrea</i>	
<i>rugulosum</i> ...	295	var. <i>affine</i> . see <i>Lastrea</i>	

<i>brachypterum</i>
<i>conifolium</i>	see Lastre		
<i>caespitosum</i>
<i>faniculaceum</i>
<i>ilicifolium</i>
<i>lachenense</i>
<i>Lonchitis</i>
<i>obliquum</i>
<i>Prescottianum</i>
var. <i>Bakerianum</i>
" <i>castaneum</i>
<i>semicordatum</i>
<i>sikkimense</i>
<i>stimulans</i>
<i>Thomsoni</i>
var. <i>gracilis</i>
<i>Pasia</i>
<i>Prosaptia</i>			
<i>contigua</i>
<i>Emersoni</i>
<i>Pteris</i>			
<i>aquilina</i>	I
var. <i>esculenta</i>	I
<i>aspericaulis</i>	I
<i>liaurita</i>

		PAGE			PAGE
<i>coadunata</i>	...	222	<i>Sphæropteris</i>		
<i>gigantea</i>	224	<i>barbata</i>	22
<i>var. minor</i>	225	<i>Stegnogramme</i>		
<i>Griffithii</i>	222	<i>aspidiodes</i>	380
<i>heterocarpa</i>	219	<i>Stenochlæna</i>		
<i>heterosora</i>	219	<i>palustre</i>	421
<i>pteropus</i>	220	<i>scandens</i>	423
<i>var. minor</i>	220	<i>sorbifolia</i>	423
<i>siifolia</i>	218	<i>Stenoloma</i>		
<i>simulans</i>	223	<i>chinensis</i>	70
<i>Thwaitesii</i>	220	<i>Struthiopteris</i>		
<i>variolosa</i>	218	<i>orientalis</i>	20
<i>Schellolepis</i>	324	<i>Syngramme</i>		
<i>Schizæa</i>			<i>alismæfolia</i>	389
<i>dichotoma</i>	452	<i>fraxinea</i>	386
<i>digitata</i>	452	<i>vestita</i>	386
<i>malaccana</i>	452	<i>Wallichii</i>	387
<i>Schizoloma</i>			<i>Tæniopsis</i>		
<i>cordata</i>	79	<i>falcata</i>	410
<i>davallioides</i>	79	<i>lineata</i>	407
<i>ensifolia</i>	80	<i>scolopendrium</i>	407
<i>Griffithiana</i>	80	<i>Tænitis</i>		
<i>gueriniana</i>	80	<i>blechnoides</i>	410
<i>heterophylla</i>	80	<i>Teratophyllum</i>		
<i>lobata</i>	78	<i>aculeatum</i>	423
<i>var. malabaricum</i>	79	<i>Thamnopteris</i>		
<i>nitens</i>	79	<i>Grevillei</i>	139
<i>recurvata</i>	79	<i>musæfolia</i>	139
<i>Selliguea</i>			<i>Nidus</i>	137
<i>avenia</i>	394	<i>var. musæfolia</i>	139
<i>caudiformis</i>	390	,, <i>phyllitidis</i>	139
<i>decurrens</i>	392	<i>Simonsiana</i>	141
<i>elliptica</i>	392	<i>Trichomanes</i>		
<i>Féei</i>	389	<i>anceps</i>	44
<i>Hamiltoniana</i>	390	<i>auriculatum</i>	44
<i>involuta</i>	393	<i>bipunctatum</i>	41
<i>lanceolata</i>	393	<i>var. insigne</i>	42
<i>Maingayi</i>	392	,, <i>plicatum</i>	42
			<i>birmanicum</i>	43

Henzarianum
<i>insigne</i>
intramarginale
javanicum
Kurzii
<i>limbatum</i>
Motleyi
muscoides
var. <i>sublimbatum</i>		
<i>nanum</i>
neilgherrense
pallidum
parvulum
<i>plicatum</i>
proliferum
pusillum
pyxidiferum
var. <i>limbatum</i>



SELECTION
FROM THE
MISCELLANEOUS PUBLICATIONS
OF
THACKER, SPINK AND Co.
CALCUTTA.

LONDON : W. THACKER & Co., 87 NEWGATE STREET.



LAYS OF IND. By ALIPH CHEEM.

COMIC, SATIRICAL, AND DESCRIPTIVE

Poems Illustrative of Anglo-Indian Life.

Seventh Edition. Enlarged. Cloth, gilt. Rs. 7.

REVIEWS OF PREVIOUS EDITIONS.

"This is a remarkably bright little book. 'Aliph Cheem,' supposed to be the *nom de plume* of an officer in the 18th Hussars, is, after his fashion, an Indian Boz Gaultier. In a few of the poems the jokes, turning on local names and customs, are somewhat esoteric, but, taken throughout, the verses are characterized by high animal spirits, great cleverness, and most excellent fooling."—*The World*.

"Aliph Cheem presents us in this volume with some highly amusing ballads and songs, which have already in a former edition warmed the hearts and cheered the lonely hours of many an Anglo-Indian, the pictures being chiefly those of Indian life. There is no mistaking the humour, and at times, indeed, the fun is both 'fast and furious.' Many portions remind us of the 'Bab Ballads.' One can readily imagine the merriment created round the camp fire by the recitation of 'The Two Thumpers,' which is irresistibly droll. . . . The edition before us is enlarged, and contains illustrations by the author, in addition to which it is beautifully printed and handsomely got up, all which recommendations are sure to make the name of Aliph Cheem more popular in India than ever."—*Liverpool Mercury*.

THACKER, SPINK & CO., CALCUTTA.

Reviews of "Lays of Ind."

"The 'Lays' are not only Anglo-Indian in origin, but out-and-out Anglo-Indian in subject and colour. To one who knows something of life at an Indian 'station' they will be especially amusing. Their exuberant fun at the same time may well attract the attention of the ill-defined individual known as 'the general reader.'"—*Scotsman*.


"To many Anglo-Indians the lively verses of 'Alph Cheem' must be very well known, while to those who have not yet become acquainted with them we can only say, read them on the first opportunity. To those not familiar with Indian life they may be specially com-



mended for the picture which they give of many of its lighter incidents and conditions, and of several of its ordinary personages. . . . We have read the volume with real pleasure, and we have only to add that it is nicely printed and elegantly finished, and that it has several charming woodcuts, of which some are by the author, whom Indian gossip, by the way, has identified with Captain Yeldham, of the 18th Hussars."—*Bath Chronicle*.

"Satire of the most amusing and inoffensive kind, humour the most genuine, and pathos the most touching pervade these 'Lays of Ind.' . . . From Indian friends we have heard of the popularity these 'Lays' have obtained in the land where they were written, and we predict for them a popularity equally great at home."—*Monthly Homoeopathic Review*.

"Former editions of this entertaining book having been received with great favour by the public and by the press, a new edition has been issued in elegant type and binding. The author, although assuming a *nom de plume*, is recognized as a distinguished cavalry officer, possessed of a vivid imagination and a sense of humour amounting sometimes to rollicking and contagious fun. Many of his 'Lays' suggest recollections of some of the best pieces in the 'Ingoldsby Legends,' or in the 'Biglow Papers' of Russell Lowell, while revealing a character of their own. Anglo-Indian terms and usages are skilfully employed, and even what appears to some the uneventful life of a secluded station is made to yield incidents for humorous description."—*Capital and Labour*, May 17th, 1876.



REVIEWS OF CAPT. HAYES'
RIDING: ON THE FLAT AND ACROSS COUNTRY.
A Guide to Practical Horsemanship.

Illustrated Sporting and Dramatic News—"The book is one that no man who has ever sat in a saddle can fail to read with interest."

The Field.—"The general directions are in most cases in accordance with our own opinions; and Mr. Hayes has supplemented his own experience of race-riding by resorting to Tom Cannon, Fordham, and other well-known jockeys for illustration. 'The Guide' is, on the whole, thoroughly reliable; and both the illustrations and the printing do credit to the publishers."

The Sporting Life.—"It has, however, been reserved for Captain Hayes to write what in our opinion will be generally accepted as the most comprehensive, enlightened, and 'all round' work on riding, bringing to bear as he does not only his own great experience, but the advice and practice of many of the best recognized horsemen of the period."

Athenæum.—"Is an eminently sensible and useful manual. . . . Is in all respects satisfactory."

Scotsman.—"A thoroughly practical treatise."

Graphic.—"Is as practical as Captain Horace Hayes' 'Veterinary Notes' and 'Guide to Horse Manage-

ment in India.' Greater praise than this it is impossible to give."

Vanity Fair.—"Three-fourths of those who indulge in what they call riding in Hyde Park would do well to learn this book by heart."


The Queen.—"The chapter devoted to 'ladies' riding' calls for notice in these columns, as in it will be found information of a kind which is frequently asked for, while it is not always easy to point out to the enquirer a trustworthy source from which it may be obtained."

Society.—"The whole book is written in a style eminently suited to the subject."


Bell's Life.—"There is left nothing unsaid in the present publication to ensure a thorough acquaintance with the subject."

Sporting Times.—" 'Riding,' we venture to say, will take its place as a manual for equestrians, and will be equally appreciated by the boy with his first Shetland pony and the practised sportsman of the shires. . . . It is written in an easy, pleasant style; not too elaborate for the youthful rider, and sufficiently instructive for the practical horseman. We heartily commend it to our readers."

The Standard.—"Captain Hayes having written an able work on the horse and how to treat him, goes on by a course of natural progress to describe how to ride him, and displays in his last book the ability which characterised his previous effort. To possess knowledge and to succeed in imparting it are two different things; but Captain Hayes is not only master of his subject, but he knows how to aid others in gaining such a mastery as may be obtained by the study of a book."



HINDU MYTHOLOGY.



By Rev. W. J. WILKINS.

THACKER, SPINK & CO., CALCUTTA.

JUST PUBLISHED.

Uniform with 'Lays of Ind.," "Riding," "Anglo-Indian Domestic Life," &c.
Imperial 16mo. Rs. 7.

HINDU MYTHOLOGY:

VEDIC AND PURANIC.



BY REV. W. J. WILKINS,

OF THE LONDON MISSIONARY SOCIETY, CALCUTTA.

*ILLUSTRATED BY VERY NUMEROUS ENGRAVINGS FROM
DRAWINGS BY NATIVE ARTISTS AND OTHERS.*

THACKER, SPINK & CO., CALCUTTA.

IN PREPARATION.

Uniform with "Lays of Ind," "Riding," "Hindu Mythology," &c.

A NATURAL HISTORY
OF THE
MAMMALIA OF INDIA.



By R. A. STERNDALÉ,

AUTHOR OF "SCENES OR CAMP LIFE ON THE SATPURA RANGE," "THE DENIZENS OF THE JUNGLE," "THE AFGHAN KNIFE," ETC.

COPIOUSLY ILLUSTRATED FROM ORIGINAL DRAWINGS
BY THE AUTHOR AND OTHERS.



IN PREPARATION.

Uniform with "Lays of Ind," "Riding," "Hindu Mythology," &c

ANGLO-INDIAN DOMESTIC LIFE.

BY THE AUTHOR OF "RURAL LIFE IN BENGAL."


PROFUSELY ILLUSTRATED.



A New Edition abridged and re-written with the Original and other Illustrations
selected by the late COLLESWORTHY GRANT.



HANDBOOK OF INDIAN FERNS.



By COLONEL R. H. BEDDOME.

THACKER, SPINK & CO., CALCUTTA.

IN THE PRESS.

Uniform with "Lays of Ind," "Natural History of the Mammalia of India,"
"Hindu Mythology," "Riding," &c.


A POPULAR HANDBOOK
OF
INDIAN FERNS.

By COLONEL R. H. BEDDOME,

AUTHOR OF "THE FERNS OF BRITISH INDIA," "THE FERNS OF SOUTHERN INDIA."

The work has been specially written in a style as free from technicalities as possible, to meet the wants of non-scientific readers, and to assist visitors to the numerous Hill Stations of India in the selection and gathering of Ferns.

Illustrated with 300 Plates from the Drawings already so well known in India.



ILLUSTRATED FROM DRAWINGS BY THE AUTHOR.

THACKER, SPINK & CO., CALCUTTA.

Second Edition, in Crown 8vo., Illustrated, Rs. 7.

VETERINARY NOTES FOR HORSE-OWNERS.

AN EVERY-DAY HORSE BOOK.

BY CAPT. M. HORACE HAYES.

OPINIONS OF THE PRESS.


Saturday Review.—"The work is written in a clear and practical way."

The Field.—"Of the many popular veterinary books which have come under our notice, this is certainly one of the most scientific and reliable. The author tells us, in the preface to the first edition, that any merit which the book may possess is to

be ascribed to the teaching of the Principal and Professors of the New Veterinary College at Edinburgh, where he studied. It is much to be desired that every student would make so much use of his opportunities as Capt. Hayes has done.

"Some notice is accorded to nearly all the diseases which are common to horses in this country, and the writer takes advantage of his Indian experience to touch upon several maladies of horses in that country, where veterinary surgeons are few and far between. The description of symptoms and the directions for the application of remedies are given in perfectly plain terms, which the tyro will find no difficulty in comprehending; and, for the purpose of further smoothing his path, a chapter is given


on veterinary medicines, their actions, uses, and doses. This information will be most acceptable to the majority of horse-owners, and may be invaluable on an emergency when no advice better than that of the village cow doctor can be obtained."



THACKER, SPINK & CO., CALCUTTA.

The Veterinary Journal.—"A handy little book for the use of horse-owners, which may prove of much service to them when they cannot obtain the assistance of a veterinary surgeon, as well as afford them some notion of many of the ailments to which the horse is exposed, and the manner in which they may be best treated."

The Sporting Times.—"It is what it professes to be—a clear and comprehensive manual for all horse-owners, and, without trenching too much upon the almost



sacred mysteries of the veterinary art, it places a valuable weapon of defence in the hands of those who may be out of the reach of professional assistance, and we should be glad to see it as a part of the 'properties' wherever a horse is kept."

Illustrated Sporting and Dramatic News.—"Simplicity is one of the most commendable features in the book. What Captain Hayes has to say he says in plain terms, and the book is a very useful one for everybody who is concerned with horses."

The Sporting Life.—"We heartily welcome the second edition of this exceedingly useful book. The first edition was brought out about two years since, but the work now under notice is fully double the size of its predecessor, and as a matter of course contains more information. Captain Hayes, the author, is not only a practical man in all things connected with the horse, but has also studied his subject from a scientific point of view."

THACKER, SPINK & CO., CALCUTTA.

New Edition. Crown 8vo., price Rs. 5.

A GUIDE TO TRAINING AND HORSE MANAGEMENT IN INDIA.

BY CAPT. M. HORACE HAYES.

OPINIONS OF THE PRESS.

The Pioneer (Allahabad).—"The merits of Captain Hayes' book are so well known to all Anglo-Indians that it would be idle to go over old ground in praising the sound practical knowledge shown by the author in all he writes concerning race-horses. The new edition, however, goes even farther than this, for the subject of horse management is no longer made subservient to training, but is dealt with fully and clearly in separate chapters."

Oriental Sporting Magazine.—"His intentions are truly sportsmanlike and praiseworthy, and if carefully followed ought to ensure the attainment of a portion of that success in training which the author himself has been renowned for."

Englishman (Calcutta).—"It is the practical part of the book with reference to the more ordinary matters connected with horses that makes it so useful."

The Veterinary Journal.—"We entertain a very high opinion of Captain Hayes' book on 'Horse Training and Management in India,' and are of opinion that no better guide could be placed in the hands of either amateur horseman or veterinary surgeon newly arrived in that important division of our empire. We had not the opportunity of seeing the first edition, but from a careful perusal of the present it appears to contain everything pertaining to training and managing horses. The author proves himself to be not only

THACKER, SPINK & CO., CALCUTTA.

well-informed theoretically and practically, as to preparing horses for racing in that hot climate, as well as to their equipment and handling for ordinary work, but he also shows that he is possessed of no mean amount of that knowledge which is derived from the study of veterinary medicine. His remarks on shoeing are most judicious and sensible, and if his recommendations were adopted and carried into effect, the pernicious farriery of the native shoeing-smiths would be replaced by a method which would greatly benefit horses and their owners.

“Everything relating to racing in India, and the rules of the turf in that country, is embodied in Captain Hayes’ book, and not the least valuable portion of it is the Hindustanee vocabulary, containing vernacular terms referring to all matters relating to the horse, as well as diseases, medicine, etc. This alone should make the guide valuable to the young veterinary surgeon, who finds himself compelled to pick up, as best he can, the rudiments of a strange language when he commences his tour of professional duty.”

Saturday Review.—“Captain Hayes has brought out a second edition of his work on the training and management of horses in India. It is, of course, specially adapted to the circumstances of that country; but the general instructions which it contains, and which are of a shrewd and practical character, render it a useful guide in regard to horses anywhere.”

Bombay Gazette.—“We have to acknowledge with thanks the receipt of the new edition, re-arranged and much enlarged, of Captain Hayes’ very valuable handbook on the above subject. Captain Hayes, by his ‘Veterinary Notes for Horse-owners,’ as well as the former edition of the work we are now noticing, has already made his name pretty familiar to the lovers of horseflesh in India, and it would therefore be quite superfluous for us to say that he is an authority to whom the horse-owner may safely turn for advice and assistance on every kind of subject connected with the horse and his use.”

BY THE SAME AUTHOR. IN THE PRESS.

INDIAN RACING REMINISCENCES.

REPRINTED, WITH ADDITIONS FROM THE INDIAN NEWSPAPERS,
WITH NUMEROUS ILLUSTRATIONS.

THACKER, SPINK & CO., CALCUTTA.

One Vol. Small 8vo. Rs. 3-8.

TALES FROM INDIAN HISTORY:

BEING THE

Annals of India retold in Narratives.

BY J. TALBOYS WHEELER.

Forms a complete History of India from the earliest period to the present day, drawn up as a series of "Narratives" for general reading in schools and families. So far it will resemble the history of Scotland as told in Sir Walter Scott's "Tales of a Grandfather," omitting all details and disquisitions which are sufficiently given in the author's larger histories of India, and seeking to tell the progress of events in the most simple and attractive manner.

Reviews of Wheeler's Tales from Indian History.

"While the work has been written for them (natives), it has also been written for the people of England, who will find in the volume, perhaps for the first time, the history of our great dependency made extremely attractive reading. Mr. Wheeler's narrative is written in a most graceful style: indeed, he is master of the English language. He does not confine himself to the mere dry details of history, but tells the adventures of Indian heroes and heroines in legends of love and war; describes the village communities of India, their organization and self-government; delineates the results of caste, infant marriage, and other Hindoo institutions and usages as seen in the family and social life of the people in villages and towns, as well as in courts and palaces. . . . The work also contains valuable observations on the foreign relations of the Indian Empire with Persia, Russia, Turkey, and China. Altogether this is a work of rare merit."—*Broad Arrow*.

"In going through an interesting book, the reader will be furnished with a good general notion of Indian history, and learn besides something about Indian modes of life."—*Queen*.

"Will absorb the attention of all who delight in thrilling records of adventure and daring. It is no mere compilation, but an earnest and brightly-written book."—*Daily Chronicle*.

"This little volume contains a history of India in the form of tales and narratives, intended by the author for the people of India as well as for those of the British Isles."—*Army and Navy Gazette*.

"No young reader who revolts at the ordinary history presented to him in his school books will hesitate to take up this. No one can read a volume such as this without being deeply interested."—*Scotsman*.

THACKER, SPINK & CO., CALCUTTA.

In Crown 8vo., Rs. 3-8.

BASIL ORMOND, AND CHRISTABEL'S LOVE.

Metrical Tales.

BY THE AUTHOR OF "LAYS OF IND."

OPINIONS OF THE PRESS.

Broad Arrow.—"Display remarkable power, coupled with melodious versification."

Graphic.—"The author of 'Lays of Ind' shows in his latest book, 'Basil Ormond, and Christabel's Love,' that he can write something more than good humorous verse—a fact which was indeed foretokened by more than one piece in his former volume, notably by 'The Loafer.' There is, perhaps, no one amongst our minor singers whose rhythm is more easy and natural, whilst in the first of the two poems there is an amount of tragic intensity for which we were hardly prepared."

Times of India.—"Both stories are prettily told, and we feel assured that these modest poems will command a large circle of readers. 'Aliph Cheem' is so well known to the Indian public through his 'Lays of Ind,' that anything from his pen requires but few words of recommendation from us; but in his present poems he has opened a new mine, and we are mistaken if the public do not agree with us in thinking that he has hit upon a vein of rich metal."

Statesman.—"We have already seen how 'Aliph Cheem' can use his pen for good-natured satire and racy fun, and are now presented with a couple of poems of a deeper vein. The plaintive strain in these prolonged ballads, for such the poems are, is maintained without a break from the first stanza to the last."

Islington News.—" 'Aliph Cheem,' the poet who received the commendation of the entire Press, both of England and India, upon his former work, has written the above volume, which comprises two love stories of an Idyllic yet striking character. . . . At the present time, when there is so much versification, which is not only inelegant but unintelligible, it is gratifying to have a book placed in our hands for review that is clear in style, and displays a genuine poetic instinct from the first page to the last."

THACKER, SPINK, & CO., CALCUTTA.

In 8vo., price Rs. 3-8.

THE
RACES OF AFGHANISTAN:

BEING A BRIEF ACCOUNT OF THE
*PRINCIPAL NATIVE TRIBES INHABITING
THAT COUNTRY.*

By **SURGEON-MAJOR H. W. BELLEW, C.S.I.,**
Late on Political Duty at Kabul.

OPINIONS OF THE PRESS.

The Examiner.—"If Lord Hartington wants to begin to understand the problem he has to deal with in Afghanistan, he should take down with him to the House of Commons this new book by Dr. Bellew. . . . Dr. Bellew perceived, by the ignorant comments of speakers and writers on the Afghan question, how very much some book was needed to enlighten the understandings of public instructors; and therefore, amid the multifarious duties he had to fulfil at Cabul, he set himself to produce a work which does not err on the side of elaborateness, but which yet contains ample information to keep journalists and Members of Parliament and platform orators from vain babblings on this intricate subject. . . . The *résumé* of the history of the Afghans, including a chapter summarising our relations with the country, leads naturally up to the story of the life of Shere Ali, which results in the conclusion that the Afghan 'is not fit to govern either himself or others, and sadly wants a master. If we don't take up that *rôle*, Russia will. For a master the Afghans want, and a master they must have sooner or later.' . . . An admirable index increases the value of the work."


Indian Mail.—"This short book is of considerable merit, and ought to seriously influence the judgment of those who have assumed the direction of England's affairs. Its value lies in the preciseness of the information given, its clear and methodical arrangement, and its trustworthiness, from the acknowledged ability and absolute knowledge of the subject possessed by the author. Dr. Bellew is not a political speculator, but a serious scholar, who has spent many years among and in connection with the people of Afghanistan; he speaks their languages with ease, is well read in the history and literature of the country, and enjoys the confidence of the tribesmen. . . . Dr. Bellew's book will be read with considerable interest, both as a valuable contribution to ethnography and as a guide in the maze in which politicians have lost themselves."

THACKER, SPINK & CO., CALCUTTA.

Prepared for the use of the Survey Department of India, and published under the authority of the Government of India.

Royal 8vo. Rs. 12.

A MANUAL OF SURVEYING FOR INDIA:



*DETAILING THE MODE OF
OPERATIONS ON THE
TRIGONOMETRICAL,*

TOPOGRAPHICAL,

AND REVENUE


SURVEYS OF INDIA.

COMPILED BY

SIR H. L. THUILLIER, K.C.S.I., ROYAL ARTILLERY,

AND

LIEUT.-COL. R. SMYTH, late BENGAL ARTILLERY.



THACKER, SPINK & CO., CALCUTTA.

In Crown 8vo., Rs. 3-8.

THE
STEEPLECHASE HORSE:

How to Select, Train and Ride Him.

**WITH NOTES ON ACCIDENTS AND DISEASES, AND
THEIR TREATMENT.**

BY CAPTAIN J. HUMFREY.

OPINIONS OF THE PRESS.

Sporting Life.—"Under this title we have to notice a very useful and interesting volume, evidently the work of a practical pen. The subject-matter applies principally to Indian experiences, but is for the most part applicable to the selection, education, and management of the horse in any country. The treatment of equine ailments is also set forth in plain language.

"With these remarks we feel justified in heartily commending the book, and its perusal cannot fail to repay."


Illustrated Sporting and Dramatic News.—"The following anecdote is an admirable specimen of what ingenuity and determination can accomplish. It is borrowed from an excellent little volume by Captain Humfrey, entitled, 'The Steeplechase,' a treatise which though primarily intended for Indian readers, may be studied with advantage by all who care to know what a practical judge has to say upon the subject."

THACKER, SPINK & CO., CALCUTTA.

Third Edition, Demy 8vo, Rs. 10.

A MANUAL OF GARDENING FOR BENGAL AND UPPER INDIA.

By THOMAS A. C. FIRMINGER, M.A.



OPERATIONS OF GARDENING.

Chap. I.—Climate—Soils—Manures.

Chap. II.—Laying-out a Garden—Lawns—Hedges—Hoeing and Digging—Drainage—Conservatories—Betel Houses—Decorations—Implements—Shades—Labels—Vermin—Weeds.

Chap. III.—Seeds—Seed Sowing—Pot Culture—Planting—Cuttings—Layers—Gootee—Grafting and Inarching—Budding—Pruning and Root Pruning—Conveyance.

Chap. IV.—Calendar of Operations.

GARDEN PLANTS.

1. Culinary Vegetables.
2. Dessert Fruits.
3. Edible Nuts.
4. Ornamental Annuals.
5. Ornamental Trees, Shrubs, and Herbaceous Perennials.

IN PREPARATION. Demy 8vo.

MANUAL OF AGRICULTURE FOR INDIA.

By J. FRED. POGSON,
AUTHOR OF "INDIAN GARDENING," "THE INDIAN AGRICULTURIST."

Principal Headings.—SOILS, PLOUGHING, MANURES, WHEAT, BARLEY, OATS, MILLET, MAIZE, PEA, BEAN, DALI, SUGAR, ROOT FIELD CROPS, &c., &c.

THACKER, SPINK & CO., CALCUTTA.

In Demy 8vo., cloth., Rs. 8.

MALARIA:

ITS CAUSES AND ITS EFFECTS;

AND

INJURIES OF THE SPLEEN.

BY E. G. RUSSELL, M.B., B.Sc., LOND.,

Surgeon, Bengal Medical Service.

OPINIONS OF THE PRESS.

The Lancet.—"In the present volume Dr. Russell discusses most exhaustively the subject of malaria and its effects, and the work should be read by all who are liable to be called upon to treat the conditions to which it gives rise. The chapter which describes the malarial lesions of the stomach and liver will perhaps be read with most interest by the English practitioner. Dr. Russell offers a useful hint not to mistake severe dyspepsia and gastric irritability associated with pale diarrhoea arising from too constant and prolonged a residence in malarious regions for hepatic disease induced by 'free living'—a caution the more needed since, we fear, all dyspeptic troubles occurring in old Indians are apt to be regarded as due 'to liver' provoked by unlimited brandy pawnee, hot curries, and tropical suns. Dr. Russell ably discusses the relationship between ague and typhoid—a question that has been raised by Sir Joseph Fayrer. . . . The second part of the volume is devoted to an analysis of fifty cases of injury of the spleen. These injuries, from the frequency of their occurrence, and the importance of their medico-legal bearings, are of peculiar interest to the medical practitioner in India, but their consideration need not detain us at present. We therefore leave Dr. Russell's volume, commending it to the notice of our readers as a book full of facts and of clinical research, and which must be considered as an authority on the subject of malaria."

The Medical Press and Circular.—"The appearance of this work is opportune at this time, when the subject of fevers in relation to climate in India is obtaining special attention. The author, from his position as civil surgeon of F'amrup in Assam, had ample opportunities of practically studying his subject, for the district of which that station is the centre is noted for its malarious character, and for the prevalence of malarial disease, including 'fever and its sequelæ, dysentery, diarrhoea, and cholera.' But besides the results of his personal observations, he summarises the views of recognised authorities on his subject, and thus presents to his readers a convenient epitome of previous investigations. His work is of considerable merit, and deserves to be extensively read."

THACKER, SPINK & CO., CALCUTTA.

ENTIRELY RE-WRITTEN. Seventh Edition. Crown 8vo., Rs. 7.

GOODEVE'S HINTS
FOR THE
MANAGEMENT AND MEDICAL TREATMENT
OF
CHILDREN IN INDIA.

RE-WRITTEN BY EDWARD A. BIRCH, M.D.,

Surgeon-Major, Bengal Establishment.

Dr. Goodeve.—"I have no hesitation in saying, that the present edition is for many reasons superior to its predecessors. It is written very carefully, and with much knowledge and experience on the author's part, whilst it possesses the great advantage of bringing up the subject to the present level of Medical Science."

The Medical Times and Gazette, in an article upon this work and Moore's 'Family Medicine for India,' says:—"New editions of these two well-known works have recently appeared. They are both intended to supply in some measure the medical wants of our numerous countrymen in India, who may be either far from professional help, in emergencies of sickness or of accident, or destitute of medical advice regarding the proper management of their own health, and especially that of their children, in the trying climate of Hindostan. Although we are, as a rule, very much opposed to popular medical instruction, believing that the result is most frequently a minimum of serviceable knowledge along with a vast preponderance of what is but partial, misleading, and dangerous, yet the peculiar circumstances of many of our countrymen in India, together with the special and insidious dangers of its varying climate, fully justify the publication of a few trustworthy popular works to warn the unwary new-comer, before it be too late, of the dangers he has to encounter, and to give judicious counsel to solitary individuals and families who cannot enjoy the advantages of personal professional advice. Moreover, the two works before us are in themselves probably about the best examples of medical works written for non-professional readers. The style of each is simple, and as free as possible from technical expressions. The modes of treatment recommended are generally those most likely to yield good results in the hands of laymen; and throughout each volume the important fact is kept constantly before the mind of the reader, that the volume he is using is but a poor substitute for personal professional advice, for which it must be discarded whenever there is the opportunity. Written with such objects, and in such a spirit, these volumes cannot fail to be of the greatest service; and that they are appreciated is shown by the rapid appearance of successive editions, the second mentioned and elder treatise having now reached the seventh edition. We would add, that although they are specially written for lay readers, there are few young medical officers proceeding to India who would not receive several useful hints from these unpretentious volumes. But it is to parents or to the guardians of European children in India that they must be of pre-eminent service."

THACKER, SPINK & CO., CALCUTTA.

PART I., Rs. 1. PART II., Rs. 2.

A TEXT BOOK
OF
INDIAN BOTANY:
MORPHOLOGICAL,
PHYSIOLOGICAL,
and SYSTEMATIC.



BY **W. H. GREGG,**
LECTURER ON BOTANY, HUGHLI
COLLEGE.

Revised by
W. BOTTING HEMSLEY,
THE HERBARIUM, KEW.



Crown 8vo, cloth, Rs. 2-8.



THE AMATEUR GARDENER IN THE HILLS

HINTS FROM VARIOUS AUTHORITIES ON
GARDEN MANAGEMENT

And adapted to the Hills; with Hints on
FOWLS, PIGEONS, AND RABBIT KEEPING;

And various Recipes connected with the above subjects
which are not commonly found in Recipe Books.



THACKER, SPINK & CO., CALCUTTA.




In Super Royal 8vo, with 166 Illustrations. Rs. 6-8.

RURAL LIFE IN BENGAL.

Illustrative of Anglo-Indian Suburban Life; the Habits of the Rural Classes; the varied Produce of the Soil and Seasons; and the Culture and Manufacture of Indigo.

By C. GRANT.



Crown 8vo, Rs. 3.

THE INDIAN COOKERY BOOK



**A PRACTICAL HANDBOOK TO THE
KITCHEN IN INDIA:**

ADAPTED TO THE THREE PRESIDENCIES.

Containing Original and Approved Recipes in every department of Indian Cookery; Recipes for Summer Beverages and Home-made Liqueurs; Medicinal and other Recipes; together with a variety of things worth knowing.

BY A THIRTY-FIVE YEARS' RESIDENT.

THACKER, SPINK & CO., CALCUTTA.

GUIDE BOOKS, &c.

AGRA.

A Handbook for Visitors to Agra and its Neighbourhood. By H. G. KEENE, C.S. Fourth Edition, revised. Maps, Plans, &c. Fcap. 8vo., cloth. Rs. 2-8.

ALLAHABAD—CAWNPORE.

A Handbook for Visitors to Allahabad, Cawnpore, and Lucknow, By H. G. KEENE, C.S. Second Edition, re-written and enlarged. Fcap. 8vo. Rs. 2-8.

CENTRAL INDIA.

A Memoir of Central India, including Malwa and adjoining Provinces, with the History, and copious Illustrations, of the Past and Present Conditions of that Country. By Major-General Sir JOHN MALCOLM, G.C.B., K.L.S. Reprinted from the Third Edition. With Map. In 2 vols., crown 8vo. Rs. 7.

DELHI.

A Handbook for Visitors to Delhi and its Neighbourhood. By H. G. KEENE, C.S. Fourth Edition. Maps. Fcap. 8vo., cloth. Rs. 2-8.

KASHMIR.

The Kashmir Handbook: A Guide for Visitors. With Map and Routes. By JOHN INCE, M.D. Third Edition. Rs. 5.

NORTH-WEST PROVINCES.

How we did "the Lions" of the North-West: A Trip in the Durga-Pujahs to Lucknow, Delhi, Agra. By F. O. B. Re. 1.

HOMEWARD, VIA AMERICA.

Calcutta to Liverpool, by China, Japan, and America, in 1877. By Lieut.-General Sir HENRY NORMAN. Second Edition. Fcap. 8vo., cloth. Rs. 2-8.

THE HIMALAYA.

Three Months' Tour through Bussahir, Kunawar, and Spiti, to Lahoul. By Mrs. J. C. MURRAY AYNLEY. Crown 8vo.

MASURI.

A Guide to Masuri Dehra, Doon, and the Hills to the North, including Chapters on Hardwar, Roorkee, &c. By J. NORTHAM.

MEDICINE.

Municipal Work in India ; or, Hints on Sanitation, General Conservancy, and Improvements in Municipalities, Towns, and Villages. By R. CRAUFUIRD STERNDALÉ, Vice-Chairman Calcutta Suburban Municipality. Crown 8vo. Rs. 5.

Blindness. Causes of Blindness in India ; Treatment and Cure. By G. C. HALL, Esq., Surgeon Ind. Medl. Service. Fcap. 8vo, cloth. Rs. 2.

Malaria : its Cause and Effects. Injuries of the Spleen : an Analysis of Thirty-nine Cases. By E. G. RUSSELL, M.B. 8vo, cloth. Rs. 8.

“ Must be considered an authority on the subject.”—*Lancet*.

Jail Hygiene in India. By T. HUME, M.B. Fcap. 8vo. Rs. 2-8.

The Himalaya : Climate and Medical Topography, in their relation to the Disease Distribution of the Himalaya and Sub-Himalayan Districts ; with reason for assigning a Malarious Origin to Goitre and other Diseases. By F. N. MACNAMARA, M.D., F.R.G.S. With Map. 8vo. Rs. 10.

The Materia Medica of the Hindus. Compiled from Sanscrit works by UDOY CHAND DUTT ; with a Glossary of Indian Plants. By G. KING, F.L.S., M.B. 8vo, cloth. Rs. 6.

A Monograph of Two Hundred and Forty-eight Cases of Lateral Lithotomy Operations performed in the North-West Provinces during a period of twelve years, by RAM NARAIN DASS. Illustrated by 256 Diagrams of Calculi, showing exterior and section, executed in Lithography. Pp. viii., 94. 8vo., cloth. Rs. 10.

A Glossary of Medical and Medico-Legal Terms, including those most frequently met with in the Law Courts. Compiled by R. F. HUTCHINSON, M.D., Surgeon-Major, Bengal Army. Fcap. 8vo., cloth. Rs. 3.

Hidayat-ul-Hukuma. A Hindustani Manual for the use of Indian Medical Officers, Native Doctors, and others. By GEORGE S. A. RANKING, B.A., M.B., Cantab., Surgeon, 6th Bengal Cavalry. 18mo., sewed. Rs. 1-4.

DOMESTIC, GARDEN, AND PRODUCE.

Goodeve's Hints for the Management and Medical Treatment of Children in India. By EDWARD A. BIRCH, M.D., Surgeon-Major. Seventh Edition. Crown 8vo., cloth. Rs. 7.

"I have no hesitation in saying that the present one is for many reasons superior to its predecessors. It is written very carefully, and with much knowledge and experience on the author's part, whilst it possesses the great advantage of bringing up the subject to the present level of Medical Science."—Dr. GOODEVE.

Indian Notes about Dogs; their Diseases and Treatment. Compiled by Major C.—, author of "Horse Notes." Fcap. 8vo., cloth. Re. 1-8.

Indian Domestic Economy and Receipt Book. With Hindustani Romanized Names; comprising numerous directions for Plain Wholesome Cookery, both Oriental and English; with much miscellaneous matter answering all general purposes of reference connected with Household Affairs likely to be immediately required by Families, Messes, and Private Individuals residing at the Presidencies or Out-Stations. By Dr. R. RIDDELL. Eighth Edition, revised. Royal 12mo. Rs. 6.

Indian Cookery Book: A Practical Handbook to the Kitchen in India, adapted to the three Presidencies. By A THIRTY-FIVE YEARS' RESIDENT. Crown 8vo. Rs. 3.

Queries at a Mess Table: What shall we Eat? What shall we Drink? By JOSHUA DUKE, Surgeon, 3rd Punjab Cavalry, Author of "Banting in India." Fcap. 8vo., cloth, gilt. Rs. 2-4.

How to get Thin; or, Banting in India. By J. DUKE. 12mo., stiff cover. Re. 1.

A Manual of Gardening for Bengal and Upper India. By T. A. C. FIRMINGER, Honorary Member Agri.-Horti. Society. Third edition. 8vo., cloth, gilt. Rs. 10.

The Amateur Gardener in the Hills: being Hints and Recipes. By AN AMATEUR. Crown 8vo. Rs. 3.

Familiar Indian Flowers. Thirty Coloured Plates and descriptive Letterpress. By LENA LOWIS. 4to., cloth. Rs. 22-8.

Tea Gardens, Coffee Plantations, Indigo, Jute, Lac, and Cinchona Concerns in India, with their Capital, Directors, Proprietors, Agents, Managers, &c., and their Factory Marks. Rs. 2-8.

INDIAN LAW.

The Negotiable Instruments Act of 1881. Edited, with Introduction, Notes, &c., by M. D. CHALMERS, Esq., Barrister-at-Law, Author of "Digest of the Law of Bills of Exchange." 8vo., cloth. Rs. 7.

The Law of Specific Relief in India : being a Commentary on Act I. of 1877. By C. COLLETT, M.C.S., Author of "The Law of Torts." 8vo. (*In the Press*).

The Indian Law Examination Manual. Second Edition. Revised, with all the New Laws, by FENDALL CURRIE. 8vo. Rs. 5.

Revenue Sale Law of Lower Bengal. The Acts, Regulations, and Rules of the Board of Revenue; with Notes. Edited by W. E. H. FORSYTH. 8vo. Rs. 5.

The Land Acquisition Act. Act X. of 1870, and Supplementary Acts. Edited, with Notes, by H. BEVERLEY, B.C.S. 8vo. Rs. 5.

The Law of Succession in India : Testamentary and Intestate. By G. S. HENDERSON, Barrister-at-Law. 8vo.

The Probate and Administration Act. Act V. of 1881. With Notes by W. E. H. FORSYTH. Edited by F. J. COLLINSON. Rs. 5.

Indian Case Law on Torts. By R. D. ALEXANDER, B.C.S. Crown 8vo. Rs. 4.

HISTORY, &c.

The Sovereign Princes and Chiefs of Central India. By G. R. ABERIGH MACKAY. Vol. I. Introduction. HOLKAR. Illustrated. Royal 8vo., cloth, gilt. Rs. 12.

Ancient India, as described by MEGASTHENES and ARRIAN. With Introduction, Notes, &c., by J. W. MCCRINDLE. 8vo. Rs. 2-8.

The Commerce and Navigation of the Erythræan Sea : being a Translation of PERIPLUS, MARIS, ERYTHRÆI, and ARRIAN'S VOYAGE OF NEARCHOS. With Introduction, Commentary, and Notes, By J. W. MCCRINDLE. 8vo. Rs. 3.

Hindu Tribes and Castes. By Rev. M. A. SHERRING. Vol. II. : Hindu and Mahommedan Tribes of North-West Provinces, and the Aborigines of the Central Provinces. Vol. III. : The Races of South India, and Dissertations on Caste, Unity of the Human Race, &c. 4to. Each Vol., Rs. 16.

Early Records of British India. A History of Early English Settlements, as told in old Records, &c. By J. TALBOYS WHEELER. Royal 8vo. Rs. 5.

THACKER, SPINK & CO.,

CALCUTTA.

*THE VARIOUS DEPARTMENTS OF THE ESTABLISHMENT
ARE AS FOLLOW:*

Bookselling.

The New Publications received by every steamer, and a large stock of Books in every department of Literature kept on hand.

Publishing.

The present Catalogue is an illustration of some of the Works published by them.

Stationery.

Plain, for offices and general use; Fancy, for presents and objects of beauty.

Printing and Engraving.

In all branches, including Books, Pamphlets, and General Letterpress; Heraldic Engraving, Die-Sinking, and Copperplate Engraving.

Miscellanies.

Scientific Instruments, Artists' Materials, Cutlery, Dressing Cases, Writing Cases, Despatch Boxes, and Fancy Goods in great variety.

Banking and Agency.

Demand and Fixed Deposit Accounts opened; Home Remittances granted; House and Personal Agency undertaken.

5 & 6 GOVERNMENT PLACE, CALCUTTA.



